

One-Pot Synthetic Approach to Dipyrrmethanes and Bis(indolyl)methanes via Nitrosoalkene Chemistry

Ana L. Cardoso, Susana M. M. Lopes, Carla Grosso, Marta Pineiro, Américo Lemos, and Teresa M. V. D. Pinho e Melo*

Cite This: *J. Chem. Educ.* 2021, 98, 2661–2666

Read Online

ACCESS |


Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: A one-pot regio- and stereoselective synthesis of a dipyrrmethane and a bis(indolyl)methane based on two consecutive reactions of nitrosoalkenes with pyrrole or indole, respectively, is described as an experiment to be carried out by upper-division undergraduate students in a laboratory classroom. Importantly, the ability of electrophilic conjugated nitrosoalkenes to react via Michael addition or hetero-Diels–Alder reactions with electron-rich heterocycles will provide an opportunity for students to acknowledge alternative reaction pathways underlying certain transformations. Reactions were performed under mild conditions using water as a solvent, followed by purification through column chromatography on silica gel, and characterization of the desired products by NMR and IR spectroscopy. This laboratory experiment combines organic synthesis, determination of the purity of compounds (TLC analysis and melting point measurements), as well as structural analysis (interpretation of 1D NMR spectra). Several important organic chemistry concepts, such as stereo- and regioselectivity, *in situ* generation and reactivity of conjugated nitrosoalkenes, conjugated 1,4-addition reactions, and cycloaddition reactions, are also discussed.


KEYWORDS: Upper-Division Undergraduate, Organic Chemistry, Hands-On Learning/Manipulatives, Heterocycles, NMR Spectroscopy, IR Spectroscopy



In past decades, the chemistry of conjugated nitrosoalkenes has been extensively explored in organic synthesis, originating a plethora of new heterocyclic and nonheterocyclic compounds, some of which exhibit promising biological activities.¹ These reactive intermediates are usually generated *in situ* through base-promoted dehydrohalogenation of α -halooximes. The conjugation of the nitroso group with the C=C double bond conveys a strong electrophilic character to the system making the carbon C-4 strongly activated to nucleophilic attack. Thus, they can participate in conjugate 1,4-additions (Michael additions) with a variety of nucleophiles (Scheme 1).² On the other hand, they can also act as electron-deficient heterodienes participating efficiently in inverse electron-demand Diels–Alder reactions with electron-rich olefins and electron-rich heterocycles to afford 1,2-oxazines (Scheme 1).¹

In the reaction of nitrosoalkenes with electron-rich heterocycles, both reactivities have been observed, with the mechanistic pathway being strongly influenced not only by the nitrosoalkene substituents but also by the type of heterocycle. 1*H*-Imidazole reacts with ethyl bromopyruvate oxime **1a** via an elimination reaction to generate nitrosoalkene **2a** followed by a conjugate addition reaction to give the corresponding alkylated product **3** (Scheme 2a).³ On the other hand, the inverse electron-demanding Diels–Alder reaction of

Scheme 1. Typical Reactivity of Nitrosoalkenes



nitrosoalkenes **2a** and **2b**, generated *in situ* from the corresponding oximes **1**, with pyrrole and indole, gives open-chain oximes **5** and **7**, respectively, as single products. The synthesis of these products can be rationalized considering the


Received: February 23, 2021

Revised: June 9, 2021

Published: July 23, 2021



Scheme 2. Conjugate 1,4-Addition and Hetero-Diels-Alder Reaction of Nitrosoalkenes with Heterocycles



formation of the Diels–Alder cycloadducts, the bicyclic 1,2-oxazines **4** and **6**, followed by oxazine ring-opening with concomitant rearomatization of the pyrrole or indole unit (Scheme 2b). In fact, the open-chain oximes were isolated as single isomers, with the hydroxyl and pyrrolylmethyl or indolylmethyl groups *cis* as expected for a mechanism pathway involving a hetero-Diels–Alder reaction and subsequent ring-opening reaction of the bicyclic 1,2-oxazine. In addition, pyrrole was functionalized at position 2 whereas the reaction with indole gave a 3-alkylated derivative which is in agreement with the expected and opposite regioselectivity of the hetero-Diels–Alder reaction.^{4–8}

Conversely, the reaction of *p*-bromophenyl oxime **8** with pyrrole, under the same reaction conditions, led to the formation of two isomeric oximes **9** and **10** (Scheme 3a).⁹ This outcome is not consistent with a process involving a hetero-Diels–Alder reaction, which should lead exclusively to isomer **9**. Instead, a process involving as first step a conjugate addition reaction pathway occurs leading to two isomeric oximes **12** and

13. Starting from nitrosoalkene **11** at the *s-cis* conformation, oxime **12** was formed, whereas **11-s-trans** afforded oxime **13**. Oximes **12** and **13** undergo a 1,5-sigmatropic hydrogen shift, to give isomeric oximes **9** and **10**, respectively (Scheme 3b).⁹ Interestingly, it was demonstrated that indole reacts with the same nitrosoalkene **11** via an inverse electron-demanding Diels–Alder reaction giving only one open-chain oxime.¹⁰


Dipyrromethanes (DPs) are valuable and versatile building blocks of a vast number of compounds such as porphyrins, *meso*-substituted corroles, chlorins, calix[4]pyrroles, and BODIPY dyes.^{11,12} These compounds display remarkable photo- and biochemical properties suitable for several applications including photonic organic-based materials, optical anion sensors, photosensitizers in photodynamic therapy, and theranostic agents.^{13–15} On the other hand, naturally occurring and synthetic bis(indolyl)methanes (BIMs) have been shown to exhibit a wide range of biological activities and have also been used as dyes and colorimetric sensors.^{16,17}

The laboratory experiment described herein involves a regioselective one-pot synthetic approach to dipyrromethanes (e.g., **16**) and bis(indolyl)methanes (e.g., **17**) based on two consecutive reactions of nitrosoalkenes, generated *in situ* from 1,1-dichloro-2-propanone oxime (**14**), with pyrrole or indole, respectively (Scheme 4).^{18–21}


■ PEDAGOGICAL GOALS AND ASSESSMENT

The purpose of this laboratory experiment is to allow the students to carry out the synthesis of important scaffolds in organic synthesis in a straightforward procedure and under very mild conditions. The pedagogical aims of this experiment include deepening the knowledge about important organic chemistry concepts, such as regioselectivity and generation and reactivity of conjugated nitrosoalkenes, namely, conjugated 1,4-addition reactions and cycloaddition reactions. The students will also acquire knowledge in one-pot reactions and atom economy as well as on TLC analysis, chromatography techniques, and NMR and IR spectroscopy. Each experiment, the synthesis of DP **16** and BIM **17**, has been carried out successfully by 28 advanced undergraduate students (BSc Biochemistry and BSc Chemistry) within the organic chemistry course, working individually or in groups of two. After carrying out the experimental work and discussing the results, the students should be able to propose the

Scheme 3. Conjugate 1,4-Addition Reaction of 3-(*p*-Bromophenyl)nitrosoalkene with Pyrrole



Scheme 4. On-Water Synthesis of DP 16 and BIM 17 via Two Consecutive Reactions of Nitrosoalkenes with Pyrrole or Indole



mechanism for the synthesis of dipyrrromethanes and bis-(indolyl)methanes based on the chemistry of nitrosoalkenes. The students will also acquire knowledge on purification and isolation of organic compounds by chromatographic techniques and demonstrate an understanding of the NMR and IR data that allowed them to structurally characterize the products.

The students' assessment is performed on the basis of the laboratory reports as well as on instructor observation of the students' laboratory skill development, postlab questions focused on spectral analysis and reaction mechanisms, and exam questions (Supporting Information).

In general, the assessment of students confirmed that the pedagogic goals of the laboratory experiment were successfully achieved, well-illustrated by the results of the students' answers to the exam questions. In question 1, students acknowledged that the replacement of 1,1-dichloro-2-propanone oxime by 1,1-dibromo-2-propanone oxime, in the reaction with pyrrole, would lead to an enhancement of the reaction rate due to the presence of a better leaving group (Br). On the other hand, answers to questions 2 and 3 clearly show that the students not only understood the mechanism underlying the synthesis of dipyrrromethanes and bis(indolyl)methanes but also were able to apply the acquired knowledge to reactions with other dienophiles (e.g., furan, question 2) and heterodienes (e.g., azoalkenes, question 3). Moreover, the correct identification of the reaction products (cycloadduct vs open-chain oxime derivative) was accomplished by the students revealing knowledge regarding the regioselectivity of the proposed reactions as well as the aromaticity–nucleophilicity of the heterocycles (furan vs pyrrole). Also, students were capable of identifying the cycloadduct arising from the reaction of ethyl bromopyruvate oxime with 2,5-dimethylpyrrole, where the ring-opening reaction pathway to yield the open-chain oxime derivative is precluded, thus demonstrating mechanistic knowledge on the topic (question 4).

Although there are several experiments reported in this *Journal* illustrating basic concepts of Diels–Alder reactions, undergraduate laboratory experiments on the application of the

hetero-Diels–Alder reaction are more limited.^{21–27} Moreover, this synthetic methodology features several aspects that make it suitable for an undergraduate laboratory experiment. The total synthesis involves four consecutive reactions carried out in one-pot, base-induced dehydrohalogenations, the hetero-Diels–Alder and Michael addition reactions, without the need of isolating or purifying any intermediate. The reactions are carried out at atmospheric pressure and room temperature using water as solvent, an environmentally friendly solvent. Furthermore, it illustrates the efficiency of hetero-Diels–Alder reactions carried out in an aqueous medium. This has been suggested to be caused not only by the hydrophobic effect in which an entropy-driven aggregation of two nonpolar molecules dissolved in water occurs, but also by hydrogen-bonding interactions. In addition, it has been observed that aqueous solvents can enhance the selectivity and regioselectivity of the reaction.²⁸ Indeed, the optimized on-water reaction conditions of these reactions afford the desired dipyrrromethane and bis(indolyl)methane in higher yields with significantly shorter reaction times than those carried out in dichloromethane or in the absence of solvent.^{18–20} Moreover, the synthetic methodology provides functionalized heterocyclic compounds in a regio- and stereoselective fashion.

EXPERIMENT OVERVIEW

In the classroom environment, upper-division undergraduate students may work either individually or in groups of two during three laboratory sessions of 3 h. Each group carries out one reaction sequence, either using pyrrole or indole, to obtain dipyrrromethane **16** or bis(indolyl)methane **17**, respectively (Scheme 4). Detailed experimental procedures are provided in the Supporting Information.

First Session

The first session involves the synthesis and isolation of dipyrrromethane **16** or bis(indolyl)methane **17** through the *in situ* generation of 4-chloronitrosoalkene **15** and subsequent reaction with pyrrole or indole, respectively. To prepare dipyrrromethane **16**, a solution of oxime **14** in pyrrole is added to a solution of sodium carbonate in water. The reaction mixture should be vigorously stirred at room temperature for 20 min. In the synthesis of bis(indolyl)methane **17**, the indole and a solution of oxime **14** in a cosolvent (dichloromethane) are added to the solution of sodium carbonate in water. The reaction is stirred vigorously for 1 h at room temperature. Upon completion, the biphasic mixture is transferred to a separator funnel and extracted with ethyl acetate, and the combined organic extracts are dried. After removal of the solvent, the composition of the crude product is evaluated by thin layer chromatography (TLC) analysis, and the appropriate eluent to be used in the following purification session is selected. The crude mixture should be stored in the refrigerator until the purification session.

Second Session

This session involves the purification of dipyrrromethane **16** or bis(indolyl)methane **17**. The flash chromatography is prepared using silica gel and the eluent selected in the first session. After isolation of the product, the solvent is evaporated off affording dipyrrromethane **16** as a light gray solid and bis(indolyl)methane **17** as an oil. However, compound **17** becomes a colorless solid upon addition of diethyl ether. The success of the isolation of the target compounds is confirmed by TLC




Figure 1. Representative parts of ^1H NMR spectrum of dipyrromethane **16**.




Figure 2. Representative parts of ^1H NMR spectrum of bis(indolyl)methane **17**.

analysis, and the compounds are stored in a desiccator until the next session.

Third Session

In this session, the yields and atom economy are calculated, and samples are prepared to record the IR and NMR spectra. Additionally, melting point measurements of the products can be carried out (optional).

HAZARDS

Students should be familiar with good laboratory practices to work safely in the laboratory. Personal protective equipment (PPE) is mandatory throughout the laboratory sessions. The entire experiment should be conducted in a fume hood. The experiments must be fully risk-assessed by all students. Proper procedures and precautions in handling waste and chemicals should be adhered to with reference to safety data sheets (SDS). All of the reactants and solvents used in the experiments should be carefully handled in a fume hood. Pyrrole [CAS 109-97-7], ethyl acetate [CAS 141-78-6],

dichloromethane [CAS 75-09-2], and diethyl ether [CAS 60-29-7] are flammable liquid and vapor, are toxic if swallowed, cause eye irritation or damage, and are harmful if inhaled. Indole [CAS 120-72-9] is a white solid which is harmful if swallowed, is toxic in contact with skin, causes serious eye irritation, and is very toxic to aquatic life. Sodium carbonate [CAS 497-19-8] is a white solid which causes serious eye irritation. Hexane [CAS 110-54-3] causes neurotoxicity. Organic solvent waste should be disposed in appropriate waste collector containers. According to our experience, oxime **14** is unstable at room temperature and should be stored in the freezer. Dipyrromethane **16** and bis(indolyl)methane **17** should be stored in the dark.

RESULTS AND DISCUSSION

Synthesis

The synthesis of dipyrromethane **16** and bis(indolyl)methane **17** was reproduced by BSc students, individually or in groups of two, during three laboratory sessions (3 h/session). The

students obtained dipyrromethane **16** in yields ranging from 30% to 55% and bis(indolyl)methane **17** with yields ranging from 34% to 49%. The atom economy was calculated as 53% and 63% for the synthesis of **16** and **17**, respectively. The TLC analysis of the crude mixture of reaction with pyrrole showed the presence of unreacted pyrrole, dipyrromethane **16**, and some impurities (Figure S1, see SI). Flash chromatography performed with the eluent ethyl acetate/hexane (1:1) allowed the isolation of pure dipyrromethane **16** as a light gray solid. Using the same purification method, bis(indolyl)methane **17** was isolated as a white solid. After purification, the purity of the products was assessed by TLC and ^1H NMR analysis of both compounds and by melting point measurements (optional).

Spectral Analysis

The identification and purity of the products was confirmed by the ^1H NMR spectra, by comparison with the previously reported data.^{18,19} The analysis of the ^1H NMR spectra allows the consolidation of concepts such as chemical shift and multiplicity. For example, in the ^1H NMR spectrum of dipyrromethane **16** (Figure 1), it is possible to identify the presence of signals corresponding to two pyrrole units. Two NH protons emerge at 10.55 ppm as a broad singlet and the OH proton at 10.48 ppm. Also, signals corresponding to CH protons of pyrrole units appear between 5.79 and 6.62 ppm while the signal of the *meso*-proton (H-5) shows a chemical shift of 4.88 ppm. Protons of the methyl group appear as a singlet at 1.70 ppm (Figure S6, see SI). The ^1H NMR spectrum of bis(indolyl)methane **17** (Figure 2) shows signals corresponding to two indole units, with the NH protons at 10.90 ppm and the OH proton at 10.43 ppm. Additionally, the CH protons of the indole units can be observed at 7.13 ppm (H-5 and H-6), as well as the *meso*-proton (H-1) at 5.40 ppm. The protons of the methyl group appear as a singlet at 1.80 ppm (Figure S11, see SI).

CONCLUSIONS

The ability of electrophilic conjugated nitrosoalkenes to participate in Michael addition reactions or hetero-Diels–Alder reactions with electron-rich heterocycles is illustrated and demonstrated in these laboratory experiments. The *in situ* generation and reactivity of conjugated nitrosoalkenes toward pyrrole or indole afforded the corresponding dipyrromethane and bis(indolyl)methane, respectively, in a stereo- and regioselective fashion.

These reactions carried out under mild conditions use water as solvent, have short reaction times, and involve simple manipulations, easy isolation, and handling procedures. The laboratory experiment includes the purification of the compounds through column chromatography, determination of their purity, and structural analysis using NMR spectroscopy.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available at <https://pubs.acs.org/doi/10.1021/acs.jchemed.1c00184>.

Detailed experimental procedures and instructions for students and spectra (IR, ^1H , and ^{13}C NMR spectroscopy) (PDF, DOCX)

AUTHOR INFORMATION

Corresponding Author

Teresa M. V. D. Pinho e Melo – University of Coimbra, Coimbra Chemistry Centre (CQC) and Department of Chemistry, 3004-535 Coimbra, Portugal; orcid.org/0000-0003-3256-4954; Email: tmelo@ci.uc.pt

Authors

Ana L. Cardoso – University of Coimbra, Coimbra Chemistry Centre (CQC) and Department of Chemistry, 3004-535 Coimbra, Portugal; orcid.org/0000-0003-0551-7255

Susana M. M. Lopes – University of Coimbra, Coimbra Chemistry Centre (CQC) and Department of Chemistry, 3004-535 Coimbra, Portugal; orcid.org/0000-0002-1580-5667

Carla Grosso – University of Coimbra, Coimbra Chemistry Centre (CQC) and Department of Chemistry, 3004-535 Coimbra, Portugal; orcid.org/0000-0003-1999-3471

Marta Pineiro – University of Coimbra, Coimbra Chemistry Centre (CQC) and Department of Chemistry, 3004-535 Coimbra, Portugal; orcid.org/0000-0002-7460-3758

Américo Lemos – CQC and Departamento de Química e Farmácia, Faculdade de Ciências e Tecnologia, Universidade do Algarve, 8005-139 Faro, Portugal; orcid.org/0000-0001-9588-4555

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.jchemed.1c00184>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Coimbra Chemistry Centre (CQC) is supported by the Portuguese Agency for Scientific Research, the Fundação para a Ciência e a Tecnologia (FCT), through project UIDB/00313/2020 and UIDP/00313/2020, cofunded by COMPETE2020-UE. C.G. also acknowledges FCT for the PhD fellowship SFRH/BD/130198/2017. The authors also acknowledge the UC-NMR facility for obtaining the NMR data (www.nmrcc.uc.pt).

REFERENCES

- (1) Lopes, S. M. M.; Cardoso, A. L.; Lemos, A.; Pinho e Melo, T. M. V. D. Recent advances in the chemistry of conjugated nitrosoalkenes and azoalkenes. *Chem. Rev.* **2018**, *118* (23), 11324–11352.
- (2) Boyko, Y. D.; Dorokhov, V. S.; Sukhorukov, A. Y.; Ioffe, S. L. Conjugated nitrosoalkenes as Michael acceptors in carbon-carbon bond forming reactions: a review and perspective. *Beilstein J. Org. Chem.* **2017**, *13*, 2214–2234.
- (3) Gilchrist, T. L.; Stretch, W.; Chrystal, E. J. T. Reaction of azoles with ethyl bromopyruvate oxime: alkylation by substitution and by elimination-addition. *J. Chem. Soc., Perkin Trans. 1* **1987**, 2235–2239.
- (4) Lopes, S. M. M.; Lemos, A.; Pinho e Melo, T. M. V. D. A hetero-Diels–Alder approach to functionalized 1H-tetrazoles: synthesis of tetrazolyl-1,2-oxazines, -oximes and 5-(1-aminoalkyl)-1H-tetrazoles. *Tetrahedron Lett.* **2010**, *51* (S1), 6756–6759.
- (5) Lopes, S. M. M.; Palacios, F.; Lemos, A.; Pinho e Melo, T. M. V. D. Diels–Alder reactions of 3-(1H-tetrazol-5-yl)-nitrosoalkenes: synthesis of functionalized 5-(substituted)-1H-tetrazoles. *Tetrahedron* **2011**, *67* (46), 8902–8909.
- (6) Panice, M. R.; Lopes, S. M. M.; Figueiredo, M. C.; Goes Ruiz, A. L. T.; Foglio, M. A.; Nazari Formaggio, A. S.; Sarragiotto, M. H.; Pinho e Melo, T. M. V. D. New 3-tetrazolyl- β -carboline and β -carboline-3-

carboxylates with anti-cancer activity. *Eur. J. Med. Chem.* **2019**, *179*, 123–132.

(7) Gilchrist, T. L.; Lemos, A. Reaction of pyrroles with ethyl 2-nitroso- and 2-azo-propenoates, and with ethyl cyanoformate *N*-oxide: A comparison of the reaction pathways. *J. Chem. Soc., Perkin Trans. 1* **1993**, 1391–1395.

(8) Gilchrist, T. L.; Roberts, T. G. Addition and cycloaddition reactions of the electrophilic vinyl nitroso compounds 3-nitrosobut-3-en-2-one, 2-nitrosopropenal, and ethyl 2-nitrosopropenoate. *J. Chem. Soc., Perkin Trans. 1* **1983**, 1283–1292.

(9) Nunes, S. C. C.; Lopes, S. M. M.; Gomes, C. S. B.; Lemos, A.; Pais, A. A. C. C.; Pinho e Melo, T. M. V. D. Reactions of nitrosoalkenes with dipyrromethanes and pyrroles: insight into the mechanistic pathway. *J. Org. Chem.* **2014**, *79* (21), 10456–10465.

(10) Lopes, S. M. M.; Nunes, S. C. C.; Caratão, C. C.; Pais, A. A. C. C.; Pinho e Melo, T. M. V. D. Reactivity of 1-arylnitrosoethylenes towards indole derivatives. *Monatsh. Chem.* **2016**, *147* (9), 1565–1573.

(11) Nascimento, B. F. O.; Lopes, S. M. M.; Pineiro, M.; Pinho e Melo, T. M. V. D. Current advances in the synthesis of valuable dipyrromethane scaffolds: classic and new methods. *Molecules* **2019**, *24* (23), 4348.

(12) Lopes, S. M. M.; Pinho e Melo, T. M. V. D. *Meso*-substituted corroles from nitrosoalkenes and dipyrromethanes. *J. Org. Chem.* **2020**, *85* (5), 3328–3335.

(13) Loudet, A.; Burgess, K. BODIPY dyes and their derivatives: syntheses and spectroscopic properties. *Chem. Rev.* **2007**, *107* (11), 4891–4932.

(14) Jorda, R.; Lopes, S. M. M.; Reznickova, E.; Krystof, V.; Pinho e Melo, T. M. V. D. Biological evaluation of dipyrromethanes in cancer cell lines: antiproliferative and pro-apoptotic properties. *ChemMedChem* **2017**, *12* (9), 701–711.

(15) Lopes, S. M. M.; Pineiro, M.; Pinho e Melo, T. M. V. D. Corroles and hexaphyrins: synthesis and application in cancer photodynamic therapy. *Molecules* **2020**, *25* (15), 3450.

(16) Praveen, P. J.; Parameswaran, P. S.; Majik, M. S. Bis(indolyl)-methane alkaloids: isolation, bioactivity, and syntheses. *Synthesis* **2015**, *47* (13), 1827–1837.

(17) Singh, A.; Kaur, G.; Banerjee, B. Recent developments on the synthesis of biologically significant bis/tris(indolyl)methanes under various reaction conditions: a review. *Curr. Org. Chem.* **2020**, *24* (6), 583–621.

(18) Pereira, N. A. M.; Lopes, S. M. M.; Lemos, A.; Pinho e Melo, T. M. V. D. On-water synthesis of dipyrromethanes via bis-hetero-Diels-Alder reaction of azo- and nitrosoalkenes with pyrrole. *Synlett* **2014**, *25* (3), 423–427.

(19) Grosso, C.; Cardoso, A. L.; Lemos, A.; Varela, J.; Rodrigues, M. J.; Custódio, L.; Barreira, L.; Pinho e Melo, T. M. V. D. Novel approach to bis(indolyl)methanes: *de novo* synthesis of 1-hydroxyiminomethyl derivatives with anti-cancer properties. *Eur. J. Med. Chem.* **2015**, *93*, 9–15.

(20) Grosso, C.; Cardoso, A. L.; Rodrigues, M. J.; Marques, C.; Barreira, L.; Lemos, A.; Pinho e Melo, T. M. V. D. Hetero-Diels-Alder approach to bis(indolyl)methanes. *Bioorg. Med. Chem.* **2017**, *25* (3), 1122–1131.

(21) Grosso, C.; Liber, M.; Brigas, A. F.; Pinho e Melo, T. M. V. D.; Lemos, A. Regioselectivity in hetero Diels-Alder Reactions. *J. Chem. Educ.* **2019**, *96* (1), 148–152.

(22) Murphy, J. J.; Driver, R. B.; Walsh, R.; Stephens, J. C. Synthesis of an imidazolidinone organocatalyst and its application in a Diels-Alder cycloaddition: a multistep experiment for the organic teaching laboratory. *J. Chem. Educ.* **2016**, *93* (9), 1626–1630.

(23) Celius, T. C. Fast hetero-Diels-Alder reactions using 4-phenyl-1,2,4-triazoline-3,5-dione (PTAD) as the dienophile. *J. Chem. Educ.* **2010**, *87* (11), 1236–1237.

(24) Sauvage, X.; Delaude, L. The synthesis of *N*-benzyl-2-azanorbornene via aqueous hetero Diels-Alder reaction. *J. Chem. Educ.* **2008**, *85* (11), 1538–1540.

(25) Baar, M. R.; Wustholz, K. Diels-Alder synthesis of *endo-cis-N*-phenylbicyclo[2.2.2]oct-5-en-2,3-dicarboximide. *J. Chem. Educ.* **2005**, *82* (9), 1393–1394.

(26) Augé, J.; Lubin-Germain, N. Hetero Diels-Alder reaction with aqueous glyoxylic acid. *J. Chem. Educ.* **1998**, *75* (10), 1285–1287.

(27) Cooley, J. H.; Williams, R. V. *Endo*- and *exo*-stereochemistry in the Diels-Alder reaction: kinetic *versus* thermodynamic control. *J. Chem. Educ.* **1997**, *74* (5), 582–585.

(28) Otto, S.; Engberts, J. B. F. N. Diels-Alder reactions in water. *Pure Appl. Chem.* **2000**, *72* (7), 1365–1372.