

HYPERSTRUCTURES ON LIE ALGEBROIDS

P. ANTUNES AND J.M. NUNES DA COSTA

ABSTRACT. We define hypersymplectic structures on Lie algebroids recovering, as particular cases, all the classical results and examples of hypersymplectic structures on manifolds. We prove a 1-1 correspondence theorem between hypersymplectic structures and (pseudo-)hyperkähler structures. We show that the hypersymplectic framework is very rich in already known compatible pairs of tensors such as Poisson-Nijenhuis, ΩN and $P\Omega$ structures.

Mathematics Subject Classification: 53D17, 53D05, 53C26

Keywords: Hypersymplectic, hyperkähler, Lie algebroid.

1. INTRODUCTION

Hypersymplectic structures on manifolds were introduced by Xu in [13]. Aiming to study hyperkähler structures on manifolds from the viewpoint of symplectic geometry, Xu was led to introduce, in a natural way, the notion of hypersymplectic structure. The aim of this paper is to define and study hypersymplectic structures on Lie algebroids. Our definition is inspired in [13] and, generalizing a result from [13] on manifolds, we prove that there exists a 1-1 correspondence between hypersymplectic and hyperkähler structures on a Lie algebroid. We also show that hypersymplectic structures on a Lie algebroid provide other type of interesting structures on the Lie algebroid, such as Poisson-Nijenhuis, ΩN and $P\Omega$ structures. The results of this paper are from [1] and were never published. We want to stress that the proofs we give here are different from those in [1]. We have improved the techniques and the proofs became much more elegant.

It is worth noticing that Hitchin [4] called hypersymplectic structures on a manifold M to what we call, in this work, para-hypersymplectic structures on the Lie algebroid TM . On the other hand, the structures which we call hypersymplectic are in 1-1 correspondence with hyperkähler structures (Theorem 6.6). For this reason, many authors refer to them simply as hyperkähler structures.

The paper is divided into six sections. Since our computations widely use the big bracket – the Poisson bracket induced by the symplectic structure on the cotangent bundle of a supermanifold, Section 1 contains a short review of Lie algebroids in the supergeometric framework and their deformation by Nijenhuis tensors and bivector fields. We also review the definition of Schouten-Nijenhuis bracket of multivectors on a Lie algebroid A as a derived bracket expression, and give a definition of the Frölicher-Nijenhuis bracket of two A -valued forms in supergeometric terms. In Section 2 we introduce the concept of ε -hypersymplectic structure on a Lie algebroid and we study the properties of the tensors induced on the Lie algebroid by this structure. One of the interesting features of ε -hypersymplectic structures, discussed in Section 3, is that pairs of these induced tensors determine well known structures on the Lie algebroid, such as $P\Omega$, ΩN and Poisson-Nijenhuis structures. In Section 4, we consider a particular case, that amounts to fix a sign in the ε -hypersymplectic structure, and we show that the number of $P\Omega$, ΩN and

Poisson-Nijenhuis structures on the Lie algebroid increases, when compared to the general case. In Section 5 we take the opposite sign, and consider two cases: hypersymplectic and para-hypersymplectic structures. In both cases we are able to define a pseudo-metric on the Lie algebroid. Using this pseudo-metric, we prove a 1-1 correspondence between (para-)hypersymplectic and (para-)hyperkähler structures on the Lie algebroid. The paper closes with an example in $T\mathbb{R}^4$ that provides many (para-)hypersymplectic structures.

2. PRELIMINARIES ON LIE ALGEBROIDS

2.1. Lie algebroids in supergeometric terms. We begin this section by introducing the supergeometric formalism, following the same approach as in [12, 10]. Given a vector bundle $A \rightarrow M$, we denote by $A[n]$ the graded manifold obtained by shifting the fibre degree by n . The graded manifold $T^*[2]A[1]$ is equipped with a canonical symplectic structure which induces a Poisson bracket on its algebra of functions $\mathcal{F} := C^\infty(T^*[2]A[1])$. This Poisson bracket is sometimes called the *big bracket* (see [5]).

Let us describe locally this Poisson algebra. Fix local coordinates $x^i, p_i, \xi^a, \theta_a$, $i \in \{1, \dots, n\}, a \in \{1, \dots, d\}$, on $T^*[2]A[1]$, where x^i, ξ^a are local coordinates on $A[1]$ and p_i, θ_a are their associated moment coordinates. In these local coordinates, the Poisson bracket is given by

$$\{p_i, x^i\} = \{\theta_a, \xi^a\} = 1, \quad i = 1, \dots, n, \quad a = 1, \dots, d,$$

while all the remaining brackets vanish.

The Poisson algebra of functions \mathcal{F} is endowed with an $(\mathbb{N} \times \mathbb{N})$ -valued bidegree. We define this bidegree (locally but it is well defined globally, see [12, 10]) as follows: the coordinates on the base manifold M , x^i , $i \in \{1, \dots, n\}$, have bidegree $(0, 0)$, while the coordinates on the fibres, ξ^a , $a \in \{1, \dots, d\}$, have bidegree $(0, 1)$ and their associated moment coordinates, p_i and θ_a , have bidegrees $(1, 1)$ and $(1, 0)$, respectively. We denote by $\mathcal{F}^{k,l}$ the $C^\infty(M)$ -module of functions of bidegree (k, l) and we verify that the big bracket has bidegree $(-1, -1)$, i.e.,

$$\{\mathcal{F}^{k_1, l_1}, \mathcal{F}^{k_2, l_2}\} \subset \mathcal{F}^{k_1+k_2-1, l_1+l_2-1}.$$

Let us recall that a *Lie algebroid* structure on a vector bundle $A \rightarrow M$ is a pair $(\rho, [\cdot, \cdot])$ where

- $\rho : A \rightarrow TM$ is a morphism of vector bundles, called the *anchor*,
- $[\cdot, \cdot]$ is a Lie bracket on the space of sections $\Gamma(A)$

satisfying the *Leibniz rule*

$$[X, fY] = f[X, Y] + (\rho(X) \cdot f)Y,$$

for all $f \in C^\infty(M)$ and $X, Y \in \Gamma(A)$.

Theorem 2.1 ([9, 11]). *There is a 1-1 correspondence between Lie algebroid structures on $A \rightarrow M$ and functions $\mu \in \mathcal{F}^{1,2}$ such that $\{\mu, \mu\} = 0$.*

The anchor and bracket associated to a given $\mu \in \mathcal{F}^{1,2}$ are defined, for all $X, Y \in \Gamma(A)$ and $f \in C^\infty(M)$, by the derived bracket expressions

$$(1) \quad \rho(X) \cdot f = \{\{X, \mu\}, f\} \quad \text{and} \quad [X, Y] = \{\{X, \mu\}, Y\}.$$

Moreover, the differential of the Lie algebroid is given by $d(\sigma) = \{\mu, \sigma\}$, for all $\sigma \in \Gamma(\wedge^\bullet A^*)$.

When using the supergeometric formalism, we shall denote a Lie algebroid by the pair (A, μ) instead of the triple $(A, \rho, [\cdot, \cdot])$.

2.2. Deformation of Lie algebroids. Let $(A, \rho, [\cdot, \cdot])$ be a Lie algebroid and μ the function in $\mathcal{F}^{1,2}$ corresponding to the Lie algebroid structure $(\rho, [\cdot, \cdot])$ on A .

Let $N \in \Gamma(A \otimes A^*)$ be a $(1, 1)$ -tensor seen as a vector bundle morphism $N : A \rightarrow A$. The deformation of the Lie bracket $[\cdot, \cdot]$ by N is defined, for all $X, Y \in \Gamma(A)$, by

$$[X, Y]_N = [NX, Y] + [X, NY] - N[X, Y].$$

The deformed structure $(\rho \circ N, [\cdot, \cdot]_N)$ is given, in supergeometric terms, by $\mu_N := \{N, \mu\} \in \mathcal{F}^{1,2}$. The deformation of μ_N by a $(1, 1)$ -tensor $S \in \Gamma(A \otimes A^*)$ is denoted by $\mu_{N,S}$, i.e.

$$(2) \quad \mu_{N,S} = \{S, \{N, \mu\}\},$$

while the deformed bracket associated to $\mu_{N,S}$ is denoted by $[\cdot, \cdot]_{N,S}$.

The Nijenhuis torsion of N , $\mathcal{T}N$, is given, for all $X, Y \in \Gamma(A)$, by

$$\mathcal{T}N(X, Y) = [NX, NY] - N[X, Y]_N.$$

A vector bundle endomorphism $N : A \rightarrow A$ is a *Nijenhuis* tensor on a Lie algebroid (A, μ) if its Nijenhuis torsion vanishes. In this case the deformed bracket $[\cdot, \cdot]_N$ is a Lie bracket and (A, μ_N) is a Lie algebroid.

When $N^2 = -\text{Id}_A$ (resp. $N^2 = \text{Id}_A$), N is said to be an *almost complex structure* (resp. *almost para-complex structure*). If moreover $\mathcal{T}N = 0$, then we can remove the prefix ‘‘almost’’ and N is a *complex structure* (resp. *para-complex structure*¹).

The deformation of the Lie bracket $[\cdot, \cdot]$ by a bivector $\pi \in \Gamma(\wedge^2 A)$ is a bracket on $\Gamma(A^*)$ defined, for all $\alpha, \beta \in \Gamma(A^*)$, by

$$[\alpha, \beta]_\pi = \mathcal{L}_{\pi^\sharp(\alpha)}\beta - \mathcal{L}_{\pi^\sharp(\beta)}\alpha - d(\pi(\alpha, \beta)),$$

where d is the differential of the Lie algebroid (A, μ) , \mathcal{L} is the Lie derivative determined by d and $\pi^\sharp : \Gamma(A^*) \rightarrow \Gamma(A)$ is defined by setting $\langle \beta, \pi^\sharp(\alpha) \rangle := \pi(\alpha, \beta)$, for all $\alpha, \beta \in \Gamma(A^*)$.

The deformed structure $(\rho \circ \pi^\sharp, [\cdot, \cdot]_\pi)$ corresponds to $\mu_\pi := \{\pi, \mu\} \in \mathcal{F}^{1,2}$. Moreover, if π is a Poisson bivector on A , $[\cdot, \cdot]_\pi$ is a Lie bracket and (A^*, μ_π) is a Lie algebroid.

Consider a $(1, 1)$ -tensor $N \in \Gamma(A \otimes A^*)$ and a bivector $\pi \in \Gamma(\wedge^2 A)$. The Magri-Morosi concomitant $C_{\pi,N}$ of π and N [8] is the difference between the deformed brackets on $\Gamma(A^*)$, $([\cdot, \cdot]_\pi)_{N^*}$ and $([\cdot, \cdot]_N)_\pi$, where N^* stands for the transpose of N . More precisely, the concomitant $C_{\pi,N}$ is given, for all $\alpha, \beta \in \Gamma(A^*)$, by

$$C_{\pi,N}(\alpha, \beta) = ([\alpha, \beta]_N)_\pi - ([\alpha, \beta]_\pi)_{N^*},$$

or, in terms of big bracket and functions on \mathcal{F} ,

$$C_{\pi,N} = \{\pi, \{N, \mu\}\} + \{N, \{\pi, \mu\}\} = \mu_{N,\pi} + \mu_{\pi,N}.$$

Notice that, when $C_{\pi,N} = 0$, applying the Jacobi identity we get

$$(3) \quad \{\pi, \{N, \mu\}\} = -\{N, \{\pi, \mu\}\} = \frac{1}{2} \{\{\pi, N\}, \mu\}.$$

¹In other works (including some of ours) these structures are called *(almost) product* instead of *(almost) para-complex*. We use the prefix ‘‘para’’ to express the fact that some sign is switched in comparison with the classical notion (without prefix). This change of terminology, when compared to previous works, enables us to express in a unified way the properties that are satisfied by both the classical structure and its para-version.

2.3. Schouten-Nijenhuis bracket and Frölicher-Nijenhuis bracket. Let (A, μ) be a Lie algebroid. The Schouten-Nijenhuis bracket is the natural extension, by derivation, of the Lie bracket on $\Gamma(A)$ to a bracket on $\Gamma(\wedge^\bullet A)$. Let $P \in \Gamma(\wedge^p A)$ and $Q \in \Gamma(\wedge^q A)$ be two multivectors on A . The Schouten-Nijenhuis bracket of P and Q is the $(p+q-1)$ -vector on A defined, in terms of big bracket, by

$$(4) \quad [P, Q]_{SN} = \{\{P, \mu\}, Q\}.$$

Notice that, in supergeometric terms, the Lie bracket $[\cdot, \cdot]$ and the Schouten-Nijenhuis bracket $[\cdot, \cdot]_{SN}$ are defined by the same² derived bracket expression (compare (4) and the second equation in (1)). Then, in order to simplify the notation and if there is no risk of confusion, the Schouten-Nijenhuis bracket on $\Gamma(\wedge^\bullet A)$ will be denoted by $[\cdot, \cdot]$, as the Lie bracket on $\Gamma(A)$.

Let $K \in \Gamma(\wedge^k A^* \otimes A)$ and $L \in \Gamma(\wedge^l A^* \otimes A)$ be two A -valued forms. We denote by $i_L K$ the section of $\wedge^{k+l-1} A^* \otimes A$ defined by

$$i_L K := \alpha_L \wedge (i_{X_L} \alpha_K) \otimes X_K,$$

where $K = \alpha_K \otimes X_K \in \Gamma(\wedge^k A^* \otimes A)$ and $L = \alpha_L \otimes X_L \in \Gamma(\wedge^l A^* \otimes A)$. The Frölicher-Nijenhuis bracket [3] of K and L , denoted by $[K, L]_{FN}$, is the section of $\wedge^{k+l} A^* \otimes A$ defined, in terms of big bracket, by the simple expression

$$(5) \quad [K, L]_{FN} = \{\{K, \mu\}, L\} + (-1)^{k(l+1)} \{i_L K, \mu\}.$$

It is known that the Nijenhuis torsion of a vector bundle endomorphism $N : A \rightarrow A$ is given by $\mathcal{T}N = -\frac{1}{2}[N, N]_{FN}$. Thus, using formula (5) and the notation introduced in (2), we get

$$\mathcal{T}N = \frac{1}{2}(\mu_{N,N} - \mu_{N^2}),$$

where $N^2 = N \circ N = i_N N$.

3. ε -HYPERSYMPLECTIC STRUCTURES ON LIE ALGEBROIDS

In this section we define an ε -hypersymplectic structure on a Lie algebroid, generalizing the definition of hypersymplectic triple given in [13] for manifolds. As we will see, an ε -hypersymplectic structure induces some other structures on the Lie algebroid. We study the main properties of these induced structures and the relations between them.

Let $A \rightarrow M$ be a vector bundle endowed with a Lie algebroid structure $(\rho, [\cdot, \cdot])$, corresponding to a function $\mu \in \mathcal{F}^{1,2}$.

Take three symplectic forms ω_1, ω_2 and $\omega_3 \in \Gamma(\wedge^2 A^*)$ with inverse Poisson bivectors π_1, π_2 and $\pi_3 \in \Gamma(\wedge^2 A)$, respectively. Then, for all $i \in \{1, 2, 3\}$ we have

$$\omega_i^b \circ \pi_i^\sharp = \text{Id}_{A^*} \quad \text{and} \quad \pi_i^\sharp \circ \omega_i^b = \text{Id}_A,$$

where $\omega_i^b : \Gamma(A) \rightarrow \Gamma(A^*)$ is defined by setting $\langle \omega_i^b(X), Y \rangle := \omega_i(X, Y)$, for all $X, Y \in \Gamma(A)$.

Let us define the *transition (1, 1)-tensors* $N_1, N_2, N_3 : \Gamma(A) \rightarrow \Gamma(A)$, by setting

$$(6) \quad N_i := \pi_{i-1}^\sharp \circ \omega_{i+1}^b,$$

considering the indices in \mathbb{Z}_3 .

²This was predictable because, for all $\Theta \in \mathcal{F}^{a,b}$, the operator $\{\Theta, \cdot\} : \mathcal{F} \rightarrow \mathcal{F}$ is a derivation of degree $a+b-2$ of the algebra (\mathcal{F}, \wedge) .

Remark 3.1. We consider 1, 2 and 3 as the representative elements of the equivalence classes of \mathbb{Z}_3 , i.e., $\mathbb{Z}_3 := \{[1], [2], [3]\}$. In what follows, although we omit the brackets, and write i instead of $[i]$, all the indices (and corresponding computations) must be thought in \mathbb{Z}_3 .

Definition 3.2. The triple $(\omega_1, \omega_2, \omega_3)$ is an ε -hypersymplectic structure on a Lie algebroid $(A, \rho, [\cdot, \cdot])$ if the transition $(1, 1)$ -tensors satisfy

$$(7) \quad N_i^2 = \varepsilon_i \text{Id}_A, \quad i = 1, 2, 3,$$

where the parameters $\varepsilon_i = \pm 1$ form the triple $\varepsilon = (\varepsilon_1, \varepsilon_2, \varepsilon_3)$.

Remark 3.3. Condition (7) can be written using only the symplectic forms and their inverse Poisson bivectors as, for example, in the following formula where the indices are treated as belonging to \mathbb{Z}_3 ,

$$(8) \quad \omega_{i+1}^b \circ \pi_{i-1}^\# = \varepsilon_i \omega_{i-1}^b \circ \pi_{i+1}^\#.$$

Proposition 3.4. Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on (A, μ) . The transition $(1, 1)$ -tensors satisfy, for all $i \in \{1, 2, 3\}$,

- (a) $(N_i)^{-1} = \varepsilon_i N_i$,
- (b) $N_3 N_2 N_1 = \text{Id}_A$ and $N_1 N_2 N_3 = \varepsilon_1 \varepsilon_2 \varepsilon_3 \text{Id}_A$.

Notice that, using the symplectic forms and their inverse Poisson bivectors, the condition (a) in Proposition 3.4 is given by

$$(9) \quad \pi_{i+1}^\# \circ \omega_{i-1}^b = \varepsilon_i \pi_{i-1}^\# \circ \omega_{i+1}^b.$$

Definition 3.5. We define $g \in \otimes^2 A^*$ by setting, for all $X, Y \in \Gamma(A)$,

$$g(X, Y) := \langle g^b X, Y \rangle,$$

where $g^b : A \rightarrow A^*$ is given by

$$(10) \quad g^b := \varepsilon_3 \varepsilon_2 \omega_3^b \circ \pi_1^\# \circ \omega_2^b.$$

In the next proposition we show that the definition of g^b is not affected by a circular permutation of the indices in equation (10) and that g is symmetric or skew-symmetric, depending on the sign of $\varepsilon_1 \varepsilon_2 \varepsilon_3$.

Proposition 3.6. Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on (A, μ) . The morphism g^b defined in (10) satisfies:

- (a) $g^b = \varepsilon_{i-1} \varepsilon_{i+1} \omega_{i-1}^b \circ \pi_i^\# \circ \omega_{i+1}^b$, for all $i \in \mathbb{Z}_3$;
- (b) $\varepsilon_{i-1} g^b = \omega_i^b \circ N_i$, for all $i \in \mathbb{Z}_3$;
- (c) $(g^b)^* = -\varepsilon_1 \varepsilon_2 \varepsilon_3 g^b$.

Proof. (a) Starting from the definition of g^b , applying formula (9) and then formula (8), we have:

$$g^b = \varepsilon_3 \varepsilon_2 \omega_3^b \circ \pi_1^\# \circ \omega_2^b = \varepsilon_2 \omega_3^b \circ \pi_2^\# \circ \omega_1^b = \varepsilon_1 \varepsilon_2 \omega_2^b \circ \pi_3^\# \circ \omega_1^b.$$

This last equality is obtained from (10), after a circular permutation of the indices. We can continue permuting the indices, using formulae (9) and (8), to get:

$$g^b = \varepsilon_1 \varepsilon_2 \omega_2^b \circ \pi_3^\# \circ \omega_1^b = \varepsilon_1 \omega_2^b \circ \pi_1^\# \circ \omega_3^b = \varepsilon_1 \varepsilon_2 \omega_1^b \circ \pi_2^\# \circ \omega_3^b.$$

Therefore, for any $i \in \mathbb{Z}_3$, $g^b = \varepsilon_{i-1} \varepsilon_{i+1} \omega_{i-1}^b \circ \pi_i^\# \circ \omega_{i+1}^b$.

- (b) Using (a) and then (8), we have

$$\varepsilon_{i-1} g^b = \varepsilon_{i+1} \omega_{i-1}^b \circ \pi_i^\# \circ \omega_{i+1}^b = \omega_i^b \circ \pi_{i-1}^\# \circ \omega_{i+1}^b.$$

Now, from the definition of N_i , given by (6), we get $\varepsilon_{i-1} g^b = \omega_i^b \circ N_i$.

- (c) Using (a) and the fact that both ω_i and π_i , $i = 1, 2, 3$, are skew-symmetric, we obtain

$$\begin{aligned} (g^b)^* &= -\varepsilon_{i-1}\varepsilon_{i+1} \omega_{i+1}^b \circ \pi_i^\# \circ \omega_{i-1}^b \\ &= -\varepsilon_{i-1} \omega_{i+1}^b \circ \pi_{i-1}^\# \circ \omega_i^b, \end{aligned}$$

where we used formula (9). Since, from (a), $g^b = \varepsilon_{i+1}\varepsilon_i \omega_{i+1}^b \circ \pi_{i-1}^\# \circ \omega_i^b$, we get

$$(g^b)^* = -\varepsilon_1\varepsilon_2\varepsilon_3 g^b.$$

□

We define the inverse of g , $g^{-1} \in \bigotimes^2 A$, by setting

$$g^{-1}(\alpha, \beta) := \langle (g^b)^{-1}(\alpha), \beta \rangle,$$

for all $\alpha, \beta \in \Gamma(A^*)$. As a direct consequence of this definition, we have $(g^{-1})^\# = (g^b)^{-1}$.

Proposition 3.7. *The morphisms g^b, ω_i^b and N_i satisfy the following:*

- (a) $\omega_i^b \circ N_i = \varepsilon_1\varepsilon_2\varepsilon_3 N_i^* \circ \omega_i^b = \varepsilon_{i-1}g^b$,
- (b) $\pi_i^\# \circ N_i^* = \varepsilon_1\varepsilon_2\varepsilon_3 N_i \circ \pi_i^\# = \varepsilon_{i+1}(g^{-1})^\#$,
- (c) $g^b \circ N_i = \varepsilon_1\varepsilon_2\varepsilon_3 N_i^* \circ g^b = \varepsilon_i\varepsilon_{i-1}\omega_i^b$,

for all indices in \mathbb{Z}_3 .

Proof. (a) Using (6), (9) and Proposition 3.4, we have

$$\omega_i^b \circ N_i = \omega_i \circ \pi_{i-1}^\# \circ \omega_{i+1}^b = \varepsilon_i \omega_i \circ \pi_{i+1}^\# \circ \omega_{i-1}^b = \varepsilon_{i-1}g^b.$$

The remaining part of the statement can be proved using Proposition 3.6(c) and then Proposition 3.6(b). Indeed,

$$\omega_i^b \circ N_i = \varepsilon_{i-1}g^b = -\varepsilon_i\varepsilon_{i+1}(g^b)^* = -\varepsilon_i\varepsilon_{i+1}(\varepsilon_{i-1}\omega_i^b \circ N_i)^* = \varepsilon_1\varepsilon_2\varepsilon_3 N_i^* \circ \omega_i^b.$$

Statements (b) and (c) can be proved directly, by the same kind of arguments, or using (a). □

Proposition 3.8. *For all sections $X, Y \in \Gamma(A)$ and for all indices in \mathbb{Z}_3 ,*

$$g(N_i X, N_i Y) = \varepsilon_{i-1}\varepsilon_{i+1} g(X, Y).$$

Proof. Let X and Y be sections of A . We have

$$\begin{aligned} g(N_i X, N_i Y) &= \langle g^b(N_i X), N_i Y \rangle = \varepsilon_1\varepsilon_2\varepsilon_3 \langle N_i^*(g^b X), N_i Y \rangle \\ &= \varepsilon_1\varepsilon_2\varepsilon_3 \langle g^b X, (N_i)^2(Y) \rangle = \varepsilon_{i-1}\varepsilon_{i+1} \langle g^b X, Y \rangle \\ &= \varepsilon_{i-1}\varepsilon_{i+1} g(X, Y), \end{aligned}$$

where we used Proposition 3.7(c) in the second equality. □

The next proposition is the continuation of Proposition 3.7, in the cases where the indices are different.

Proposition 3.9. *The morphisms induced by an ε -hypersymplectic structure $(\omega_1, \omega_2, \omega_3)$ satisfy the following relations, for all indices $i, k \in \mathbb{Z}_3, i \neq k$,*

- (a) $\omega_i^b \circ N_k = N_k^* \circ \omega_i^b = \begin{cases} \omega_{i-1}^b & , k = i + 1 \\ \varepsilon_{i-1}\omega_{i+1}^b & , k = i - 1; \end{cases}$
- (b) $\pi_i^\# \circ N_k^* = N_k \circ \pi_i^\# = \begin{cases} \varepsilon_{i+1}\pi_{i-1}^\# & , k = i + 1 \\ \pi_{i+1}^\# & , k = i - 1; \end{cases}$
- (c) $N_i \circ N_k = \varepsilon_1\varepsilon_2\varepsilon_3 N_k \circ N_i = \begin{cases} \varepsilon_i\varepsilon_{i+1}N_{i-1} & , k = i + 1 \\ \varepsilon_{i+1}N_{i+1} & , k = i - 1. \end{cases}$

Proof. The proofs of all statements are done by direct computations. We prove (a) to illustrate the kind of computations we use.

(a) If $k = i + 1$,

$$\omega_i^b \circ N_k = \omega_i^b \circ N_{i+1} = \omega_i^b \circ \pi_i^\# \circ \omega_{i-1}^b = \omega_{i-1}^b.$$

On the other hand, if $k = i - 1$,

$$\begin{aligned} \omega_i^b \circ N_k &= \omega_i^b \circ N_{i-1} = \omega_i^b \circ \pi_{i+1}^\# \circ \omega_i^b \\ &= \varepsilon_{i-1} \omega_i^b \circ \pi_i^\# \circ \omega_{i+1}^b = \varepsilon_{i-1} \omega_{i+1}^b, \end{aligned}$$

where we used (9) in the third equality. In both cases, $\omega_i^b \circ N_k$ is skew-symmetric, so that $\omega_i^b \circ N_k = -(\omega_i^b \circ N_k)^*$, i.e.,

$$\omega_i^b \circ N_k = N_k^* \circ \omega_i^b.$$

□

The next diagram shows all the relations between the morphisms induced by an ε -hypersymplectic structure on a Lie algebroid, recorded in Proposition 3.9.




FIGURE 1.

This is to be understood as the pattern for a tetrahedron $ABCD$, where the vertices A , B and C are labeling the Lie algebroid, while the vertex D labels the dual of the Lie algebroid. The morphism g^b does not appear in Figure 1 but, as shown in Figure 2, g^b is the altitude of the tetrahedron $ABCD$.




FIGURE 2.

The next proposition is a collection of already proved relations between ω_i^b , π_i^\sharp , N_k and g . The novelty is that we write these relations using the big bracket and functions on \mathcal{F} .

Proposition 3.10. *For all indices i, k in \mathbb{Z}_3 ,*

$$\begin{aligned}
 \text{(a)} \quad \{\omega_i, \pi_k\} &= \begin{cases} \text{Id}_A & , k = i \\ N_{i-1} & , k = i + 1 \\ \varepsilon_{i+1} N_{i+1} & , k = i - 1; \end{cases} \\
 \text{(b)} \quad \{N_k, N_{k+1}\} &= -\{N_{k+1}, N_k\} = \varepsilon_{k-1} (1 - \varepsilon_1 \varepsilon_2 \varepsilon_3) N_{k-1}; \\
 \text{(c)} \quad \{N_k, \omega_i\} &= \begin{cases} \varepsilon_{i-1} (1 + \varepsilon_1 \varepsilon_2 \varepsilon_3) g & , k = i \\ 2\omega_{i-1} & , k = i + 1 \\ 2\varepsilon_{i-1} \omega_{i+1} & , k = i - 1; \end{cases} \\
 \text{(d)} \quad \{N_k, \pi_i\} &= \begin{cases} -\varepsilon_{i+1} (1 + \varepsilon_1 \varepsilon_2 \varepsilon_3) g^{-1} & , k = i \\ -2\varepsilon_{i+1} \pi_{i-1} & , k = i + 1 \\ -2\pi_{i+1} & , k = i - 1. \end{cases}
 \end{aligned}$$

Proof. Let us prove (c) to exemplify how the already proved results translate in terms of big bracket and functions on \mathcal{F} . For a better understanding of these computations, see [9, 1].

(c) We can identify the function $\{N_k, \omega_i\} \in \mathcal{F}$ with a 2-form on A such that $\{N_k, \omega_i\}^b = \omega_i^b \circ N_k + N_k^* \circ \omega_i^b$. Then, using Proposition 3.7(a) and Proposition 3.9(a), we have:

$$\begin{aligned}
 \{N_k, \omega_i\}^b &= \omega_i^b \circ N_k + N_k^* \circ \omega_i^b \\
 &= \begin{cases} (1 + \varepsilon_1 \varepsilon_2 \varepsilon_3) \omega_i^b \circ N_i & , k = i \\ 2\omega_i^b \circ N_k & , k \neq i \end{cases} \\
 &= \begin{cases} \varepsilon_{i-1} (1 + \varepsilon_1 \varepsilon_2 \varepsilon_3) g^b & , k = i \\ 2\omega_{i-1}^b & , k = i + 1 \\ 2\varepsilon_{i-1} \omega_{i+1}^b & , k = i - 1. \end{cases}
 \end{aligned}$$

□

We recall an useful lemma from [7].

Lemma 3.11. *A bivector π , a 2-form ω and a $(1,1)$ -tensor N on a Lie algebroid (A, μ) , related by $N = \pi^\sharp \circ \omega^\flat$, satisfy the relation*

$$(11) \quad \{[\pi, \pi], \omega\}, \omega\} = \{\{\{\pi, d\omega\}, \pi\}, \omega\} - \{\{\pi, N\}, d\omega\} + 2\{\pi, \{\omega, \{N, \mu\}\}\} + 4\mathcal{T}N.$$

The next theorem, which is the main result of this section, shows that the transition tensors of an ε -hypersymplectic structure are Nijenhuis tensors.

Theorem 3.12. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) . The transition $(1,1)$ -tensors are Nijenhuis tensors, i.e., for all $i \in \{1, 2, 3\}$,*

$$\mathcal{T}N_i = 0.$$

Proof. By definition, $N_i = \pi_{i-1}^\sharp \circ \omega_{i+1}^\flat$ and, using the fact that ω_{i+1} is closed and π_{i-1} is a Poisson bivector, the formula (11) reduces to

$$\mathcal{T}N_i = -\frac{1}{2}\{\pi_{i-1}, \{\omega_{i+1}, \{N_i, \mu\}\}\}.$$

Now, using the Jacobi identity, Proposition 3.10(c) and the fact that ω_{i-1} is closed, we have

$$\{\omega_{i+1}, \{N_i, \mu\}\} = \{\{\omega_{i+1}, N_i\}, \mu\} = \{-2\varepsilon_i \omega_{i-1}, \mu\} = 0.$$

Therefore,

$$\mathcal{T}N_i = 0.$$

□

Notice that the fact that the 2-forms ω_i , $i = 1, 2, 3$, are closed is crucial in the proof of Theorem 3.12. All the other proofs in this section don't need this assumption.

4. INDUCED COMPATIBLE STRUCTURES

In this section we show that an ε -hypersymplectic structure induces many pairs of compatible structures such as, amongst others, pairs of compatible Poisson bivectors and Poisson-Nijenhuis structures.

Let (A, μ) be a Lie algebroid. Recall that a pair (π, N) , where π is a bivector and N is a $(1,1)$ -tensor on A is a *Poisson-Nijenhuis structure* (PN structure, for short) on (A, μ) if

$$(12) \quad [\pi, \pi] = 0, \quad \mathcal{T}N = 0, \quad N \circ \pi^\sharp = \pi^\sharp \circ N^* \quad \text{and} \quad C_{\pi, N} = 0.$$

A pair (ω, N) formed by a 2-form ω and a $(1,1)$ -tensor N on A is an ΩN structure on (A, μ) if

$$(13) \quad d\omega = 0, \quad \mathcal{T}N = 0, \quad \omega^\flat \circ N = N^* \circ \omega^\flat \quad \text{and} \quad d(\omega_N) = 0,$$

where $\omega_N(\cdot, \cdot) = \omega(N\cdot, \cdot)$ or, equivalently, $\omega_N^\flat = \omega^\flat \circ N$.

A pair (π, ω) formed by a bivector π and a 2-form ω on A is a $P\Omega$ structure on (A, μ) if

$$(14) \quad [\pi, \pi] = 0, \quad d\omega = 0 \quad \text{and} \quad d(\omega_N) = 0,$$

where N is the $(1,1)$ -tensor on A defined by $N = \pi^\sharp \circ \omega^\flat$.

Proposition 4.1. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) . For all $i \in \mathbb{Z}_3$,*

- (a) (π_{i+1}, ω_i) is a $P\Omega$ structure,
- (b) (π_{i-1}, ω_i) is a $P\Omega$ structure.

Proof. (a) We only need to prove that $d((\omega_i)_{\pi_{i+1} \circ \omega_i}) = 0$. Equation (6) gives $\pi_{i+1} \circ \omega_i = N_{i-1}$ and from Proposition 3.9 (a), we get $\omega_i^b \circ N_{i-1} = \varepsilon_{i-1} \omega_{i+1}^b$. Then, $d((\omega_i)_{\pi_{i+1} \circ \omega_i}) = d(\varepsilon_{i-1} \omega_{i+1}) = 0$. Therefore, (π_{i+1}, ω_i) is a $P\Omega$ structure.

(b) Analogously, (π_{i-1}, ω_i) is a $P\Omega$ structure. In fact,

$$d((\omega_i)_{\pi_{i-1} \circ \omega_i}) = d(\varepsilon_i(\omega_i)_{N_{i+1}}) = d(\varepsilon_i \omega_{i-1}) = 0.$$

□

Recall the following result from [7, 1]:

Proposition 4.2. *Let (π, ω) be a $P\Omega$ structure on a Lie algebroid (A, μ) and define $N := \pi^\sharp \circ \omega^b$. Then,*

- (a) (π, N) is a PN structure on A ,
- (b) (ω, N) is an ΩN structure on A .

Then, Proposition 4.1 has an immediate corollary.

Corollary 4.3. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) . For all $i, k \in \{1, 2, 3\}$, with $i \neq k$,*

- (a) (π_i, N_k) is a PN structure,
- (b) (ω_i, N_k) is an ΩN structure.

Notice that, in general, an ε -hypersymplectic structure induces 6 PN structures. In fact, contrary to what is claimed in [2], the pairs (π_i, N_i) , $i = 1, 2, 3$, are PN structures only when $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$, as we will see in the next section.

The remaining results of this section deal with compatibility between two bivectors or two $(1, 1)$ -tensors induced by an ε -hypersymplectic structure on a Lie algebroid (A, μ) . First we recall a result on PN structures.

Proposition 4.4 ([6]). *Let (π, N) be a PN structure on a Lie algebroid (A, μ) . Then π is a Poisson bivector on the Lie algebroid (A, μ_N) .*

Proposition 4.5. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) . For all $i, j \in \{1, 2, 3\}$, the Poisson bivectors π_i and π_j are compatible in the sense that $\pi_i + \pi_j$ is a Poisson bivector.*

Proof. We prove that, for all $i \in \mathbb{Z}_3$, the Poisson bivectors π_{i-1} and π_{i+1} are compatible. Since they are both Poisson bivectors, it is equivalent to prove that $[\pi_{i-1}, \pi_{i+1}] = 0$. From Corollary 4.3, for any $i \in \mathbb{Z}_3$, (π_{i+1}, N_i) is a PN structure. Then, from Proposition 4.4, π_{i+1} is a Poisson bivector on (A, μ_{N_i}) which means that $[\pi_{i+1}, \pi_{i+1}]_{N_i} = 0$, or equivalently,

$$\{\pi_{i+1}, \{\pi_{i+1}, \{N_i, \mu\}\}\} = 0.$$

The pairs (π_{i+1}, N_i) , $i \in \mathbb{Z}_3$, being PN structures, the concomitants C_{π_{i+1}, N_i} vanish and, using (3), we obtain

$$\left\{ \pi_{i+1}, \frac{1}{2} \{ \{ \pi_{i+1}, N_i \}, \mu \} \right\} = 0.$$

Applying Proposition 3.10(d) we get

$$\{\pi_{i+1}, \{\pi_{i-1}, \mu\}\} = 0,$$

which is equivalent to $[\pi_{i-1}, \pi_{i+1}] = 0$. □

Theorem 4.6. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) , such that $\varepsilon_1 \varepsilon_2 \varepsilon_3 = -1$. The Nijenhuis tensors N_i and N_j are compatible, for all $i, j \in \{1, 2, 3\}$, in the sense that $N_i + N_j$ is a Nijenhuis tensor.*

Proof. We know, from Theorem 3.12, that the transition (1,1)-tensors N_i , $i = 1, 2, 3$, are Nijenhuis tensors. We only need to prove that N_i and N_{i+1} are compatible, for all $i \in \mathbb{Z}_3$. Since $\mathcal{T}N = -\frac{1}{2}[N, N]_{FN}$, for all $N \in \Gamma(A \otimes A^*)$, the Nijenhuis tensors N_i and N_{i+1} are compatible if and only if $[N_i, N_{i+1}]_{FN} = 0$. Using formula (5), we have

$$(15) \quad \begin{aligned} [N_i, N_{i+1}]_{FN} &= \{\{N_i, \mu\}, N_{i+1}\} + \{i_{N_{i+1}}N_i, \mu\} \\ &= \varepsilon_{i+1} \{\{N_i, \mu\}, \{\omega_i, \pi_{i-1}\}\} + \varepsilon_i \varepsilon_{i+1} \{N_{i-1}, \mu\}, \end{aligned}$$

where we used Proposition 3.10(a) and Proposition 3.9(c). If we apply the Jacobi identity in the first term of the right hand side of (15), we get

$$(16) \quad [N_i, N_{i+1}]_{FN} = \varepsilon_{i+1} \{\{\{N_i, \mu\}, \omega_i\}, \pi_{i-1}\} + \varepsilon_{i+1} \{\omega_i, \{\{N_i, \mu\}, \pi_{i-1}\}\} + \varepsilon_i \varepsilon_{i+1} \mu_{N_{i-1}}.$$

Applying the Jacobi identity in the first term of the right hand side of (16) and taking into account the fact that ω_i is closed, we get

$$(17) \quad [N_i, N_{i+1}]_{FN} = \varepsilon_{i+1} \{\{\{N_i, \omega_i\}, \mu\}, \pi_{i-1}\} - \varepsilon_{i+1} \{\omega_i, \{\pi_{i-1}, \{N_i, \mu\}\}\} + \varepsilon_i \varepsilon_{i+1} \mu_{N_{i-1}}.$$

Let us do some computations on the second term of the right hand side of (17). Using (3), Proposition 3.10 and the closeness of ω_i , we have

$$(18) \quad \begin{aligned} \varepsilon_{i+1} \{\omega_i, \{\pi_{i-1}, \{N_i, \mu\}\}\} &= \varepsilon_{i+1} \left\{ \omega_i, \frac{1}{2} \{\{\pi_{i-1}, N_i\}, \mu\} \right\} \\ &= \varepsilon_{i+1} \{\omega_i, \{\varepsilon_i \pi_{i+1}, \mu\}\} = \varepsilon_i \varepsilon_{i+1} \{\{\omega_i, \pi_{i+1}\}, \mu\} \\ &= \varepsilon_i \varepsilon_{i+1} \{N_{i-1}, \mu\} = \varepsilon_i \varepsilon_{i+1} \mu_{N_{i-1}}. \end{aligned}$$

Replacing (18) in (17), we get

$$(19) \quad [N_i, N_{i+1}]_{FN} = \varepsilon_{i+1} \{\{\{N_i, \omega_i\}, \mu\}, \pi_{i-1}\}.$$

From Proposition 3.10(c), since $\varepsilon_1 \varepsilon_2 \varepsilon_3 = -1$, $\{N_i, \omega_i\} = 0$ and the proof is complete. \square

5. CASE $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$: MORE COMPATIBLE STRUCTURES

When $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$, from Proposition 3.6(b) we deduce that g is a 2-form on A , i.e., $g \in \Gamma(\wedge^2 A^*)$, and $g^{-1} \in \Gamma(\wedge^2 A)$ is a bivector on A . In the next theorem, the bivector g^{-1} is induced by a PN structure. Recall that, given a bivector $\pi \in \Gamma(\wedge^2 A)$ and a morphism $\varphi : A^* \rightarrow A^*$, we define the bivector $i_\varphi \pi$ by setting

$$i_\varphi \pi(\alpha, \beta) = \pi(\varphi\alpha, \beta) - \pi(\varphi\beta, \alpha),$$

for all $\alpha, \beta \in \Gamma(A^*)$.

Theorem 5.1. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) , such that $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$. The pair (π_i, N_i) is a PN structure and $i_{N_i^*} \pi_i = 2\varepsilon_{i+1} g^{-1}$, for all $i \in \{1, 2, 3\}$.*

The proof of Theorem 5.1 uses the following lemma:

Lemma 5.2. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) . For all $i \in \mathbb{Z}_3$,*

- (a) $\{\pi_i, \{N_i, \mu\}\} = -\varepsilon_i \{N_{i+1}, \{\pi_{i+1}, \mu\}\}$,
- (b) $\{N_i, \{\pi_i, \mu\}\} = -\varepsilon_i \{\pi_{i+1}, \{N_{i+1}, \mu\}\}$.

Proof. (a) From Proposition 3.10(a), we have

$$\{\pi_i, \{N_i, \mu\}\} = \varepsilon_i \{\pi_i, \{\{\omega_{i-1}, \pi_{i+1}\}, \mu\}\}.$$

Applying twice the Jacobi identity and using the facts that ω_{i-1} is closed and $[\pi_i, \pi_{i+1}] = 0$, we get

$$\begin{aligned} \{\pi_i, \{N_i, \mu\}\} &= \varepsilon_i \{\pi_i, \{\omega_{i-1}, \{\pi_{i+1}, \mu\}\}\} \\ &= \varepsilon_i \{\{\pi_i, \omega_{i-1}\}, \{\pi_{i+1}, \mu\}\}. \end{aligned}$$

Finally, using Proposition 3.10(a), we obtain

$$\{\pi_i, \{N_i, \mu\}\} = -\varepsilon_i \{N_{i+1}, \{\pi_{i+1}, \mu\}\}.$$

(b) The proof is similar to the proof of (a). □

The proof of Theorem 5.1 is now straightforward.

Proof. We already know from Proposition 3.7 that, when $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$,

$$\pi_i^\sharp \circ N_i^* = N_i \circ \pi_i^\sharp = \varepsilon_{i+1} (g^{-1})^\sharp,$$

so that

$$i_{N_i^*} \pi_i = 2\varepsilon_{i+1} g^{-1}.$$

It only remains to prove that $C_{\pi_i, N_i} = 0$. Using alternately Lemma 5.2(a) and Lemma 5.2(b), we obtain

$$\begin{aligned} \{\pi_i, \{N_i, \mu\}\} &= -\varepsilon_i \{N_{i+1}, \{\pi_{i+1}, \mu\}\} \\ &= \varepsilon_i \varepsilon_{i+1} \{\pi_{i-1}, \{N_{i-1}, \mu\}\} \\ &= -\varepsilon_i \varepsilon_{i+1} \varepsilon_{i-1} \{N_i, \{\pi_i, \mu\}\}. \end{aligned}$$

Since $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$, we get

$$\{\pi_i, \{N_i, \mu\}\} + \{N_i, \{\pi_i, \mu\}\} = 0,$$

i.e.,

$$C_{\pi_i, N_i} = 0.$$

Therefore, (π_i, N_i) is a PN structure. □

Recall the following result.

Proposition 5.3 ([6]). *If (π, N) is a PN structure on a Lie algebroid (A, μ) , then $i_{N^*} \pi$ is a Poisson bivector.*

As a direct consequence, from Theorem 5.1, we have:

Corollary 5.4. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) , such that $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$. The bivector g^{-1} is a Poisson bivector. Equivalently, the 2-form g is symplectic.*

Using Corollary 5.4, we can give a general version of Theorem 4.6 that includes both cases $\varepsilon_1 \varepsilon_2 \varepsilon_3 = -1$ and $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$.

Theorem 5.5. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) . The Nijenhuis tensors N_i and N_j are compatible, for all $i, j \in \{1, 2, 3\}$, in the sense that $N_i + N_j$ is a Nijenhuis tensor.*

Proof. The case $\varepsilon_1 \varepsilon_2 \varepsilon_3 = -1$ was treated in Theorem 4.6, where we proved, without using any assumption on the sign of $\varepsilon_1 \varepsilon_2 \varepsilon_3$, equation (19):

$$[N_i, N_{i+1}]_{FN} = \varepsilon_{i+1} \{\{\{N_i, \omega_i\}, \mu\}, \pi_{i-1}\}.$$

If $\varepsilon_1 \varepsilon_2 \varepsilon_3 = 1$, we may use Proposition 3.10(c) to get

$$[N_i, N_{i+1}]_{FN} = 2\varepsilon_{i+1} \varepsilon_{i-1} \{\{g, \mu\}, \pi_{i-1}\} = 0,$$

because g is a closed 2-form. □

When $\varepsilon_1\varepsilon_2\varepsilon_3 = 1$, taking into account the fact that g is a symplectic form and g^{-1} is a Poisson bivector, we can add 6 new (and non-trivial) $P\Omega$ structures to the ones of Proposition 4.1.

Proposition 5.6. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) , such that $\varepsilon_1\varepsilon_2\varepsilon_3 = 1$. Then, for all $i \in \{1, 2, 3\}$, the pairs (π_i, g) and (g^{-1}, ω_i) are $P\Omega$ structures.*

Proof. We only need to show that $d(g_{\pi_i^\sharp \circ g^b}) = 0$ and $d((\omega_i)_{(g^{-1})^\sharp \circ \omega_i^b}) = 0$. From Proposition 3.7(a), we deduce that $\pi_i^\sharp \circ g^b = \varepsilon_{i-1}N_i$ and we have

$$g_{\pi_i^\sharp \circ g^b} = \varepsilon_{i-1}g_{N_i} = \varepsilon_i\omega_i,$$

where we used Proposition 3.7(c) in the last equality. The form ω_i being closed, we conclude that

$$d(g_{\pi_i^\sharp \circ g^b}) = 0.$$

In a similar way, we have

$$d((\omega_i)_{(g^{-1})^\sharp \circ \omega_i^b}) = d(\varepsilon_{i+1}(\omega_i)_{N_i}) = d(\varepsilon_{i+1}\varepsilon_{i-1}g) = 0.$$

□

From Proposition 4.2, the 6 $P\Omega$ structures of Proposition 5.6 induce 6 PN structures and 6 ΩN structures as stated in the following two corollaries. Notice that for 3 of these induced PN structures we proved it directly in Theorem 5.1.

Corollary 5.7. *For all $i \in \{1, 2, 3\}$, the pairs (π_i, N_i) and (g^{-1}, N_i) are PN structures.*

Corollary 5.8. *For all $i \in \{1, 2, 3\}$, the pairs (ω_i, N_i) and (g, N_i) are ΩN structures.*

In the last result of this section we prove that, when $\varepsilon_1\varepsilon_2\varepsilon_3 = 1$, the new Poisson bivector, g^{-1} , is compatible with any π_i , $i = 1, 2, 3$.

Proposition 5.9. *Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) , such that $\varepsilon_1\varepsilon_2\varepsilon_3 = 1$. For all $i \in \{1, 2, 3\}$, the Poisson bivectors π_i and g^{-1} are compatible.*

Proof. Using Proposition 3.10(d), we have

$$[\pi_i, g^{-1}] = \{\{\pi_i, \mu\}, g^{-1}\} = -\frac{1}{2}\varepsilon_{i+1}\{\{\pi_i, \mu\}, \{N_i, \pi_i\}\}.$$

Applying the Jacobi identity and using the fact that π_i is a Poisson bivector, i.e., $\{\{\pi_i, \mu\}, \pi_i\} = 0$, we get

$$\begin{aligned} [\pi_i, g^{-1}] &= -\frac{1}{2}\varepsilon_{i+1}(\{\{\{\pi_i, \mu\}, N_i\}, \pi_i\} + \{N_i, \{\{\pi_i, \mu\}, \pi_i\}\}) \\ &= \frac{1}{2}\varepsilon_{i+1}\{\{N_i, \{\pi_i, \mu\}\}, \pi_i\}. \end{aligned}$$

From Theorem 5.1, we know that $C_{\pi_i, N_i} = 0$ or, equivalently, $\{N_i, \{\pi_i, \mu\}\} = -\{\pi_i, \{N_i, \mu\}\}$ and so the last equality becomes

$$[\pi_i, g^{-1}] = -\frac{1}{2}\varepsilon_{i+1}\{\{\pi_i, \{N_i, \mu\}\}, \pi_i\} = -\frac{1}{2}\varepsilon_{i+1}[\pi_i, \pi_i]_{N_i} = 0,$$

where we used, in the last equality, Proposition 4.4 and the fact that (π_i, N_i) is a PN structure. Therefore, π_i and g^{-1} are compatible Poisson bivectors. □

6. CASE $\varepsilon_1\varepsilon_2\varepsilon_3 = -1$: (PARA)-HYPERKÄHLER STRUCTURES

In this section we consider ε -hypersymplectic structures with $\varepsilon_1\varepsilon_2\varepsilon_3 = -1$. As we will see, these structures are in 1-1 correspondence with (para-)hyperkähler structures, a notion we will define later. First, let us consider two different cases of an ε -hypersymplectic structure with $\varepsilon_1\varepsilon_2\varepsilon_3 = -1$.

Definition 6.1. Let $(\omega_1, \omega_2, \omega_3)$ be an ε -hypersymplectic structure on a Lie algebroid (A, μ) , such that $\varepsilon_1\varepsilon_2\varepsilon_3 = -1$.

- If $\varepsilon_1 = \varepsilon_2 = \varepsilon_3 = -1$, then $(\omega_1, \omega_2, \omega_3)$ is said to be a *hypersymplectic* structure on A .
- Otherwise, $(\omega_1, \omega_2, \omega_3)$ is said to be a *para-hypersymplectic* structure on A .

It is clear that all para-hypersymplectic structures satisfy, eventually after a cyclic permutation of the indices, $\varepsilon_1 = \varepsilon_2 = 1$ and $\varepsilon_3 = -1$. In the sequel, every para-hypersymplectic structures will be considered in such form.

Definition 6.2. Let (A, μ) be a Lie algebroid. A *pseudo-metric* on $A \rightarrow M$ is a symmetric and non-degenerate $C^\infty(M)$ -linear map $\mathbf{g} : \Gamma(A) \rightarrow \Gamma(A^*)$. Furthermore, if \mathbf{g} is *positive definite*, i.e., if g satisfies

$$\langle \mathbf{g}(X), X \rangle > 0,$$

for all non vanishing sections $X \in \Gamma(A)$, then \mathbf{g} is a *metric* on A .

Because $\varepsilon_1\varepsilon_2\varepsilon_3 = -1$, the proof of the next result is a consequence of Proposition 3.6.

Proposition 6.3. Let $(\omega_1, \omega_2, \omega_3)$ be a (para-)hypersymplectic structure on a Lie algebroid (A, μ) . The morphism g^b defined in (10) is a pseudo-metric on A .

Note. In the sequel, we do not require the metric to be positive definite. However, in order to simplify the terminology we will omit the “pseudo” prefix.

Definition 6.4. A (para-)hermitian structure is a pair (\mathbf{g}, I) where \mathbf{g} is a metric and I is a (para-)complex tensor such that, for all $X, Y \in \Gamma(A)$,

$$\langle \mathbf{g}(IX), IY \rangle = -\langle \mathbf{g}(X), I^2(Y) \rangle.$$

Notice that the condition above may be written simply as $\mathbf{g} \circ I + I^* \circ \mathbf{g} = 0$.

Definition 6.5. A (para-)hyperkähler structure on a Lie algebroid (A, μ) is a quadruple $(\mathbf{g}, I_1, I_2, I_3)$, where the pairs $(\mathbf{g}, I_j)_{j=1,2}$ are both (para-)hermitian structures on A , the morphisms I_1 and I_2 anti-commute, $I_3 = I_1I_2$, and $\omega_i^b := \mathbf{g} \circ I_i$ are closed 2-forms, for $i = 1, 2, 3$.

The main result of this section is a direct consequence of Proposition 6.3, Proposition 3.8 and Proposition 3.9.

Theorem 6.6. Let (A, μ) be a Lie algebroid. The triple $(\omega_1, \omega_2, \omega_3)$ is a (para-)hypersymplectic structure on A if and only if (g, N_1, N_2, N_3) is a (para-)hyperkähler structure on A .

Note. The 1-1 correspondence in Theorem 6.6 concerns pseudo (para-)hyperkähler structures. We may restrict ourselves to the more usual (para-)hyperkähler structures (g, N_1, N_2, N_3) , with a positive definite metric g . These are in 1-1 correspondence with (para-)hypersymplectic structures $(\omega_1, \omega_2, \omega_3)$, such that g (defined by (10)) is positive definite.

To conclude, we address a simple example in $T\mathbb{R}^4$ which provide many (para-)hypersymplectic structures.

Example 6.7. Consider the coordinates (x, y, p, q) on \mathbb{R}^4 and the following six 2-forms

$$\begin{aligned}\omega_1 &= dx \wedge dp + dy \wedge dq; & \omega_4 &= dx \wedge dp - dy \wedge dq; \\ \omega_2 &= dx \wedge dq + dp \wedge dy; & \omega_5 &= dx \wedge dq - dp \wedge dy; \\ \omega_3 &= dx \wedge dy - dp \wedge dq; & \omega_6 &= dx \wedge dy + dp \wedge dq.\end{aligned}$$

The 2-forms $\omega_i \in \Gamma(\wedge^2(T^*\mathbb{R}^4))$, $i = 1, \dots, 6$, are symplectic forms on \mathbb{R}^4 . For all pairwise different indices $i, j, k \in \{1, \dots, 6\}$, the triple $(\omega_i, \omega_j, \omega_k)$ is a (para-)hypersymplectic structure on the Lie algebroid $T\mathbb{R}^4$. More precisely:

- (1) The triples $(\omega_1, \omega_2, \omega_3)$ and $(\omega_4, \omega_5, \omega_6)$ are hypersymplectic structures.
- (2) The 9 triples $(\omega_i, \omega_j, \omega_k)$ where $1 \leq i < j \leq 3$ and $k \in \{4, 5, 6\}$ are para-hypersymplectic structures.
- (3) The 9 triples $(\omega_i, \omega_j, \omega_k)$ where $4 \leq i < j \leq 6$ and $k \in \{1, 2, 3\}$ are para-hypersymplectic structures.

Acknowledgments. This work was partially supported by CMUC-FCT (Portugal) and FCT grants PTDC/MAT/099880/2008 and PEst-C/MAT/UI0324/2011 through European program COMPETE/FEDER.

REFERENCES

- [1] P. Antunes, *Crochets de Poisson gradués et applications: structures compatibles et généralisations des structures hyperkählériennes*, Thèse de doctorat de l'École Polytechnique, (2010).
- [2] J. R. Camacaro and J. F. Cariñena, Hyper-complex structures and Lie algebroids, Proceedings of the XI Fall Workshop on Geometry and Physics, Publ. R. Soc. Mat. Esp., vol. 6, R. Soc. Mat. Esp., Madrid, 2004, pp. 197–202.
- [3] A. Frölicher and A. Nijenhuis, Theory of vector-valued differential forms. I. Derivations of the graded ring of differential forms, *Nederl. Akad. Wetensch. Proc. Ser. A.* **59** = *Indag. Math.* **18** (1956) 338–359.
- [4] N. J. Hitchin, Hypersymplectic quotients, La "Mécanique analytique" Lagrange et son héritage, Atti della Accademia delle Scienze di Torino, Classe de Scienze fisiche, Matematiche e Naturali, Supplemento al numero 124, Accademia delle Scienze di Torino, Torino, 1990, pp. 169–180.
- [5] Y. Kosmann-Schwarzbach, Jacobian quasi-bialgebras and quasi-Poisson Lie groups. In *Mathematical aspects of classical field theory*, Contemp. Math., 132, Amer. Math. Soc., 1992, pp. 459–489.
- [6] Y. Kosmann-Schwarzbach and F. Magri, Poisson-Nijenhuis structures, *Ann. Inst. H. Poincaré, Phys. Théor.* **53** (1) (1990) 35–81.
- [7] Y. Kosmann-Schwarzbach and V. Rubtsov, Compatible structures on Lie algebroids and Monge-Ampère operators, *Acta Appl. Math.* **109** (1) (2010) 101–135.
- [8] F. Magri and C. Morosi, A geometrical characterization of integrable Hamiltonian systems through the theory of Poisson-Nijenhuis manifolds, Quaderno S 19, Univ. of Milan, 1984.
- [9] D. Roytenberg, *Courant algebroids, derived brackets and even symplectic supermanifolds*. PhD thesis, UC Berkeley, 1999. math.DG/9910078.
- [10] D. Roytenberg, Quasi-Lie bialgebroids and twisted Poisson manifolds, *Lett. Math. Phys.* **61** (2) (2002) 123–137.
- [11] A. Vaintrob, Lie algebroids and homological vector fields, *Russian Math. Surveys* **52** (2) (1997) 428–429.
- [12] T. Voronov, Graded manifolds and Drinfeld doubles for Lie bialgebroids, in *Quantization, Poisson brackets and beyond*, T. Voronov, ed., Contemp. Math. 315, Amer. Math. Soc., Providence, RI, 2002, pp. 131–168.
- [13] P. Xu, Hyper-Lie Poisson structures, *Ann. Sc. l'École Normale Sup.* **30** (3) (1997) 279–302.

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3001-454 COIMBRA, PORTUGAL

E-mail address: `pantunes@mat.uc.pt`

CMUC, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF COIMBRA, 3001-454 COIMBRA, PORTUGAL

E-mail address: `jmcosta@mat.uc.pt`