

Homological properties of quantised Borel-Schur algebras and resolutions of quantised Weyl modules

Stephen Donkin*

*Department of
Mathematics
University of York
Heslington, York
UK*

stephen.donkin@york.ac.uk

Ana Paula Santana[†]

*CMUC, Department of
Mathematics
University of Coimbra
Coimbra
Portugal*

aps@mat.uc.pt

Ivan Yudin[‡]

*CMUC, Department of
Mathematics
University of Coimbra
Coimbra
Portugal*

yudin@mat.uc.pt

Abstract

We continue the development of the homological theory of quantum general linear groups previously considered by the first author. The development is used to transfer information to the representation theory of quantised Schur algebras. The acyclicity of induction from some rank-one modules for quantised Borel-Schur subalgebras is deduced. This is used to prove the exactness of the complexes recently constructed by Boltje and Maisch, giving resolutions of the co-Specht modules for Hecke algebras.

1 Introduction

In [SY12] the last two authors constructed characteristic free projective resolutions of the Weyl modules for the classical Schur algebra. Then, using the Schur functor, obtained resolutions by permutation modules of the co-Specht modules for the symmetric group. This last result allowed them to prove Conjecture 3.4 of Boltje and Hartman [BH11].

*The first author wishes to thank CMUC for hospitality in May 2012, when the major part of the research for this paper was done, and in April 2013, when the final version was produced.

[†]This work was partially supported by the Centro de Matemática da Universidade de Coimbra (CMUC), funded by the European Regional Development Fund through the program COMPETE and by the Portuguese Government through the FCT - Fundação para a Ciência e a Tecnologia under the project PEst-C/MAT/UI0324/2011.

[‡]The third author's work is supported by the FCT Grant SFRH/BPD/31788/2006.

The key ingredients of [SY12] are the use of the normalised bar resolution in the context of Borel-Schur algebras and Woodcock's Theorem [Woo94b], which reduces the construction of projective resolutions for Weyl modules to the construction of projective resolutions for rank-one modules for the Borel-Schur algebra. The original motivation of the present paper was to extend the results of [SY12] to the context of quantised Schur algebras and Hecke algebras. This is easily achieved once one has a quantised version of the generalization of Woodcock's theorem given in [SY12].

Fix positive integers n and r , a commutative ring R and an invertible element q in R . Consider the quantised Schur algebra $S_{R,q}(n, r)$ and the quantised positive Borel-Schur algebra $S_{R,q}^+(n, r)$. For each partition $\lambda = (\lambda_1, \dots, \lambda_n)$ of r there is a rank-one module R_λ for $S_{R,q}^+(n, r)$. The induced module

$$W_\lambda^{R,q} := S_{R,q}(n, r) \otimes_{S_{R,q}^+(n,r)} R_\lambda$$

is the Weyl module associated with λ . Following [SY12], we work in the category of $S_{R,q}^+(n, r)$ -modules and use the normalised bar resolution to construct a projective resolution of R_λ . Next we apply the induction functor $S_{R,q}(n, r) \otimes_{S_{R,q}^+(n,r)} -$ to this resolution and obtain a complex $B_{*,\lambda}^{R,q}$ of finite length

$$\cdots \rightarrow B_{1,\lambda}^{R,q} \rightarrow B_{0,\lambda}^{R,q} \rightarrow W_\lambda^{R,q} \rightarrow 0,$$

where each $B_{k,\lambda}^{R,q}$ is a projective $S_{R,q}(n, r)$ -module.

To show that this complex is exact we use Theorem 8.4, which is the quantised version of Woodcock's Theorem. So $B_{*,\lambda}^{R,q}$ is a projective resolution of the quantised Weyl module $W_\lambda^{R,q}$ and it is simple to see that this resolution is universal, that is

$$B_{*,\lambda}^{R,q} \cong B_{*,\lambda}^{\mathcal{Z},t} \otimes_{\mathcal{Z}} R,$$

where $\mathcal{Z} = \mathbb{Z}[t, t^{-1}]$ is the universal quantization ring.

Write $\mathcal{H}_{R,q}$ for the Hecke algebra over R associated with the symmetric group Σ_r . In [BM12], Boltje and Maisch constructed, for each composition $\lambda = (\lambda_1, \dots, \lambda_n)$ of r , a complex $\tilde{\mathcal{C}}_*^\lambda$ of left $\mathcal{H}_{R,q}$ -modules and proved that it is exact in degrees 0 and -1 . Specializing to $q = 1$, $\tilde{\mathcal{C}}_*^\lambda$ coincides with the complex constructed in [BH11]. Suppose that λ is a partition of r . Then the last module in $\tilde{\mathcal{C}}_*^\lambda$ is the dual of the Specht module S^λ over $\mathcal{H}_{R,q}$. It was proved in [SY12] that in this situation upon specializing to $q = 1$ the resulting complex is exact. It is natural to conjecture that the same should be true for an arbitrary q .

Returning to our setting, we choose $n \geq r$ and fix λ a partition of r into at most n parts. We apply the Schur functor

$$F: S_{R,q}(n, r)\text{-mod} \rightarrow \mathcal{H}_{R,q}\text{-mod}$$

to our resolution $B_{*,\lambda}^{R,q}$ and obtain an exact complex $F(B_{*,\lambda}^{R,q})$ which we prove to be isomorphic to $\tilde{\mathcal{C}}_*^\lambda$. This proves the exactness of $\tilde{\mathcal{C}}_*^\lambda$.

We approach the quantisation of Woodcock's Theorem, as described above, via the representation theory of the quantum general linear group $G(n)$ of degree n , introduced in [DD91]. In fact we take this opportunity to develop the homological theory previously considered in [Don96] and [Don98]. The focus here is on a comparison between the homological algebra in the category of polynomial modules and in the full category of modules for the quantum group. We work over an arbitrary field \mathbb{K} and non-zero parameter $q \in \mathbb{K}$.

Let $B(n)$ be the negative Borel (quantum) subgroup of G . We prove in particular that the derived functors of induction $R^i \text{ind}_{B(n)}^{G(n)}$ take polynomial modules to polynomial modules, Corollary 7.7. Furthermore we show that if V is a homogeneous polynomial $B(n)$ -module of degree r then $R^i \text{ind}_{B(n)}^{G(n)} V = 0$ for all $i > r$, Lemma 6.2. In general the tensor product is not commutative in the category of modules for a quantum group. However, we show that if L and M are B -modules and L is one dimensional then the B -module $L \otimes M$ and $M \otimes L$ are isomorphic, Proposition 7.1. Using this property and a Koszul resolution we show that the polynomial part of the coordinate algebra of $B(n)$ is acyclic for the induction functor. This leads to the fact that the derived functors of induction applied to a polynomial $B(n)$ -module are the same whether computed in the polynomial category or the full module category, Theorem 7.5. Kempf's Vanishing Theorem for representations of quantum groups, when expressed in the polynomial category, is essentially the quantised version of Woodcock's Theorem, over a field. Some further work is needed to express this in terms of the acyclicity theorem for induction over Schur algebras, over an arbitrary coefficient ring, mentioned above, Theorem 8.4.

Though not needed for the application to resolutions we also take the opportunity to give the generalisation to the quantum Borel subgroup $B(n)$ of another theorem of Woodcock, [Woo92], Theorem 7 and [Woo94a] (see also [Woo97] for related material obtained by working with global bases). This theorem asserts that the extension groups between polynomial $B(n)$ -modules of the same degree whether calculated in the polynomial category or the full $B(n)$ -module category are the same, Theorem 5.2. We approach the quantised version by considering the derived functors of the functor pol , which takes a $B(n)$ -module to its largest polynomial submodule. Though in detail it looks quite different it is in spirit rather close to the approach of [Woo92], and we gratefully acknowledge the influence of this unpublished work.

The organization of the present paper is as follows. We first study the homological results for quantum $G(n)$ and its negative Borel subgroup. Then we use this to obtain the quantised version of Woodcock's Theorem, Theorem 8.4. In the last part of the paper we construct universal projective resolutions for quantised Weyl modules. Using these resolutions, we prove the exactness of Boltje and Maisch complexes for dominant weights.

2 Restriction and induction of comodules

We fix a field \mathbb{K} . For a vector space V over \mathbb{K} we write V^* for the linear dual $\text{Hom}_{\mathbb{K}}(V, \mathbb{K})$ and if W is also a vector space over \mathbb{K} we write simply $V \otimes W$ for the tensor product $V \otimes_{\mathbb{K}} W$. We write id_X for the identity map on a set X .

For a coalgebra $A = (A, \delta_A, \epsilon_A)$ over \mathbb{K} we write $\text{Comod}(A)$ for the category of right A -comodules and write $\text{comod}(A)$ for the category of finite dimensional right A -comodules. We recall for future use the definition of the coefficient space of an A -comodule. Let $V = (V, \tau)$ be a right A -comodule and let $\{v_i : i \in I\}$, be a \mathbb{K} -basis of V . The coefficient space $\text{cf}(V)$ is the \mathbb{K} -span of the elements $f_{ij} \in A$ defined by the equations

$$\tau(v_i) = \sum_{j \in I} v_j \otimes f_{ji}$$

for $i \in I$. (This space is independent of the choice of basis. For further properties see [Gre76].)

Let $B = (B, \delta_B, \epsilon_B)$ also be a coalgebra and suppose $\phi: A \rightarrow B$ is a coalgebra map. Recall that for $V = (V, \tau) \in \text{Comod}(A)$ we have

$$\phi_0(V) = (V, (\text{id}_V \otimes \phi) \circ \tau) \in \text{Comod}(B).$$

If $f: V \rightarrow V'$ is a morphism of right comodules then the same map $f: V \rightarrow V'$ is also a morphism of B -comodules. In this way we have an exact functor $\phi_0: \text{Comod}(A) \rightarrow \text{Comod}(B)$, with $\phi_0(f) = f$, for f a morphism of A -comodules. We call ϕ_0 the ϕ -restriction (or just restriction) functor.

More interestingly perhaps, we have the ϕ -induction functor $\phi^0: \text{Comod}(B) \rightarrow \text{Comod}(A)$. This is described in [Don80], Section 3, and we briefly recall the construction and some properties. If X is a \mathbb{K} -vector space (possibly with extra structure) we write $|X| \otimes A$ for the vector space $X \otimes A$ regarded as an A -comodule with structure map $\text{id}_X \otimes \delta_A$. Let $(W, \mu) \in \text{Comod}(B)$. The set of all $s \in W \otimes A$ such that

$$(\mu \otimes \text{id}_A)(s) = (\text{id}_W \otimes (\phi \otimes \text{id}_A) \circ \delta_B)(s) \in W \otimes B \otimes A$$

is an A -subcomodule of $|W| \otimes A$, which we denote $\phi^0(W)$. If $f: W \rightarrow W'$ is a morphism of B -comodules then the map $f \otimes \text{id}_A$ restricts to an A -comodule map $\phi^0(f): \phi^0(W) \rightarrow \phi^0(W')$. In this way we obtain a left exact functor $\phi^0: \text{Comod}(B) \rightarrow \text{Comod}(A)$. Let $V = (V, \lambda) \in \text{Comod}(A)$ and $W = (W, \mu) \in \text{Comod}(B)$. We have a natural isomorphism $\text{Hom}_B(\phi_0(V), W) \rightarrow \text{Hom}_A(V, \phi^0(W))$, taking $\alpha \in \text{Hom}_B(\phi_0(V), W)$ to $\tilde{\alpha} = (\alpha \otimes \text{id}_A) \circ \lambda$.

Suppose now that A is finite dimensional. We consider the dual algebra $S = A^* = \text{Hom}_{\mathbb{K}}(A, \mathbb{K})$. Given a right A -comodule V with structure map $\tau: V \rightarrow V \otimes A$ we may also regard V as a left S -module with action $\alpha v = (\text{id}_V \otimes \alpha)\tau(v)$. If $\theta: V \rightarrow V'$ is a morphism of right A -comodules then, regarding V and V' as left S -modules, $\theta: V \rightarrow V'$ is also a morphism in the category of left S -modules. In this way

we have an equivalence between the categories of finite dimensional right A -comodules and of finite dimensional left S -modules. For finite dimensional right A -comodules V, V' this equivalence of categories induces a \mathbb{K} -linear isomorphism $\text{Ext}_A^i(V, V') \rightarrow \text{Ext}_S^i(V, V')$ in each degree i .

If S is a \mathbb{K} -algebra and V is a left (resp. right) A -module then the linear dual V^* is naturally a right (resp. left) S -module. Now suppose $\phi: A \rightarrow B$ is a morphism of finite dimensional \mathbb{K} -coalgebras and let $T = B^*$. The linear dual $\phi^*: T \rightarrow S$ is a \mathbb{K} -algebra map. Now A is naturally an (S, S) -bimodule with left action $\alpha a = (\text{id}_A \otimes \alpha)\delta_A(a)$ and right action $a\beta = (\beta \otimes \text{id}_A)\delta(a)$, for $a \in A, \alpha, \beta \in S$. We view an S -module also as a T -module via ϕ^* .

We have the natural linear isomorphism $\eta: V \otimes A \rightarrow (V^* \otimes A^*)^*$. The tensor product $V^* \otimes_T A^*$ is a quotient of $(V^* \otimes A^*)^*$ and we thus identify $(V^* \otimes_T A^*)^*$ with a subspace of $(V^* \otimes A^*)^*$. From the definitions one checks that an element y of $V \otimes A$ lies in $\phi^0(V)$ if and only if $\eta(y)$ lies in $(V^* \otimes_T A^*)^*$. The map η restricts to an isomorphism of left A -modules

$$\phi^0 V \rightarrow (V^* \otimes_T A^*)^*.$$

It follows that the derived functors of ϕ^0 are given as follows.

Proposition 2.1. *Let $\phi: A \rightarrow B$ be a morphism of finite dimensional coalgebras over \mathbb{K} . Then for $V \in \text{comod}(B)$ we have*

$$R^i \phi^0 V = (\text{Tor}_i^{B^*}(V^*, A^*))^*$$

for $i \geq 0$.

3 The quantum polynomial algebra in n^2 variables

We shall work with the quantum general linear groups defined in [DD91]. We briefly recall the construction and some properties, starting with the construction of the quantum polynomial algebra. We fix $n \geq 1$. Let R be a commutative ring and let $q \in R$. We write $A_{R,q}(n)$ for the R -algebra given by generators c_{ij} , $1 \leq i, j \leq n$, and relations:

$$\begin{aligned} c_{ir}c_{is} &= c_{is}c_{ir}, \text{ for } 1 \leq i, r, s \leq n; \\ c_{jr}c_{is} &= qc_{is}c_{jr}, \text{ for } 1 \leq i < j \leq n, 1 \leq r \leq s < n; \\ c_{js}c_{ir} &= c_{ir}c_{js} - (q-1)c_{is}c_{jr}, \text{ for } 1 \leq i < j \leq n, 1 \leq r < s \leq n. \end{aligned}$$

We call the elements c_{ij} the (i, j) coordinate elements of $A_{R,q}(n)$. Since the relations are homogeneous, $A_{R,q}(n)$ has an R -algebra grading $A_{R,q}(n) = \bigoplus_{r \geq 0} A_{R,q}(n, r)$ in which each coordinate element has degree 1. Then by [DD91], Theorem 1.1.8 the elements

$$c_{11}^{m_{11}} c_{12}^{m_{12}} \cdots c_{1n}^{m_{1n}} c_{21}^{m_{21}} \cdots c_{nn}^{m_{nn}},$$

with $m_{11}, \dots, m_{nn} \geq 0$, form an R -basis of $A_{R,q}(n)$. We make this slightly more formal.

Let $r \geq 0$. As in [Gre07], we write $I(n, r)$ for the set of maps $i: \{1, \dots, r\} \rightarrow \{1, \dots, n\}$. We identify $i \in I(n, r)$ with the sequence (i_1, \dots, i_r) in the obvious way. For $i, j \in I(n, r)$ we write c_{ij} for the product $c_{i_1 j_1} \dots c_{i_r j_r}$. We write $i \leq j$ if $i_a \leq j_a$, for all $1 \leq a \leq r$, and write $i < j$ if $i \leq j$ and $i \neq j$. We write $Y(n, r)$ for the set of all pairs $(i, j) \in I(n, r)$ such that $i_1 \leq \dots \leq i_r$ and whenever, for some $1 \leq a < r$, we have $i_a = i_{a+1}$ then $j_a \leq j_{a+1}$. We write $Y(n)$ for the disjoint union of the sets $Y(n, r)$, $r \geq 0$.

Lemma 3.1. *The elements c_{ij} , with $i, j \in Y(n)$ form an R -basis of $A_{R,q}(n)$ and, for $r \geq 0$, the elements c_{ij} , with $i, j \in Y(n, r)$, form an R -basis of $A_{R,q}(n, r)$.*

We write I for the ideal of $A_{R,q}(n)$ generated by all c_{ij} , with $1 \leq i < j \leq n$. We leave it to the reader to check (by an easy induction argument using the defining relations) the following result.

Lemma 3.2. *The ideal I has R -basis c_{ij} , with $(i, j) \in Y(n, r)$ for some r and $i_a < j_a$, for some $1 \leq a \leq r$.*

We set $\bar{A}(n) = A(n)/I$. For $f \in A(n)$ we set $\bar{f} = f + I \in \bar{A}(n)$. For $r \geq 0$, we write $\bar{Y}(n, r)$ for the set of all $(i, j) \in Y(n, r)$ such that $i \geq j$. We set $\bar{Y}(n) = \bigcup_{r \geq 0} \bar{Y}(n, r)$. As an R -module we have $A_{R,q}(n) = I \oplus D$, where $D = \bigoplus_{(i,j) \in \bar{Y}(n)} R c_{ij}$. Hence we have the following.

Lemma 3.3. *$\bar{A}_{R,q}(n)$ has R -basis \bar{c}_{ij} , $(i, j) \in \bar{Y}(n)$, and, for $r \geq 0$, $\bar{A}_{R,q}(n, r)$ has R -basis \bar{c}_{ij} , $(i, j) \in \bar{Y}(n, r)$.*

4 Quantum general linear groups

Let \mathbb{K} be a field. The category of quantum groups over \mathbb{K} is the dual of the category of Hopf algebras over \mathbb{K} . More informally, we shall use the expression “ G is a quantum group over \mathbb{K} ” to indicate that we have in mind a Hopf algebra over \mathbb{K} , which we will denote $\mathbb{K}[G]$ and call the coordinate algebra of G . By the expression “ $\theta: G \rightarrow H$ is a morphism of quantum groups (over \mathbb{K})” we indicate that G and H are quantum groups and that we have in mind a Hopf algebra morphism from $\mathbb{K}[H]$ to $\mathbb{K}[G]$, which we call the comorphism of θ and denote $\theta^\#$. We shall say that a quantum group H is a (quantum) subgroup of a quantum group G over \mathbb{K} to indicate that $\mathbb{K}[H] = \mathbb{K}[G]/I_H$ for some Hopf ideal I_H of $\mathbb{K}[G]$, which we call the defining ideal of H in G . If H is a quantum subgroup of the quantum group G then by the inclusion map $i: H \rightarrow G$ we mean the quantum group homomorphism such that $i^\#: \mathbb{K}[G] \rightarrow \mathbb{K}[H]$ is the natural map.

Let G be a quantum group over \mathbb{K} . By the category of left G -modules we mean the category of right $\mathbb{K}[G]$ -comodules. We write $\text{Mod}(G)$ for the category of left G -modules and $\text{mod}(G)$ for the category of finite dimensional left G -modules. For $V, W \in \text{Mod}(G)$ and

$i \geq 0$ we write $\text{Ext}_G^i(V, W)$ for $\text{Ext}_{\mathbb{K}[G]}^i(V, W)$. Let H be a quantum subgroup of G . Then we have the induction functor $\text{ind}_H^G = \phi^0: \text{Mod}(H) \rightarrow \text{Mod}(G)$, where $\phi = i^\#$ is the comorphism of the inclusion map $i: H \rightarrow G$. The functor ind_H^G is left exact so we have the derived functors $R^i \text{ind}_H^G: \text{Mod}(H) \rightarrow \text{Mod}(G)$, for $i \geq 0$.

We work with the quantum coordinate algebra $A_{R,q}(n)$ of the previous section, now taking $R = \mathbb{K}$ and $q \neq 0$. We write Σ_r for the symmetric group on $\{1, 2, \dots, r\}$, for r a positive integer. To simplify notation we will omit \mathbb{K} and q in subscript in the objects defined in the previous section, where confusion seems unlikely.

By [DD91], Theorem 1.4.2, $A(n)$ has a unique structure of a bialgebra with comultiplication $\delta: A(n) \rightarrow A(n) \otimes A(n)$ and counit $\epsilon: A(n) \rightarrow \mathbb{K}$, satisfying

$$\delta(c_{ij}) = \sum_{r=1}^n c_{ir} \otimes c_{rj}, \quad \epsilon(c_{ij}) = \delta_{ij}$$

for $1 \leq i, j \leq n$ and where δ_{ij} is the Kronecker delta.

The quantum determinant

$$d = \sum_{\pi \in \Sigma_n} \text{sgn}(\pi) c_{1,\pi(1)} c_{1,\pi(2)} \cdots c_{n,\pi(n)}$$

is a group-like element of $A(n)$. Here $\text{sgn}(\pi)$ denotes the sign of a permutation π . Furthermore, we have $c_{ij}d = q^{i-j}dc_{ij}$ for $1 \leq i, j \leq n$ (see [DD91, Section 4]). It follows that we can form the Ore localisation $A(n)_d$. The bialgebra structure on $A(n)$ extends to $A(n)_d$ and indeed the localisation $A(n)_d$ is a Hopf algebra. We write $G(n)$ for the quantum group with coordinate algebra $\mathbb{K}[G(n)] = A(n)_d$.

We write $B(n)$ for the quantum subgroup whose defining ideal $I_{B(n)}$ is generated by all c_{ij} , with $1 \leq i < j \leq n$. We write $T(n)$ for the quantum subgroup whose defining ideal is generated by all c_{ij} with $1 \leq i, j \leq n$ and $i \neq j$. The inclusion map $A(n) \rightarrow \mathbb{K}[G(n)]$ gives rise to an injective map $\bar{A}(n) \rightarrow \mathbb{K}[B(n)]$ by which we identify $\bar{A}(n)$ with a subbialgebra of $\mathbb{K}[B(n)]$. A $G(n)$ -module V is called polynomial (resp. polynomial of degree r) if $\text{cf}(V) \leq A(n)$ (resp. $\text{cf}(V) \leq A(n, r)$) and a $B(n)$ -module M is called polynomial (resp. polynomial of degree r) if $\text{cf}(M) \leq \bar{A}(n)$ (resp. $\text{cf}(M) \leq \bar{A}(n, r)$). We shall often identify a polynomial $G(n)$ -module (resp. $B(n)$ -module) with the corresponding $A(n)$ -comodule (resp. $\bar{A}(n)$ -comodule).

We shall also need the parabolic (quantum) subgroups containing $B(n)$. We fix a string $a = (a_1, \dots, a_m)$ of positive integers whose sum is n . We let $I(a)$ be the ideal of $k[G(n)]$ generated by all c_{ij} such that $1 \leq i < j \leq a_1$ or $a_1 + \dots + a_r < i < j \leq a_1 + \dots + a_{r+1}$ for some $1 \leq r < m$. Then $I(a)$ is a Hopf ideal and we denote by $P(a)$ the quantum subgroup of $G(n)$ with defining ideal $I(a)$. Thus we have $P(1, 1, \dots, 1) = G(n)$ and $P(n) = B(n)$. For $1 \leq i < m$ we shall write P_i for the ‘‘minimal parabolic’’ $P(a)$, where $a = (1, 1, \dots, 2, 1, \dots, 1)$ (with 2 in the i th position).

We now introduce certain combinatorial objects associated with the representation theory of $G(n)$ and its subgroups, following [Don96].

We set $X(n) = \mathbb{Z}^n$. We shall write δ_n , or simply δ , for $(n-1, n-2, \dots, 1, 0) \in X(n)$. For $1 \leq i \leq n$ we set $\epsilon_i = (0, \dots, 0, 1, 0, \dots, 0)$ (with 1 in the i th position). We have the dominance order \leq on $X(n)$: for $\lambda = (\lambda_1, \dots, \lambda_n), \mu = (\mu_1, \dots, \mu_n)$ we write $\lambda \leq \mu$ if $\lambda_1 + \dots + \lambda_i \leq \mu_1 + \dots + \mu_i$, for $1 \leq i < n$, and $\lambda_1 + \dots + \lambda_n = \mu_1 + \dots + \mu_n$.

We write $X^+(n)$ for the set of all $\lambda = (\lambda_1, \dots, \lambda_n) \in X(n)$ with $\lambda_1 \geq \dots \geq \lambda_n$. Elements of $X(n)$ will sometimes be called weights and elements of $X^+(n)$ called dominant weights. We write $\Lambda(n)$ for the set of polynomial weights, i.e., the set of $\lambda = (\lambda_1, \dots, \lambda_n) \in X(n)$ with all $\lambda_i \geq 0$, and write $\Lambda^+(n)$ for the set of dominant weights, i.e., $X^+(n) \cap \Lambda(n)$. We define the degree of a polynomial weight $\lambda = (\lambda_1, \dots, \lambda_n)$ by $\deg(\lambda) = \lambda_1 + \dots + \lambda_n$. For $r \geq 0$ we define $\Lambda(n, r) \subset \Lambda(n)$ to be the set of all polynomial weights of degree r (or compositions of r). We define the length, $\text{len}(\lambda)$, of a polynomial weight λ to be 0 if $\lambda = 0$ and to be the number of non-zero entries of λ if $\lambda \neq 0$.

For $\lambda = (\lambda_1, \dots, \lambda_n) \in X(n)$ we have a one dimensional $B(n)$ -module \mathbb{K}_λ : the comodule structure map $\tau: \mathbb{K}_\lambda \rightarrow \mathbb{K}_\lambda \otimes k[B(n)]$ takes $v \in \mathbb{K}_\lambda$ to $v \otimes (c_{11}^{\lambda_1} \dots c_{nn}^{\lambda_n} + I_{B(n)})$. We regard \mathbb{K}_λ also as a $T(n)$ -module by restriction.

The modules $\mathbb{K}_\lambda, \lambda \in X(n)$, form a complete set of pairwise non-isomorphic irreducible $T(n)$ -modules. For a $T(n)$ -module V we have the weight space decomposition $V = \bigoplus_{\lambda \in X(n)} V^\lambda$, where V^λ is a direct sum of copies of $\mathbb{K}_\lambda, \lambda \in X(n)$.

For $\lambda \in X(n)$ the induced module $\text{ind}_{B(n)}^{G(n)} \mathbb{K}_\lambda$ is non-zero if and only if $\lambda \in X^+(n)$. We set $\nabla(\lambda) = \text{ind}_{B(n)}^{G(n)} \mathbb{K}_\lambda$, for $\lambda \in X^+(n)$. The socle $L(\lambda)$ of $\nabla(\lambda)$ is simple. The modules $L(\lambda), \lambda \in X^+(n)$, form a complete set of pairwise non-isomorphic irreducible $G(n)$ -modules and the modules $L(\lambda), \lambda \in \Lambda^+(n)$, form a complete set of pairwise non-isomorphic irreducible polynomial $G(n)$ -modules. We will write D for the determinant module, i.e., the (one dimensional) left $G(n)$ -module $L(1, \dots, 1)$.

Let $1 \leq i < n$. Let $\lambda = (\lambda_1, \dots, \lambda_n) \in X(n)$ and suppose that $m = \lambda_i - \lambda_{i+1} \geq 0$. We define $\nabla_i(\lambda) = \text{ind}_{B(n)}^{P_i(n)} \mathbb{K}_\lambda$. Then $\nabla_i(\lambda)$ has weights $\lambda - r(\epsilon_i - \epsilon_{i+1}), 0 \leq r \leq m$, each occurring with multiplicity one (see [Don96], p251).

We shall need that a $G(n)$ -module whose composition factors have the form $L(\lambda)$ with $\lambda \in \Lambda^+(n)$ (resp. $\lambda \in \Lambda^+(n, r)$) is polynomial (resp. polynomial of degree r). Given the results of [Don96] this follows from the arguments in the classical case in [Don86]. We make this explicit.

Let $\pi \subseteq X^+(n)$. We say that a $G(n)$ -module V belongs to π if each composition factor of V belongs to $\{L(\lambda) \mid \lambda \in \pi\}$. For an arbitrary $G(n)$ -module we write $O_\pi(V)$ for the largest $G(n)$ -submodule of V belonging to π . Regarding $\mathbb{K}[G(n)]$ as the left regular $G(n)$ -module we define $A(\pi) = O_\pi(\mathbb{K}[G])$. Then, by the arguments for the classical case, [Don86], Section 1.2, one has the following.

Lemma 4.1. *$A(\pi)$ is a subcoalgebra of $\mathbb{K}[G(n)]$ and a $G(n)$ -module V belongs to π if and only if $\text{cf}(V) \leq A(\pi)$.*

In the case $\pi = \Lambda^+(n, r), r \geq 0$, we have $A(\pi) = A(n, r)$, see

[Don96], p263, and taking $\pi = \Lambda^+(n)$, since $\Lambda^+(n) = \bigcup_{r \geq 0} \Lambda^+(n, r)$, we have $A(\pi) = A(n)$. Hence the above lemma gives:

Lemma 4.2. *A $G(n)$ -module V is polynomial (resp. polynomial of degree r) if and only if each composition factor of V belongs to $\{L(\lambda) \mid \lambda \in \Lambda^+(n)\}$ (resp. $\{L(\lambda) \mid \lambda \in \Lambda^+(n, r)\}$).*

Remark 4.3. *We note that if M is a polynomial B -module then $\text{ind}_B^G M$ is a polynomial G -module. It is enough to check this for M finite dimensional since induction commutes with direct limits. By the left exactness of induction and the Lemma 4.2 it is enough to check this for M one dimensional. So we may assume that $M = \mathbb{K}_\lambda$ for some $\lambda \in \Lambda(n)$. But now we have*

$$\text{ind}_B^G \mathbb{K}_\lambda = \begin{cases} \nabla(\lambda), & \text{if } \lambda \in \Lambda^+(n); \\ 0, & \text{if } \lambda \notin \Lambda^+(n) \end{cases}$$

and, in particular, $\text{ind}_B^G M$ is polynomial.

5 Extensions of B -modules and polynomial B -modules

Though it is not needed for the application to resolutions of modules for the Borel-Schur algebras, we take this opportunity to put on record the quantised version of [Woo97, Theorem 7] giving that, for homogeneous polynomial $B(n)$ -modules, the extension groups $\text{Ext}^i(V, X)$ are the same whether calculated in the module category of the Borel-Schur algebra or the full $B(n)$ -module category. Though the proof given here looks rather different it is similar at key points to that of Woodcock in the classical case, [Woo92] and we gratefully acknowledge the influence of [Woo92]. A later proof was given in [Woo94a] using the deep theory of cohomology of line bundles on Schubert varieties due to van der Kallen, [vdK89] and related results are to be found in the later work [Woo97] using the theory of global bases.

In this section we adopt the following notation. We put $B = B(n)$, $A = \bar{A}(n)$ and $A_m = \mathbb{K}[\bar{c}_{m1}, \bar{c}_{m2}, \dots, \bar{c}_{mm}]$, $x_m = \bar{c}_{mm}$, $y_m = \bar{c}_{mm}^{-1}$, for $1 \leq m \leq n$. For $\alpha = (\alpha_1, \dots, \alpha_n) \in \Lambda(n)$ we put $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ and $y^\alpha = y_1^{\alpha_1} \dots y_n^{\alpha_n}$. We write simply d for the restriction of the determinant to the quantum subgroup $B(n)$, i.e., $d = x_1 \dots x_n$.

We have $A = A_1 \otimes \dots \otimes A_n$ and it is easy to check that

$$A_m/x_m A_m \cong \begin{cases} A_{m-1}, & \text{for } 1 < m \leq n; \\ \mathbb{K}, & \text{for } m = 1 \end{cases}$$

as $B(n)$ -modules.

We shall need the following result.

Lemma 5.1. *Let $\lambda \in \Lambda(n)$ and suppose $1 \leq m \leq n$ is such that $\lambda_m \neq 0$. Let Z be a polynomial B -module such that for each weight μ of Z , we have $\mu_m = \mu_{m+1} = \dots = \mu_n = 0$. Then we have*

$$\text{Ext}_A^i(\mathbb{K}_\lambda, Z \otimes A_{m+1} \otimes \dots \otimes A_n) = 0$$

for all $i \geq 0$. In particular, we have

$$\mathrm{Ext}_A^i(\mathbb{K}_\lambda, A/x_m A) = 0$$

for all $i \geq 0$.

Proof. Suppose not and let i be minimal for which the lemma fails. Since $\mathrm{Ext}_A^i(\mathbb{K}_\lambda, -)$ commutes with direct limits, the lemma fails for some finite dimensional Z and by the long exact sequence we may assume that $Z = \mathbb{K}_\mu$, for some $\mu \in \Lambda(n)$, with $\mu_m = \mu_{m+1} = \cdots = \mu_n = 0$. Now $\mathbb{K}_\mu \otimes A_{m+1} \otimes \cdots \otimes A_n$ has socle $\mathbb{K}_\mu \otimes \mathbb{K}[x_{m+1}, \dots, x_n]$ and so for each weight ν of the socle we have $\nu_m = 0$. Since $\lambda_m \neq 0$ there can be no non-zero image of \mathbb{K}_λ in the socle of $\mathbb{K}_\mu \otimes A_{m+1} \otimes \cdots \otimes A_n$ and therefore $\mathrm{Hom}_A(\mathbb{K}_\lambda, \mathbb{K}_\mu \otimes A_{m+1} \otimes \cdots \otimes A_n) = 0$. Thus we must have $i > 0$.

Now we have a short exact sequence of A -comodules (or polynomial B -modules)

$$0 \rightarrow \mathbb{K}_\mu \rightarrow A_1 \otimes \cdots \otimes A_{m-1} \rightarrow Q \rightarrow 0$$

and for each weight ν of $A_1 \otimes \cdots \otimes A_{m-1}$, and hence Q , we have $\nu_m = 0$. Tensoring with $A_{m+1} \otimes \cdots \otimes A_n$ we obtain the short exact sequence

$$\begin{aligned} 0 \rightarrow \mathbb{K}_\mu \otimes A_{m+1} \otimes \cdots \otimes A_n &\rightarrow A_1 \otimes \cdots \otimes A_{m-1} \otimes A_{m+1} \otimes \cdots \otimes A_n \\ &\rightarrow Q \otimes A_{m+1} \otimes \cdots \otimes A_n \rightarrow 0. \end{aligned}$$

But now $A_1 \otimes \cdots \otimes A_{m-1} \otimes A_{m+1} \otimes \cdots \otimes A_n$ is an injective A -comodule (since it is a direct summand of A , viewed as the right regular comodule) so we get

$$\mathrm{Ext}_A^i(\mathbb{K}_\lambda, \mathbb{K}_\mu \otimes A_{m+1} \otimes \cdots \otimes A_n) = \mathrm{Ext}_A^{i-1}(\mathbb{K}_\lambda, Q \otimes A_{m+1} \otimes \cdots \otimes A_n)$$

from the long exact sequence. This is 0, by the minimality of i , and so we are done. \square

We now consider the functor $\mathrm{pol} : \mathrm{Mod}(B) \rightarrow \mathrm{Comod}(A)$, taking $X \in \mathrm{Mod}(B)$ to the largest polynomial submodule of X . For a morphism $\theta : X \rightarrow X'$, of B -modules, $\mathrm{pol}(\theta) : \mathrm{pol}(X) \rightarrow \mathrm{pol}(X')$ is the restriction of θ .

For $V \in \mathrm{Comod}(A)$, $X \in \mathrm{Mod}(B)$, since the image of any B -module homomorphism from V to X is contained in $\mathrm{pol}(X)$, we have $\mathrm{Hom}_B(V, X) = \mathrm{Hom}_A(V, \mathrm{pol}(X))$. Thus we get a factorisation of left exact functors

$$\mathrm{Hom}_B(V, -) = \mathrm{Hom}_A(V, -) \circ \mathrm{pol}.$$

Moreover, $\mathrm{pol}(\mathbb{K}[B]) = A$ and it follows that pol takes injective B -modules to injective A -comodules. Thus, for $V \in \mathrm{Comod}(A)$, $X \in \mathrm{Mod}(B)$, we have a Grothendieck spectral sequence, with second page $\mathrm{Ext}_A^i(V, R^j \mathrm{pol} X)$, converging to $\mathrm{Ext}_B^*(V, X)$. In particular, if $k > 0$ and $R^j \mathrm{pol}(X) = 0$ for all $0 < j < k$, then we have the 5-term exact sequence

$$\begin{aligned} 0 \rightarrow \mathrm{Ext}_A^k(V, \mathrm{pol} X) &\rightarrow \mathrm{Ext}_B^k(V, X) \rightarrow \mathrm{Hom}_A(V, R^k \mathrm{pol} X) \\ &\rightarrow \mathrm{Ext}_A^{k+1}(V, \mathrm{pol} X) \rightarrow \mathrm{Ext}_B^{k+1}(V, X). \end{aligned} \quad (1)$$

Theorem 5.2. (i) Let X be a polynomial B -module. Then we have $R^i \text{pol}(X) = 0$, for all $i > 0$.

(ii) If V is also a polynomial B -module then the above spectral sequence degenerates and we have $\text{Ext}_A^i(V, X) = \text{Ext}_B^i(V, X)$, for all $i \geq 0$.

Proof. For $k > 0$ we prove by induction the statement $P(k)$: for all polynomial B -modules V', X' we have $R^i \text{pol} X' = 0$ for all $0 < i < k$ and $\text{Ext}_A^i(V', X') = \text{Ext}_B^i(V', X')$, for all $0 \leq i < k$.

Note that $P(1)$ is true since $\text{Hom}_A(V', X') = \text{Hom}_B(V', X')$ for polynomial B -modules V', X' . We now assume $P(k)$ and deduce $P(k+1)$.

We claim that $R^k \text{pol} A = 0$. Assume, for a contradiction, that this is not the case. Then the B -socle of $R^k \text{pol} A$ is not zero so we have $\text{Hom}_B(\mathbb{K}_\lambda, R^k \text{pol} A) \neq 0$ for some $\lambda \in \Lambda(n)$. Dimension shifting, using the short exact sequence

$$0 \rightarrow A \rightarrow \mathbb{K}[B] \rightarrow \mathbb{K}[B]/A \rightarrow 0$$

gives $\text{Ext}_B^{k-1}(\mathbb{K}_\lambda, \mathbb{K}[B]/A) \neq 0$. Now $\mathbb{K}[B]$ has an ascending exhaustive filtration $A \subseteq d^{-1}A \subseteq d^{-2}A \subseteq \dots$ and $\text{Ext}_B^{k-1}(\mathbb{K}_\lambda, -)$ commutes with direct limits so we must have $\text{Ext}_B^{k-1}(\mathbb{K}_\lambda, d^{-s}A/A) \neq 0$ for some $s > 0$. Hence we have $\text{Ext}_B^{k-1}(\mathbb{K}_\lambda, y^\alpha A/A) \neq 0$, for some $\alpha \in \Lambda(n)$. We choose $\alpha, \beta \in \Lambda(n)$ with $\beta_i \leq \alpha_i$, for $1 \leq i \leq n$, and with $\deg(\alpha) - \deg(\beta)$ minimal subject to the condition $\text{Ext}_B^{k-1}(\mathbb{K}_\lambda, y^\alpha A/y^\beta A) \neq 0$. Note that in fact we must have $\deg(\alpha) = \deg(\beta) + 1$ since if $\gamma \in \Lambda(n)$ with $\beta_i \leq \gamma_i \leq \alpha_i$, for all i , then we get a short exact sequence

$$0 \rightarrow y^\gamma A/y^\beta A \rightarrow y^\alpha A/y^\beta A \rightarrow y^\alpha A/y^\gamma A \rightarrow 0$$

and so we must have

$$\text{Ext}_B^{k-1}(\mathbb{K}_\lambda, y^\gamma A/y^\beta A) \neq 0 \text{ or } \text{Ext}_B^{k-1}(\mathbb{K}_\lambda, y^\alpha A/y^\gamma A) \neq 0.$$

Thus we have $\alpha = \beta + \epsilon_m$, for some $1 \leq m \leq n$. Hence we have $\text{Ext}_B^{k-1}(\mathbb{K}_\lambda, y^{\beta+\epsilon_m} A/y^\beta A) \neq 0$ and so

$$\text{Ext}_B^{k-1}(\mathbb{K}_\lambda \otimes \mathbb{K}_{\beta+\epsilon_m}, \mathbb{K}_{\beta+\epsilon_m} \otimes (y^{\beta+\epsilon_m} A/y^\beta A)) \neq 0.$$

Thus we have $\text{Ext}_B^{k-1}(\mathbb{K}_\nu, A/x_m A) \neq 0$, where $\nu = \lambda + \beta + \epsilon_m$. By the inductive hypothesis, we have

$$\text{Ext}_A^{k-1}(\mathbb{K}_\nu, A/x_m A) = \text{Ext}_B^{k-1}(\mathbb{K}_\nu, A/x_m A) \neq 0$$

and this contradicts Lemma 5.1. Hence we have $R^k \text{pol} A = 0$. Since $R^k \text{pol}$ commutes with direct limits we also have $R^k \text{pol} Z = 0$, where Z is a direct sum of copies of the right regular comodule A . Let X' be any polynomial B -module. Then X' embeds in a direct sum of copies of A , via the comodule structure map. Thus we have a short exact sequence $0 \rightarrow X' \rightarrow Z \rightarrow Y \rightarrow 0$, where Z is a direct sum of copies of

A and Y is a polynomial B -module. Now the derived functors of pol give the exact sequence

$$R^{k-1}\text{pol } Y \rightarrow R^k\text{pol } X' \rightarrow R^k\text{pol } Z = 0.$$

But also, we have $R^{k-1}\text{pol } Y = 0$, from the inductive hypothesis, so that $R^k\text{pol } X' = 0$.

Now for $V', X' \in \text{Comod}(A)$ the 5-term exact sequence (1) gives an isomorphism $\text{Ext}_A^k(V', X') \rightarrow \text{Ext}_B^k(V', X')$. This completes the proof of $P(k+1)$. Hence $P(k)$ is true for all k . Thus we have $R^i\text{pol } X = 0$ for all $i > 0$. This proves (i).

(ii) follows from (i). □

Corollary 5.3. *A B -module is polynomial if and only if all its weights are polynomial.*

Proof. If V is a polynomial B -module then V embeds, via the comodule structure map, into a direct sum of copies of $A(n)$ and it follows that all weights of V are polynomial. To prove that a B -module V with all weights polynomial is polynomial it suffices, by local finiteness, to consider the case in which V is finite dimensional. If V is one dimensional then it is isomorphic to \mathbb{K}_λ , for some $\lambda \in \Lambda(n)$, and hence polynomial. Suppose now that V has dimension bigger than one and let L be a one dimensional submodule. We may assume inductively that V/L is polynomial. We have a natural isomorphism $\text{Ext}_{A(n)}^1(V/L, L) \rightarrow \text{Ext}_B^1(V/L, L)$, by the theorem and it follows that every extension of V/L by L arises from an $A(n)$ -comodule, in particular V is polynomial. □

Let $r \geq 0$. We define the negative (quantised) Borel-Schur algebra $S^-(n, r)$ to be the dual algebra of $\bar{A}(n, r)$. We now obtain the quantised version of a theorem of Woodcock, [Woo92], Theorem 7.

Corollary 5.4. *Let V and X be polynomial B -modules which are homogeneous of degree r . Then we have $\text{Ext}_{S^-(n, r)}^i(V, X) = \text{Ext}_B^i(V, X)$, for all $i \geq 0$.*

Proof. We have

$$\text{Ext}_{S^-(n, r)}^i(V, X) = \text{Ext}_{\bar{A}(n, r)}^i(V, X) = \text{Ext}_{A(n)}^i(V, X) = \text{Ext}_B^i(V, X).$$

□

6 A vanishing theorem for polynomial modules

To save on notation we shall abbreviate $G(n), B(n), T(n)$ to G, B, T where confusion seems unlikely.

We shall need a bound for the vanishing of $R^i \text{ind}_B^G \mathbb{K}_\lambda$, for $\lambda \in \Lambda(n)$. We do this by an inductive argument using the function b that we

now introduce. For each $\lambda \in \Lambda(n)$ we shall define a non-negative integer $b(\lambda)$. We define b on $\Lambda(n, r)$, for $r \geq 0$ by descending induction on the dominance order. If λ is dominant or if $\lambda_j - \lambda_{j+1} = -1$ for some $1 \leq j < n$ we set $b(\lambda) = 0$. In particular this defines $b(\lambda)$ for $\lambda = (r, 0, \dots, 0)$. If $\lambda \in \Lambda(n, r)$ is not of the form already considered then we have $\lambda_j - \lambda_{j+1} = -m_j$, with $m_j \geq 2$, for some $1 \leq j < n$. We define

$$b_j(\lambda) = \max\{b(\lambda + t(\epsilon_j - \epsilon_{j+1})) \mid 0 < t < m_j\} + 1.$$

and

$$b(\lambda) = \min\{b_j(\lambda) \mid 1 \leq j < n, \lambda_j - \lambda_{j+1} \leq -2\}. \quad (2)$$

By an easy induction one sees that if $\lambda = (\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ with $\lambda_1, \dots, \lambda_m \neq 0$ then $b(\lambda) = b(\mu)$, where $\mu = (\lambda_1 - 1, \dots, \lambda_m - 1, 0, \dots, 0)$.

Lemma 6.1. *For $\lambda \in \Lambda(n)$ we have $b(\lambda) \leq \deg(\lambda) - \text{len}(\lambda)$.*

Proof. Since $\Lambda(1)$ consists of dominant weights the result holds for $n = 1$. Suppose that it is false and let n be minimal for which it fails. Let $\lambda \in \Lambda(n)$ be a counterexample of smallest possible degree. If $\lambda = (\lambda_1, \dots, \lambda_m, 0, \dots, 0)$ with $\lambda_1, \dots, \lambda_m \neq 0$ then

$$\begin{aligned} b(\lambda) &= b(\lambda_1 - 1, \dots, \lambda_m - 1, 0, \dots, 0) \\ &\leq \deg(\lambda_1 - 1, \dots, \lambda_m - 1, 0, \dots, 0) \\ &\quad - \text{len}(\lambda_1 - 1, \dots, \lambda_m - 1, 0, \dots, 0) \\ &= \deg(\lambda) - m - \text{len}(\lambda_1 - 1, \dots, \lambda_m - 1, 0, \dots, 0) \\ &\leq \deg(\lambda) - m = \deg(\lambda) - \text{len}(\lambda). \end{aligned}$$

Hence there exists some $1 \leq j < n$ such that $\lambda_j = 0$, $\lambda_{j+1} > 0$. If $\lambda_{j+1} = 1$ then $b(\lambda) = 0$ and λ is not a counterexample. Hence we have $\lambda_{j+1} = m \geq 2$. But now we have

$$\begin{aligned} b(\lambda) &\leq b_j(\lambda) \\ &= \max\{b(\lambda + t(\epsilon_i - \epsilon_{i+1})) \mid 0 < t < m\} + 1. \end{aligned}$$

We consider $\mu = \lambda + t(\epsilon_j - \epsilon_{j+1})$ with $0 < t < m$. Note that μ has entry $t \neq 0$ in the j th position and entry $\lambda_{j+1} - t \geq \lambda_{j+1} - (m - 1) = 1$ in the $(j + 1)$ st position. Moreover, λ and μ agree in all positions other than j and $j + 1$. Hence we have $\text{len}(\mu) = \text{len}(\lambda) + 1$. Moreover, μ is greater than λ , in the dominance order. Hence we have

$$b(\mu) \leq \deg(\mu) - \text{len}(\mu) = \deg(\lambda) - \text{len}(\lambda) - 1$$

i.e., $b(\mu) + 1 \leq \deg(\lambda) - \text{len}(\lambda)$.

Since this is true for all μ of the form $\lambda + t(\epsilon_i - \epsilon_{i+1})$ with $0 < t < m$, from (2), we have $b(\lambda) \leq \deg(\lambda) - \text{len}(\lambda)$. □

Lemma 6.2. (i) *For $\lambda \in \Lambda(n)$ we have $R^i \text{ind}_B^G \mathbb{K}_\lambda = 0$ for all $i > b(\lambda)$, and hence for $i \geq \deg(\lambda) > 0$.*

(ii) *If V is a polynomial B -module of degree r then $R^i \text{ind}_B^G V = 0$ for $i > r$.*

Proof. (i) We argue by induction on $b(\lambda)$. If $b(\lambda) = 0$ then either λ is dominant or $\lambda_j - \lambda_{j+1} = -1$ for some $1 \leq j < n$, and $R^i \text{ind}_B^G \mathbb{K}_\lambda = 0$ for $i > 0$, [Don96], Theorem 3.4 and Lemma 3.1, (ii), and the result holds. So suppose $b(\lambda) > 0$ and the result holds for all $\mu \in \Lambda(n)$ with $b(\mu) < b(\lambda)$. We have $b(\lambda) = b_j(\lambda) + 1$ for some $1 \leq j < n$ with $\lambda_j - \lambda_{j+1} = -m$, $m \geq 2$. Consider the module $\nabla_j(\lambda + (m-1)(\epsilon_j - \epsilon_{j+1}) + \delta)$. Writing $\mu = \lambda + (m-1)(\epsilon_j - \epsilon_{j+1}) + \delta$ we have

$$\mu_j - \mu_{j+1} = -m + 2(m-1) + 1 = m-1.$$

Hence $\nabla_j(\lambda + (m-1)(\epsilon_j - \epsilon_{j+1}) + \delta)$ has weights $\lambda + (m-1)(\epsilon_j - \epsilon_{j+1}) + \delta$, $\lambda + (m-2)(\epsilon_j - \epsilon_{j+1}) + \delta, \dots, \lambda + \delta$, each occurring with multiplicity 1. Hence the module $\nabla_j(\lambda + (m-1)(\epsilon_j - \epsilon_{j+1}) + \delta) \otimes \mathbb{K}_{-\delta}$ has bottom weight λ and we have a short exact sequence of B -modules

$$0 \rightarrow \mathbb{K}_\lambda \rightarrow \nabla_j(\lambda + m(\epsilon_j - \epsilon_{j+1}) + \delta) \otimes \mathbb{K}_{-\delta} \rightarrow Q \rightarrow 0. \quad (3)$$

where Q has weights $\lambda + i(\epsilon_j - \epsilon_{j+1})$, with $1 \leq i \leq m-1$.

Now we have $R^i \text{ind}_B^{P_j} \mathbb{K}_{-\delta} = 0$ for all i , by [Don96], Lemma 3.1 (ii) and so by the tensor identity and [Don96], Proposition 1.3 (iii), we have

$$R^i \text{ind}_B^{P_j} (\nabla_j(\lambda + (m-1)(\epsilon_j - \epsilon_{j+1}) + \delta) \otimes \mathbb{K}_{-\delta}) = 0$$

for all i . By the spectral sequence arising from the transitivity of induction, [Don96], Proposition 1.2, we get $R^i \text{ind}_B^G (\nabla_j(\lambda + m(\epsilon_j - \epsilon_{j+1}) + \delta) \otimes \mathbb{K}_{-\delta}) = 0$ for all i . Hence from (3) we get $R^i \text{ind}_B^G \mathbb{K}_\lambda = R^{i-1} \text{ind}_B^G Q$.

But a weight ν of Q has the form $\lambda + t(\epsilon_j - \epsilon_{j+1})$, with $1 \leq t \leq m-1$, and $b(\nu) \leq b(\lambda) - 1$. So that for $i > b(\lambda)$ we have $i-1 > b(\nu)$ and hence $R^{i-1} \text{ind}_B^G \mathbb{K}_\nu = 0$, by the inductive hypothesis. Since this holds for all weights of Q , i.e., for all composition factors \mathbb{K}_ν of Q , we get $R^{i-1} \text{ind}_B^G Q = 0$, from the long exact sequence, and hence $R^i \text{ind}_B^G \mathbb{K}_\lambda = 0$

(ii) This follows from (i) and the long exact sequence. \square

7 Kempf vanishing for quantised Schur algebras

At this point we introduce the natural left G -module for use later in this section. We write E for the \mathbb{K} -vector space with basis e_1, \dots, e_n . Then E is a G -module via the comodule structure map $\tau: E \rightarrow E \otimes \mathbb{K}[G]$ defined by $\tau(e_i) = \sum_{j=1}^n e_j \otimes c_{ji}$, $1 \leq i \leq n$. We shall also need the symmetric powers $S^r E$ and exterior powers $\bigwedge^r E$ of E . We recall the construction from [DD91] and [Don98]. Let $T(E)$ be the tensor algebra $\bigoplus_{r \geq 0} E^{\otimes r}$. Thus $T(E)$ is a graded \mathbb{K} -algebra, in such a way that each $e_i \in E$ has degree 1. The ideal generated by all $e_i e_j - e_j e_i$, $1 \leq i, j \leq n$, is homogeneous and is a G -submodule, so the (usual) symmetric

algebra $S(E)$ inherits a grading $S(E) = \bigoplus_{r \geq 0} S^r(E)$ and each $S^r(E)$ is a G -submodule of $S(E)$. Also, the ideal of $T(E)$ generated by the elements $e_i^2, e_k e_l + q e_l e_k, 1 \leq i \leq n, 1 \leq k < l \leq n$, is homogeneous and a G -submodule and we write $\Lambda(E)$ for the quotient algebra. Thus $\Lambda(E)$ inherits a grading $\Lambda(E) = \bigoplus_{r \geq 0} \Lambda^r(E)$ and each $\Lambda^r(E)$ is a G -submodule.

For $i = (i_1, \dots, i_r) \in I(n, r)$ we write e_i for $e_{i_1} \otimes \dots \otimes e_{i_r} \in E^{\otimes r}$ and \hat{e}_i for the image of e_i in $\Lambda^r(E)$. The module $\Lambda^r(E)$ has basis \hat{e}_i , with $i \in I(n, r)$, running over all maps with $i_1 > \dots > i_r$.

Proposition 7.1. (i) *Let $1 \leq r \leq n$ and let L_r be the simple B -module with weight ϵ_r . Then for any B -module M the \mathbb{K} -linear map $\phi_M: M \otimes L_r \rightarrow L_r \otimes M$ given by*

$$\phi(m \otimes l) = q^{\alpha_1 + \dots + \alpha_r} l \otimes m$$

for $\alpha = (\alpha_1, \dots, \alpha_n) \in X(n)$ and $m \in M^\alpha$, is a B -module isomorphism.

(ii) *For any B -module M and one dimensional B -module L the B -modules $M \otimes L$ and $L \otimes M$ are isomorphic.*

Proof. (i) Certainly ϕ_M is a linear isomorphism so it remains to show that it is B -module homomorphism. We shall call a B -module M admissible if ϕ_M is a B -module homomorphism. So the point is to show that all B -modules are admissible. Note also that admissibility is preserved by isomorphism. Let M and N be B -modules. Then the map $\phi_{M \otimes N}: M \otimes N \otimes L \rightarrow L \otimes M \otimes N$ factorizes as $(\phi_M \otimes \text{id}_N) \circ (\text{id}_M \otimes \phi_N)$ so that admissibility is preserved under tensor products.

Suppose now that M is a submodule of N . Then the map $\phi_N: N \otimes L_r \rightarrow L_r \otimes N$ restricts to $\phi_M: M \otimes L_r \rightarrow M \otimes L_r$. Thus if N is admissible then so is M . Similarly, if N is admissible then so is the quotient N/M . By the local finiteness of B -modules it suffices to prove that finite dimensional B -modules are admissible.

We now prove that all G -modules are admissible. Note that if M is one dimensional then the twisting map $\theta: M \otimes L_r \rightarrow L_r \otimes M$, given by $\theta(m \otimes l) = (l \otimes m)$, for $l \in L_r, m \in M$, is a B -module map and ϕ_M is a scalar multiple of θ . Hence one dimensional B -modules are admissible. Now if M is any finite dimensional G -module then $D^{\otimes r} \otimes M$ is polynomial for some $r \geq 0$. Hence M is isomorphic to a module of the form $Z \otimes N$, where Z is the dual of $D^{\otimes r}$ and N is polynomial. Hence it suffices to prove that finite dimensional polynomial modules are admissible.

Note that if $M = M_1 \oplus \dots \oplus M_t$, for G -modules M_1, \dots, M_t then $\phi_M = \phi_{M_1} \oplus \dots \oplus \phi_{M_t}$ so that if each M_i is admissible then so is M . Now if M is a polynomial G -module then, for some $r \geq 0$, we may write $M = M_0 \oplus \dots \oplus M_s$, where M_r is polynomial of degree r , for $0 \leq r \leq s$. Hence it suffices to prove that for each r , all polynomial G -modules of degree r are admissible.

We now check that the natural module E is admissible. Let Z be the subspace of $\mathbb{K}[B]$ spanned by the elements $\bar{c}_{ni} \bar{c}_{rr}, 1 \leq i \leq n$. It is seen from the defining relations that Z is also spanned by $\bar{c}_{rr} \bar{c}_{ni}$,

$1 \leq i \leq n$. We fix a non-zero element l_0 of L_r . The subspace Z is a left B -submodule of $\mathbb{K}[B]$ and we have B -module isomorphisms $\theta: E \otimes L_r \rightarrow Z$, $\eta: L_r \otimes E \rightarrow Z$ satisfying $\theta(e_i \otimes l_0) = \bar{c}_{ni}\bar{c}_{rr}$ and $\eta(l_0 \otimes e_i) = \bar{c}_{rr}\bar{c}_{ni}$, for $1 \leq i \leq n$.

Hence we have an isomorphism $\psi = \eta^{-1} \circ \theta: E \otimes L_r \rightarrow L_r \otimes E$. We consider first the case $r = n$. The element \bar{c}_{nn} commutes with the elements \bar{c}_{ni} and $\psi: E \otimes L_n \rightarrow L_n \otimes E$ is the twisting map, taking $e_i \otimes l_0 \rightarrow l_0 \otimes e_i$, for $1 \leq i \leq n$. The map ϕ_E is $q\psi$ and hence is a homomorphism. Now suppose that $r < n$. For $1 \leq i \leq r$ we have $\theta(e_i \otimes l_0) = \bar{c}_{ni}\bar{c}_{rr} = q\bar{c}_{rr}\bar{c}_{ni} = \eta(ql_0 \otimes e_i)$ and hence $\psi(e_i \otimes l_0) = ql_0 \otimes e_i$. For $i > r$ we have $\theta(e_i \otimes l_0) = \bar{c}_{rr}\bar{c}_{ni} = \bar{c}_{ni}\bar{c}_{rr}$ (from the defining relations) and so $\theta(e_i \otimes l_0) = \eta(l_0 \otimes e_i)$. We have shown that the B -module homomorphism $\psi: E \otimes L_r \rightarrow L_r \otimes E$ is given by

$$\psi(e_i \otimes l_0) = \begin{cases} ql_0 \otimes e_i, & \text{if } 1 \leq i \leq r; \\ l_0 \otimes e_i, & \text{if } r < i \leq n. \end{cases}$$

Thus $\psi = \phi_E$ and therefore ϕ_E is a B -module homomorphism.

Since the class of admissible modules is closed under taking tensor products and quotients we get that the j th symmetric power $S^j E$ is admissible for all $j \geq 0$. Now we get that for any $\beta = (\beta_1, \dots, \beta_n) \in \Lambda(n)$ the module $S^\beta E = S^{\beta_1} E \otimes \dots \otimes S^{\beta_n} E$ is admissible. But these modules $S^\beta E$ are injective in the category of polynomial G -modules and every finite dimensional polynomial G -module embeds in a direct sum of copies of the modules $S^\beta E$, [Don98], Section 2.1. Hence every finite dimensional polynomial module is admissible and hence all G -modules are admissible.

The left regular B -module $\mathbb{K}[B]$ is admissible since it is the image of the restriction homomorphism $\mathbb{K}[G] \rightarrow \mathbb{K}[B]$. Hence a direct sum of copies of $\mathbb{K}[B]$ is admissible. Let M be a B -module. Then the structure map $\tau: M \rightarrow M \otimes \mathbb{K}[B]$ embeds M into a direct sum of copies of the left regular B -module and hence M is admissible. This complete the proof of (i).

(ii) We have $L = \mathbb{K}_\lambda$, for some $\lambda \in \Lambda(n)$. If $\lambda = 0$ there is nothing to prove. If $\lambda \neq 0$ and $\lambda \in \Lambda(n)$ we may write $\lambda = \mu + \epsilon_r$, for some $1 \leq r \leq n$ and $\mu \in \Lambda(n)$. Then we have $\mathbb{K}_\lambda \cong \mathbb{K}_\mu \otimes L_r$ so we get

$$\mathbb{K}_\lambda \otimes M \cong \mathbb{K}_\mu \otimes L_r \otimes M \cong \mathbb{K}_\mu \otimes M \otimes L_r$$

by part (i) and now it follows by induction on degree that $\mathbb{K}_\lambda \otimes M$ is isomorphic to $M \otimes \mathbb{K}_\lambda$. Finally, if $\lambda = \mu - \tau$, where $\mu, \tau \in \Lambda(n)$, then we have that

$$\begin{aligned} \mathbb{K}_\tau \otimes \mathbb{K}_\lambda \otimes M &\cong \mathbb{K}_\mu \otimes M \cong M \otimes \mathbb{K}_\mu \\ &\cong M \otimes \mathbb{K}_\tau \otimes \mathbb{K}_\lambda \cong \mathbb{K}_\tau \otimes M \otimes \mathbb{K}_\lambda. \end{aligned}$$

So we have that $\mathbb{K}_\tau \otimes \mathbb{K}_\lambda \otimes M$ is isomorphic to $\mathbb{K}_\tau \otimes M \otimes \mathbb{K}_\lambda$ and tensoring on the left with the dual of \mathbb{K}_τ gives the desired result. \square

Corollary 7.2. *Let $t \geq 1$. Let V_j be a polynomial B -module of degree r_j , for $1 \leq j \leq t$ and let M_j be a G -module for $1 \leq j \leq t+1$. Then we have*

$$R^i \operatorname{ind}_B^G(M_1 \otimes V_1 \otimes \cdots \otimes M_t \otimes V_t \otimes M_{t+1}) = 0$$

for $i > r_1 + \cdots + r_t$.

Proof. By the long exact sequence we may assume that each V_i is one dimensional. Then by the Proposition $M_1 \otimes V_1 \otimes \cdots \otimes M_t \otimes V_t \otimes M_{t+1}$ is isomorphic to $M \otimes V$, where $M = M_1 \otimes \cdots \otimes M_{t+1}$ and $V = V_1 \otimes \cdots \otimes V_t$ and so the result follows from the tensor identity and Lemma 6.2(ii). \square

Remark 7.3. *Recall (or check, by dimension shifting) that if $m \geq 0$ and*

$$0 \rightarrow X_r \rightarrow X_{r-1} \rightarrow \cdots \rightarrow X_0 \rightarrow M \rightarrow 0$$

is an exact sequence of B -modules such that $R^i \operatorname{ind}_B^G X_j = 0$ for all $i > m+j$ then $R^i \operatorname{ind}_B^G M = 0$ for all $i > m$. In particular if $R^i \operatorname{ind}_B^G X_j = 0$ for all $i > j$ then $R^i \operatorname{ind}_B^G M = 0$ for all $i > 0$.

Proposition 7.4. *We have*

$$R^i \operatorname{ind}_B^G \bar{A}(n) = \begin{cases} A(n), & \text{if } i = 0; \\ 0, & \text{if } i > 0. \end{cases}$$

Proof. We first consider the case $i = 0$. The natural map $\pi: A(n) \rightarrow \bar{A}(n)$ gives rise to a G -module map $\tilde{\pi}: A(n) \rightarrow \operatorname{ind}_B^G \bar{A}(n)$, given by

$$\tilde{\pi}(f) = \sum_{i=1}^m \pi(f_i) \otimes f'_i$$

where $\delta_{\mathbb{K}[G]}(f) = \sum_{i=1}^m f_i \otimes f'_i \in A(n) \otimes \mathbb{K}[G]$. Now if $\tilde{\pi}(f) = 0$ then applying $\epsilon_{\mathbb{K}[B]} \otimes \operatorname{id}_{\mathbb{K}[G]}$ we get

$$0 = \sum_{i=1}^m \epsilon_{\mathbb{K}[B]} \pi(f_i) f'_i = \sum_{i=1}^m \epsilon_{\mathbb{K}[G]}(f_i) f'_i = f.$$

Hence $\tilde{\pi}$ is injective. Now the inclusion $\bar{A}(n)$ of $\mathbb{K}[B]$ gives rise to an injective G -module homomorphism $\operatorname{ind}_B^G \bar{A}(n) \rightarrow \operatorname{ind}_B^G \mathbb{K}[B] = \mathbb{K}[G]$. Moreover, $\operatorname{ind}_B^G \bar{A}(n)$ is polynomial, by Remark 4.3, so that this map goes onto $A(n)$. Hence we have a composition of injective G -module homomorphisms

$$A(n) \rightarrow \operatorname{ind}_B^G \bar{A}(n) \rightarrow A(n).$$

But now, restricting to degree r we get an injective homomorphism $A(n, r) \rightarrow A(n, r)$ and since $A(n, r)$ is finite dimensional this map is surjective, for all $r \geq 0$. Hence the composite $A(n) \rightarrow \operatorname{ind}_B^G \bar{A}(n) \rightarrow A(n)$ is surjective and the second map $\operatorname{ind}_B^G \bar{A}(n) \rightarrow A(n)$ is a G -module isomorphism.

We now suppose $i > 0$. Let A_r be the subalgebra of $\bar{A}(n)$ generated by the elements $\bar{c}_{r1}, \dots, \bar{c}_{rr}$. Then A_r is a B -submodule and the multiplication map $A_1 \otimes \dots \otimes A_n \rightarrow \bar{A}(n)$ is a B -module isomorphism.

Let $0 \leq m < n$ and let V_m be the B -submodule of E spanned by e_j , with $m < j \leq n$. Let $V = V_m$. For $r \geq 0$ we write $\bigwedge^r V$ for the subspace of $\bigwedge^r E$ spanned by \hat{e}_i , with $i \in I(n, r)$, $i_a > m$ for $1 \leq a \leq r$. Since this is the image of the B -submodule $V^{\otimes r}$ of $E^{\otimes r}$ under the natural map $E^{\otimes r} \rightarrow \bigwedge^r E$ we have that $\bigwedge^r V$ is a B -submodule of $E^{\otimes r}$. Similarly we have that the ideal J , say, of $S(E)$ generated by V is a B -submodule. We write $S(E/V)$ for the \mathbb{K} -algebra and B -module $S(E)/J$. We write $S^r(E/V)$ for the r th homogeneous component of $S(E/V)$.

Let U_m be the B -submodule of $A(n)$ spanned by $\bar{c}_{m1}, \dots, \bar{c}_{mm}$. Now we have a B -module homomorphism $\theta: E \rightarrow U_m$, sending e_i to \bar{c}_{mi} , for $1 \leq i \leq m$, and to 0 for $m < i \leq n$. Then θ induces a B -module isomorphism $S(E/V_m) \rightarrow A_m$. Hence we have

$$\bar{A}(n, r) \cong \bigoplus_{r=r_1+\dots+r_n} S^{r_1}(E/V_1) \otimes \dots \otimes S^{r_n}(E/V_n). \quad (4)$$

By [Don96], Lemma 3.3(ii), we have that, for $r > 0$, the \mathbb{K} -linear map $\psi: \bigwedge^r E \rightarrow E \otimes \bigwedge^{r-1} E$, given by

$$\psi(\hat{e}_i) = \sum_{a=1}^r (-1)^{a-1} e_{i_a} \otimes \hat{e}_{i_1 \dots \hat{i}_a \dots i_r}$$

for $i \in I(n, r)$ with $i_1 > \dots > i_r$ (where \hat{i}_a indicates that i_a is omitted) is a G -module homomorphism. Combining these maps with the multiplication maps $E \otimes S^b(E) \rightarrow S^{b+1}(E)$, $b \geq 0$, (also G -module maps) in the usual way, for $a \geq 0$, we obtain the Koszul resolution

$$0 \rightarrow \bigwedge^a E \rightarrow \dots \rightarrow \bigwedge^2 E \otimes S^{a-2} E \rightarrow E \otimes S^{a-1} E \rightarrow S^r(E) \rightarrow 0.$$

By restricting the maps in the above we obtain, in the usual way, the Koszul resolution (cf. [Jan03, II 12.12 (i)])

$$\begin{aligned} 0 \rightarrow \bigwedge^a V_m \rightarrow \dots \rightarrow \bigwedge^{a-j} V_m \otimes S^j V_m \\ \rightarrow \dots \rightarrow S^a(E) \rightarrow S^a(E/V_m) \rightarrow 0. \end{aligned}$$

Tensoring all such together, for $1 \leq m \leq n$, we obtain a resolution

$$\dots \rightarrow Y_1 \rightarrow Y_0 \rightarrow S^{r_1}(E/V_1) \otimes \dots \otimes S^{r_n}(E/V_n) \rightarrow 0$$

where each term Y_s is a direct sum of modules of the form $M_1 \otimes Z_1 \otimes \dots \otimes M_t \otimes Z_t$ with each M_i a G -module and each Z_j polynomial of degree d_j , say, $d_1 + \dots + d_t = s$. Now from Corollary 7.2, we have that $R^i \text{ind}_B^G Y_s = 0$ for $i > s$ and hence by Remark 7.3 we have $R^i(S^{r_1}(E/V_1) \otimes \dots \otimes S^{r_n}(E/V_n)) = 0$ for all $i > 0$. Hence by (4) above we have $R^i \text{ind}_B^G \bar{A}(n) = 0$, for all $i > 0$. \square

Theorem 7.5. *Let $\phi: \bar{A}(n) \rightarrow k[B]$ and $\psi: A(n) \rightarrow k[G]$ be the inclusion maps. Let $\pi: A(n) \rightarrow \bar{A}(n)$ be the restriction map. Then, for $V \in \text{Comod}(\bar{A}(n))$, we have $R^i \text{ind}_B^G(\phi_0 V) \cong \psi_0 R^i \pi^0 V$ for all $i \geq 0$.*

Proof. If M is a polynomial B -module then $\text{ind}_B^G M$ is a polynomial G -module, by Remark 4.3. Hence we have $\text{ind}_B^G \circ \phi_0 = \psi_0 \circ \pi^0: \text{Comod}(\bar{A}(n)) \rightarrow \text{Mod}(G)$. We write $F = \text{ind}_B^G \circ \phi_0 = \psi_0 \circ \pi^0$. Now ψ_0 is exact so we have $RF^i V = \psi_0 R^i \pi^0 V$. An injective $\bar{A}(n)$ -comodule is a direct summand of a direct sum of copies of the left regular comodule $\bar{A}(n)$. So it follows from the Proposition 7.4 that ϕ_0 takes injective objects to ind_B^G -acyclic objects. Hence we have a Grothendieck spectral sequence, with second page $R^i \text{ind}_B^G \circ R^j \phi_0 V$ converging to $R^* FV$. But ϕ_0 is exact, so the spectral sequence degenerates and we have $R^i FV \cong R^i \text{ind}_B^G(\phi_0 V) = \psi_0 R^i \pi^0 V$. \square

Remark 7.6. *Slightly less formally, identifying $\text{Comod}(\bar{A}(n))$ with the full subcategory of B -modules whose objects are the polynomial modules and identifying $\text{Comod}(A(n))$ with the subcategory of G -modules whose objects are the polynomial modules, we have $R^i \pi^0 V \cong R^i \text{ind}_B^G V$ for a polynomial B -module V .*

The Theorem 7.5 has the following corollary, generalising Remark 4.3, but which may also be proved by a straightforward dimensional shifting argument.

Corollary 7.7. *If V is a polynomial B -module then $R^i \text{ind}_B^G V$ is a polynomial G -module, for all $i \geq 0$.*

However, the main point of the discussion is to demonstrate the following result, which follows from Kempf's Vanishing Theorem for G , as in [Don96], Theorem 3.4.

Corollary 7.8. *(Kempf Vanishing for polynomial modules.) Let $\pi: A(n) \rightarrow \bar{A}(n)$ be the restriction map. For $\lambda \in \Lambda^+(n)$ we have*

$$R^i \pi^0 \mathbb{K}_\lambda = \begin{cases} \nabla(\lambda), & \text{if } i = 0; \\ 0, & \text{if } i > 0. \end{cases}$$

Let $\pi(r): A(n, r) \rightarrow \bar{A}(n, r)$ be the restriction of π . Now $\pi = \bigoplus_{r=0}^{\infty} \pi(r): A(n) = \bigoplus_{r=0}^{\infty} A(n, r) \rightarrow \bar{A}(n)$. If $V \in \text{Comod}(\bar{A}(n))$ then we may write V uniquely as $V = \bigoplus_{r=0}^{\infty} V(r)$, where $V(r) \in \text{Comod}(\bar{A}(n, r))$ (or less formally, $V(r)$ is polynomial of degree r). It follows that $R^i \pi^0 V = \bigoplus_{r=0}^{\infty} R^i \pi(r)^0 V(r)$. Hence we get:

Corollary 7.9. *(Kempf Vanishing for homogeneous polynomial modules.) Let $r \geq 0$ and let $\pi(n, r): A(n, r) \rightarrow \bar{A}(n, r)$ be the restriction map. For $\lambda \in \Lambda^+(n, r)$ we have*

$$R^i \pi(n, r)^0 \mathbb{K}_\lambda = \begin{cases} \nabla(\lambda), & \text{if } i = 0; \\ 0, & \text{if } i > 0. \end{cases}$$

Let $S(n, r) = A(n, r)^*$ and $S^-(n, r) = \bar{A}(n, r)^*$. Then from Proposition 2.1 we get :

Corollary 7.10. (*Kempf Vanishing for Schur algebras*) For $\lambda \in \Lambda^+(n, r)$ we have

$$\mathrm{Tor}_i^{S^-(n, r)}(\mathbb{K}_\lambda^*, S(n, r)) = \begin{cases} \nabla(\lambda)^*, & \text{if } i = 0; \\ 0, & \text{if } i > 0. \end{cases}$$

(Here \mathbb{K}_λ^* denotes the right $S^-(n, r)$ -dual module of \mathbb{K}_λ .)

8 General coefficient rings

We shall work with Schur algebras over general coefficient rings. We will use the universal coefficient ring $\mathcal{Z} = \mathbb{Z}[t, t^{-1}]$. First we consider the Schur algebra $S_{\mathbb{Q}(t), t}(n, r)$ over the field of rational functions in the parameter t . We define $S_{\mathcal{Z}, t}(n, r)$ to be

$$\{\xi \in S_{\mathbb{Q}(t), t}(n, r) \mid \xi(f) \in \mathcal{Z} \text{ for all } f \in A_{\mathcal{Z}, t}(n, r)\}$$

which, by Lemma 3.1, is a \mathcal{Z} -form of $S_{\mathbb{Q}(t), t}(n, r)$. For an arbitrary commutative ring and a unit q in R we define, by base change via the ring homomorphism from \mathcal{Z} to R , taking t to q , the R -algebra

$$S_{R, q}(n, r) = R \otimes_{\mathcal{Z}} S_{\mathcal{Z}, t}(n, r).$$

It is easy to check that, for R a field and q a unit in R this is consistent with our earlier definition, i.e., that the homomorphism $\mathcal{Z} \rightarrow R$, taking t to q induces an isomorphism $A_{R, q}(n, r)^* \rightarrow R \otimes_{\mathcal{Z}} S_{\mathcal{Z}, t}(n, r)$.

In the same way we define the negative (quantised) Borel-Schur subalgebra $S_{R, q}^-(n, r)$ of $S_{R, q}(n, r)$. We define $S_{\mathbb{Q}(t), t}^-(n, r) = \bar{A}_{\mathbb{Q}(t), t}(n, r)^*$. The coalgebra $\bar{A}_{\mathbb{Q}(t), t}(n, r)$ has a \mathcal{Z} -form $\bar{A}_{\mathcal{Z}, t}(n, r)$ spanned as a \mathcal{Z} -module by the elements $\bar{c}_{ij, \mathbb{Q}(t), t}$, with $i, j \in I(n, r)$. We define $S_{\mathcal{Z}, t}^-(n, r)$ to be

$$\{\xi \in S_{\mathbb{Q}(t), t}^-(n, r) \mid \xi(f) \in \mathcal{Z} \text{ for all } f \in \bar{A}_{\mathcal{Z}, t}(n, r)\}$$

which, by Lemma 3.3 is a \mathcal{Z} -form of $S_{\mathbb{Q}(t), t}^-(n, r)$. For an arbitrary commutative ring and a unit q in R we define, by base change via the ring homomorphism from \mathcal{Z} to R taking t to q , the R -algebra

$$S_{R, q}^-(n, r) = R \otimes_{\mathcal{Z}} S_{\mathcal{Z}, t}^-(n, r).$$

It is easy to check that, for R a field and q a unit in R this is consistent with our earlier definition, i.e., that the homomorphism $\mathcal{Z} \rightarrow R$, taking t to q induces an isomorphism $\bar{A}_{R, q}(n, r)^* \rightarrow R \otimes_{\mathcal{Z}} S_{\mathcal{Z}, t}^-(n, r)$.

The positive Borel-Schur algebra $S_{R, q}^+(n, r)$ is defined in an analogous way. We define $A_{\mathbb{Q}(t), t}^+(n) = A_{\mathbb{Q}(t), t}(n)/I$, where I is the ideal of $A_{\mathbb{Q}(t), t}(n)$ generated by the elements c_{ij} with $1 \leq j < i \leq n$. Then $A_{\mathbb{Q}(t), t}^+(n)$ has a natural coalgebra grading

$$A_{\mathbb{Q}(t), t}^+(n) = \bigoplus_{r \geq 0} A_{\mathbb{Q}(t), t}^+(n, r).$$

For every nonnegative r we define $S_{\mathbb{Q}(t),t}^+(n,r)$ to be the $\mathbb{Q}(t)$ -algebra dual of $A_{\mathbb{Q}(t),t}^+(n,r)$. We write $A_{\mathcal{Z},t}^+(n,r)$ for the image of $A_{\mathcal{Z},t}(n,r)$ under the natural map $A_{\mathbb{Q}(t),t}(n,r) \rightarrow A_{\mathbb{Q}(t),t}^+(n,r)$. Then $A_{\mathcal{Z},t}^+(n,r)$ is a \mathcal{Z} -form of $A_{\mathbb{Q}(t),t}^+(n,r)$ and we define $S_{\mathcal{Z},t}^+(n,r)$ to be

$$\{\xi \in S_{\mathbb{Q}(t),t}^+(n,r) \mid \xi(f) \in \mathcal{Z} \text{ for all } f \in A_{\mathcal{Z},t}^+(n,r)\}.$$

For an arbitrary commutative ring and a unit q in R we define, by base change via the ring homomorphism from \mathcal{Z} to R taking t to q , the R -algebra

$$S_{R,q}^+(n,r) = R \otimes_{\mathcal{Z}} S_{\mathcal{Z},t}^+(n,r).$$

We identify $S_{R,q}^-(n,r)$ and $S_{R,q}^+(n,r)$ with R -subalgebras of $S_{R,q}(n,r)$ in the obvious way.

We now generalise Corollary 7.10 to an arbitrary commutative ground ring from a general result. This is presumably well known but we include it here since we were unable to find a suitable reference. For an algebra S over a commutative ring R and maximal ideal M of R with residue field $\mathbb{K} = R/M$ we write $S_{\mathbb{K}}$ for the \mathbb{K} -algebra $\mathbb{K} \otimes_R S$ obtained by base change. Further, if D is a left (resp. right) S -module we write $D_{\mathbb{K}}$ for the left (resp. right) $S_{\mathbb{K}}$ -module $\mathbb{K} \otimes_R D$ obtained by base change.

Proposition 8.1. *Let R be a commutative Noetherian ring. Let S be an R -algebra which is finitely generated and projective as an R -module. Let D be a right S -module and E a left S -module. Suppose that D and E are finitely generated and projective as R -modules. Suppose further that for each maximal ideal M of R we have*

$$\mathrm{Tor}_i^{S_{\mathbb{K}}}(D_{\mathbb{K}}, E_{\mathbb{K}}) = 0$$

for all $i > 0$ (where $\mathbb{K} = R/M$). Then we have

$$\mathrm{Tor}_i^S(D, E) = 0$$

for all $i > 0$.

Proof. We first make a reduction to the case in which R is local. So we first assume the result in the local case. Let M be a maximal ideal of R and $\mathbb{K} = R/M$. Then we have the R_M algebra S_M obtained by localising at M . The R_M module D_M (resp. E_M) obtained by localisation is naturally a left (resp. right) S_M -module. Also, for $i \geq 0$, we have the localisation $\mathrm{Tor}_i^S(D, E)_M$ of the R -module $\mathrm{Tor}_i^S(D, E)$. Moreover, by (the argument of) [Mat70], (3.E), we have

$$\mathrm{Tor}_i^S(D, E)_M \cong \mathrm{Tor}_i^{S_M}(D_M, E_M) \quad (5)$$

and

$$\mathrm{Tor}_i^{\mathbb{K} \otimes_R S_M}(\mathbb{K} \otimes_R D_M, \mathbb{K} \otimes_R E_M) \cong \mathrm{Tor}_i^{S_{\mathbb{K}}}(D_{\mathbb{K}}, E_{\mathbb{K}}) = 0$$

for $i > 0$. Thus, for $i > 0$, we get $\mathrm{Tor}_i^S(D, E)_M = 0$ for all maximal ideals. Since $\mathrm{Tor}_i^S(D, E)$ is a finitely generated R -module, this implies $\mathrm{Tor}_i^S(D, E) = 0$.

We now assume that R is local with maximal ideal M and $\mathbb{K} = R/M$. We make a reduction to the case $i = 1$. Suppose that $\mathrm{Tor}_1^S(D, E)$ is zero for all D, E as above but that the result is false. We choose $i > 1$ as small as possible such that $\mathrm{Tor}_i^S(D, E) \neq 0$ for some D, E as above. We choose an epimorphism from a finitely generated projective S -module P onto E and consider the corresponding short exact sequence of S -modules

$$0 \rightarrow N \rightarrow P \rightarrow E \rightarrow 0.$$

Then N is finitely generated and projective as an R -module. Hence we have a short exact sequence of $S_{\mathbb{K}}$ -modules

$$0 \rightarrow N_{\mathbb{K}} \rightarrow P_{\mathbb{K}} \rightarrow E_{\mathbb{K}} \rightarrow 0$$

with $P_{\mathbb{K}}$ projective. Hence we have

$$\mathrm{Tor}_j^S(D, N) = \mathrm{Tor}_{j+1}^S(D, E) \quad \text{and} \quad \mathrm{Tor}_j^{S_{\mathbb{K}}}(D_{\mathbb{K}}, N_{\mathbb{K}}) = \mathrm{Tor}_{j+1}^{S_{\mathbb{K}}}(D_{\mathbb{K}}, E_{\mathbb{K}})$$

for $j \geq 1$. So by the minimality of i we have $\mathrm{Tor}_{i-1}^S(D, N) = 0$ and therefore also $\mathrm{Tor}_i^S(D, E) = 0$, a contradiction.

Hence it suffices to prove that $\mathrm{Tor}_1^S(D, E) = 0$ for all D, E satisfying the hypotheses. We now consider the right exact functor \mathcal{F} from the category of finitely generated S -modules to \mathbb{K} -spaces, $\mathcal{F}(X) = X_{\mathbb{K}} \otimes_{S_{\mathbb{K}}} E_{\mathbb{K}}$. Note that \mathcal{F} factorizes: \mathcal{F} is isomorphic to $\mathcal{G} \circ \mathcal{H}$, where \mathcal{H} is a functor from S -modules to $S_{\mathbb{K}}$ -modules, $\mathcal{H}(X) = X_{\mathbb{K}}$ and \mathcal{G} is a functor from the category of $S_{\mathbb{K}}$ -modules to \mathbb{K} -spaces $\mathcal{G}(Y) = Y \otimes_{S_{\mathbb{K}}} E_{\mathbb{K}}$. Moreover, the functors \mathcal{G} and \mathcal{H} are right exact and \mathcal{H} takes projective S -modules to projective $S_{\mathbb{K}}$ -modules. Hence, for $X \in \mathrm{mod}(S)$, there is a Grothendieck spectral sequence with second page $(L_i \mathcal{G} \circ L_j \mathcal{H})X$ converging to $(L_* \mathcal{F})X$. Taking $X = D$, since D is projective as an R -module we have $(L_j \mathcal{H})D = 0$ for all $j > 0$. Hence the spectral sequence degenerates and we have $(L_i \mathcal{F})D = (L_i \mathcal{G})(\mathcal{H}(D))$ for all $i \geq 0$. Hence we have $(L_i \mathcal{F})D = \mathrm{Tor}_i^{S_{\mathbb{K}}}(D_{\mathbb{K}}, E_{\mathbb{K}})$, and from the hypotheses, $(L_i \mathcal{F})D = 0$ for all $i > 0$.

But also, for a right S -module X we have $\mathcal{F}(X) = X_{\mathbb{K}} \otimes_{S_{\mathbb{K}}} E_{\mathbb{K}} = \mathbb{K} \otimes_R (X \otimes_S E)$. This gives another factorisation: \mathcal{F} is the composite $\mathcal{P} \circ \mathcal{Q}$, where \mathcal{Q} is the functor from right S -modules to R -modules, $\mathcal{Q}(X) = X \otimes_S E$ and \mathcal{P} is the functor from R -modules to \mathbb{K} -spaces $\mathcal{P}(Y) = \mathbb{K} \otimes_R Y$. For X projective, $X \otimes_S E$ is a projective R -module. Hence, for X a right S -module, there is a Grothendieck spectral sequence, with second page $(L_i \mathcal{P} \circ L_j \mathcal{Q})X$ converging to $(L_* \mathcal{F})X$. In particular (see [Wei94], Corollary 5.8.4), we have the 5-term exact sequence

$$\begin{aligned} (L_2 \mathcal{F})X &\rightarrow (L_2 \mathcal{P})(\mathcal{Q}(X)) \rightarrow \mathcal{P}(L_1 \mathcal{Q}(X)) \\ &\rightarrow (L_1 \mathcal{F})X \rightarrow (L_1 \mathcal{P})\mathcal{Q}(X) \rightarrow 0. \end{aligned}$$

Taking $X = D$ we obtain the exact sequence

$$\begin{aligned} \mathrm{Tor}_2^{S_{\mathbb{K}}}(D_{\mathbb{K}}, E_{\mathbb{K}}) &\rightarrow \mathrm{Tor}_2^R(\mathbb{K}, D \otimes_S E) \rightarrow \mathbb{K} \otimes_R \mathrm{Tor}_1^S(D, E) \\ &\rightarrow \mathrm{Tor}_1^{S_{\mathbb{K}}}(D_{\mathbb{K}}, E_{\mathbb{K}}) \rightarrow \mathrm{Tor}_1^R(\mathbb{K}, D \otimes_S E) \rightarrow 0. \end{aligned}$$

But $\mathrm{Tor}_i^{S_{\mathbb{K}}}(D_{\mathbb{K}}, E_{\mathbb{K}}) = 0$, for $i > 0$, and so $\mathrm{Tor}_1^R(\mathbb{K}, D \otimes_S E) = 0$. Hence $D \otimes_R E$ is a projective R -module, see [Mat70], Section 18, Lemma 4. Hence $\mathrm{Tor}_2^R(\mathbb{K}, D \otimes_S E) = 0$ and hence $\mathbb{K} \otimes_R \mathrm{Tor}_1^S(D, E) = 0$, and hence $\mathrm{Tor}_1^S(D, E) = 0$. \square

Let R be a commutative ring with Noetherian subring R_0 . Let $\phi: X \rightarrow Y$ be an R_0 -module homomorphism. It is easy to check (and we leave this to the reader) that if for every subring R' of R containing R_0 which is finitely generated over R_0 , the R' -module homomorphism $\phi_{R'}: R' \otimes_{R_0} X \rightarrow R' \otimes_{R_0} Y$ is injective then the R -module homomorphism $\phi_R: X_R \rightarrow Y_R$ (obtained by base change) is injective.

Lemma 8.2. *Let R be a commutative ring and let R_0 be a Noetherian subring. Let S be an R_0 -algebra, finitely generated and projective as an R_0 -module. Let M be a right S -module and N a left S -module and suppose that M and N are finitely generated and projective over R_0 . If $\mathrm{Tor}_i^{S_{R'}}(M_{R'}, N_{R'}) = 0$ for all $i > 0$, and all subrings R' of R containing R_0 and finitely generated over R_0 , then $\mathrm{Tor}_i^{S_R}(M_R, N_R) = 0$ for all $i > 0$.*

Proof. Choose an S -module surjection $P \rightarrow N$, where P is a finitely projective S -module and let

$$0 \rightarrow H \rightarrow P \rightarrow N \rightarrow 0 \quad (6)$$

be the corresponding short exact sequence. Then we have that $M_{R'} \otimes_{S_{R'}} N_{R'} \rightarrow H_{R'} \otimes_{S_{R'}} P_{R'}$, is injective, i.e.,

$$R' \otimes_{R_0} (M \otimes_{R_0} H) \rightarrow R' \otimes_{R_0} (M \otimes_{R_0} P)$$

is injective (whenever R' is a subring of R finitely generated over R_0 , since $\mathrm{Tor}_1^{S_{R'}}(M_{R'}, N_{R'}) = 0$). Hence we have that

$$R \otimes_{R_0} (M \otimes_{R_0} H) \rightarrow R \otimes_{R_0} (M \otimes_{R_0} P)$$

is injective, i.e., $M_R \otimes_{S_R} N_R \rightarrow H_R \otimes_{S_R} P_R$ is injective and therefore $\mathrm{Tor}_1^{S_R}(M_R, N_R) = 0$. Now for $i > 1$ it follows that $\mathrm{Tor}_i^{S_R}(M_R, N_R) = 0$ using (6) and dimension shifting. \square

Let $\lambda \in \Lambda(n, r)$. Then we have the one dimensional module \mathbb{K}_λ for the quantised Borel subgroup $B(n)$ over $\mathbb{K} = \mathbb{Q}(t)$. Thus \mathbb{K}_λ is naturally a left $S_{\mathbb{Q}(t), t}^-(n, r)$ -module and we obtain an $S_{R, q}^-(n, r)$ -module R_λ , free of rank one over R , by base change. We write R_λ^* for the right $S_{R, q}(n, r)$ -module dual of R_λ . Similarly we construct an $S_{R, q}^+(n, r)$ -module, also denoted R_λ , free of rank one over R .

Theorem 8.3. *Let R be a commutative ring and let q be a unit in R . Let $\lambda \in \Lambda^+(n, r)$. Then we have*

$$\mathrm{Tor}_i^{S_{R,q}^-(n,r)}(R_\lambda^*, S_{R,q}(n, r)) = 0$$

for all $i > 0$.

Proof. The result for R Noetherian follows from Corollary 7.10 and Proposition 8.1 and the result for general R follows from Lemma 8.2. \square

We shall also give the version of this result for the positive Borel-Schur algebra. Suppose J is an anti-automorphism of a ring S and that S has subrings S^- and S^+ interchanged by J . Given a right S^- -module (resp. left) M we write M^J for the same group M regarded as a left (resp. right) S^+ -module with action $xm = mJ(x)$ (resp. $mx = J(x)m$), $m \in M$, $x \in S^+$. Note that if M is S , regarded as a right S^- -module via right multiplication, then M^J is S regarded as a left S^+ -module via left multiplication. Similarly if M is S , regarded as a left S^- -module via left multiplication, then M^J is S regarded as a right S^+ -module via right multiplication. If M is a right S^- -module and N is a left S^- -module then we have

$$\mathrm{Tor}_i^{S^-}(M, N) \cong \mathrm{Tor}_i^{S^+}(N^J, M^J)$$

for all $i \geq 0$.

Recall that, by [Don98], pg. 82, we have an involutory anti-automorphism J of the Schur algebra $S_{\mathbb{K},q}(n, r)$ over a field \mathbb{K} . For $i \in I(n, r)$ we write $d(i)$ for the number of pairs (a, b) such that $1 \leq a < b \leq r$ and $i_a < i_b$. Then, for $i, j \in I(n, r)$, we have

$$c_{ji}(\xi)q^{d(j)} = c_{ij}(J(\xi))q^{d(i)}$$

(see [Don98], p83) and clearly J is determined by this property. Taking $\mathbb{K} = \mathbb{Q}(t)$ and $q = t$, it is easy to check that J preserves $S_{\mathbb{Z},t}(n, r)$ and interchanges $S_{\mathbb{Z},t}^-(n, r)$ and $S_{\mathbb{Z},t}^+(n, r)$. Hence J induces, for a general commutative ring R and unit $q \in R$, an anti-automorphism, which we also denote J , of $S_{R,q}(n, r)$ which interchanges $S_{R,q}^-(n, r)$ and $S_{R,q}^+(n, r)$. Moreover, for $\lambda \in \Lambda(n, r)$, starting with the left $S_{R,q}^-(n, r)$ -module R_λ , we have that $(R_\lambda^*)^J$ is the left $S_{R,q}^+(n, r)$ -module, also denoted by R_λ . Thus from Theorem 8.3 we get our quantised version of Woodcock's Theorem.

Theorem 8.4. *Let R be a commutative ring and let q be a unit in R . Let $\lambda \in \Lambda^+(n, r)$. Then we have*

$$\mathrm{Tor}_i^{S_{R,q}^+(n,r)}(S_{R,q}(n, r), R_\lambda) = 0$$

for all $i > 0$.

Recall that $A_{\mathbb{Q}(t),t}(n, r)$ has basis $\{c_{ij} \mid (i, j) \in Y(n, r)\}$. Let

$$\{\xi_{ij} \mid (i, j) \in I(n, r)\}$$

be the dual basis of $S_{\mathbb{Q}(t),t}(n, r)$, that is

$$\xi_{ij}(c_{i',j'}) = \begin{cases} 1, & i = i', j = j' \\ 0, & (i', j') \in Y(n, r), (i, j) \neq (i', j'). \end{cases}$$

Then it is straightforward that $\{\xi_{ij} \mid (i, j) \in Y(n, r)\}$ is a \mathcal{Z} -basis of $S_{\mathbb{Z},t}(n, r)$. We will also denote by the same symbol ξ_{ij} the image of ξ_{ij} in $S_{R,q}(n, r)$ under base change. For $\lambda \in \Lambda(n, r)$, we will write ξ_λ for $\xi_{l(\lambda),l(\lambda)}$.

Note that in Section 2 of [Don98] there is used a slightly different parameterisation of the set $\{\xi_{ij} \mid (i, j) \in Y(n)\}$. As we will refer the results of [Don98], we will explain this in more detail. Let U be the subset of $I(n, r) \times I(n, r)$ of pairs $(l(\lambda), j)$ such that $j_1 \geq \dots \geq j_{\lambda_1}$, $j_{\lambda_1+1} \geq \dots \geq j_{\lambda_1+\lambda_2}$ and so on. Now for every $\lambda \in \Lambda(n, r)$, there is a permutation $\pi_\lambda \in \Sigma_\lambda$ of order 2 such that $(l(\lambda), j) \in Y(n, r)$ if and only if $(l(\lambda), j\pi_\lambda) \in U$. Since the generators c_{ab} and $c_{ab'}$ commute for any b and b' , we have $c_{l(\lambda),j} = c_{l(\lambda),j\pi_\lambda}$. Thus we get for any $(i, j) \in Y(n, r)$ and $(i', j') \in U$ that

$$\xi_{ij}(c_{i',j'}) = \xi_{ij}(c_{i',j'\pi_\lambda}) = \begin{cases} 1, & i = i', j = j'\pi_\lambda \\ 0, & \text{otherwise.} \end{cases}$$

Therefore ξ_{ij} in our notation corresponds to $\xi_{i,j\pi_\lambda}$ in the notation of [Don98].

Using the above identification, from [Don98], page 38, for $\lambda, \mu \in \Lambda(n, r)$ we obtain

$$\xi_\lambda \xi_{ij} \xi_\mu = \begin{cases} \xi_{ij}, & i \in \lambda, j \in \mu, \\ 0, & \text{otherwise.} \end{cases}$$

Moreover, $1 = \sum_{\lambda \in \Lambda(n, r)} \xi_\lambda$ is an orthogonal idempotent decomposition of the identity.

Similarly to Lemma 3.2, we have that the kernel of the projection

$$f: A_{\mathbb{Q}(t),t}(n, r) \twoheadrightarrow A_{\mathbb{Q}(t),t}^+(n, r)$$

has basis

$$\{c_{ij} \mid (i, j) \in Y(n, r) \text{ but not } i \leq j\}.$$

Thus for any $(i, j) \in Y(n, r)$ such that $i \leq j$, we get that the restriction of ξ_{ij} to $\text{Ker}(f)$ is zero. Therefore we can consider ξ_{ij} as an element of $S_{\mathbb{Q}(t),t}^+(n, r) = A_{\mathbb{Q}(t),t}^+(n, r)^*$. Using a dimension argument we get that

$$\{\xi_{ij} \mid (i, j) \in Y(n, r), i \leq j\} \quad (7)$$

is a $\mathbb{Q}(t)$ -basis of $S_{\mathbb{Q}(t),t}^+(n, r)$. Obviously, it is also a \mathcal{Z} -basis of $S_{\mathbb{Z},t}^+(n, r)$ and, by base change, an R -basis for any $S_{R,q}^+(n, r)$.

We now recall the normalised bar construction. Note that this is a special case of the construction described in Chapter IX, §7 of [ML63] and its detailed treatment can be found in Section 3 of [SY12].

Let S be a ring with identity and S' a subring of S . We assume that there is an epimorphism of rings $p: S \rightarrow S'$ that splits the natural inclusion of S' into S . Write \mathcal{K} for the kernel of p . Then \mathcal{K} is an S' -bimodule.

For every S -module M we define the chain complex

$$B_*(S, S', M) = (B_k(S, S', M), d_k)_{k \geq -1}$$

as follows:

$$B_{-1}(S, S', M) = M, \quad B_0(S, S', M) = S \otimes M,$$

$$B_k(S, S', M) = S \otimes \mathcal{K}^{\otimes k} \otimes M, \quad \forall k \geq 1,$$

$$d_k = \sum_{t=0}^k (-1)^t d_{kt}: B_k(S, S', M) \rightarrow B_{k-1}(S, S', M),$$

where all the tensor products are over S' and the S -module homomorphisms $d_{kt}: B_k(S, S', M) \rightarrow B_{k-1}(S, S', M)$, $k \geq 0$, $0 \leq t \leq k$ are given by

$$d_{00}(s \otimes m) = sm$$

$$d_{k0}(s \otimes s_1 \otimes \cdots \otimes s_k \otimes m) = ss_1 \otimes s_2 \otimes \cdots \otimes s_k \otimes m$$

$$d_{kt}(s \otimes s_1 \otimes \cdots \otimes s_k \otimes m) = s \otimes \cdots \otimes s_t s_{t+1} \otimes \cdots \otimes m, \quad 1 \leq t \leq k-1$$

$$d_{kk}(s \otimes s_1 \otimes \cdots \otimes s_k \otimes m) = s \otimes s_1 \otimes \cdots \otimes s_{k-1} \otimes s_k m, \quad k \geq 0.$$

The complex $(B(S, S', M), d)$ is exact and is called the *normalised bar resolution* of M over S . Now we specialize this construction to the case of the quantised Borel-Schur algebra.

Define

$$L = L_{R,q} = \bigoplus_{(i,i) \in Y(n,r)} R\xi_{ii} = \bigoplus_{\lambda \in \Lambda(n,r)} R\xi_\lambda$$

and

$$\mathcal{J} = \mathcal{J}_{R,q} = \bigoplus_{\substack{(i,j) \in Y(n,r) \\ i < j}} R\xi_{ij}.$$

Then $L \oplus \mathcal{J} = S_{R,q}^+(n, r)$.

Proposition 8.5. *The R -module $L_{R,q}$ is a split subalgebra of $S_{R,q}^+(n, r)$ and $\mathcal{J}_{R,q}$ is a split ideal of $S_{R,q}^+(n, r)$.*

Proof. It is obvious that $L_{R,q}$ is a subalgebra of $S_{R,q}^+(n, r)$.

Now, we will check that $\mathcal{J}_{R,q}$ is an ideal of $S_{R,q}^+(n, r)$. By a base change argument, it is enough to check that $\mathcal{J}_{\mathcal{Z},t}$ is an ideal of $S_{\mathcal{Z},t}^+(n, r)$ and this can be reduced to showing that $\mathcal{J}_{\mathbb{Q}(t),t}(n, r)$ is an ideal of $S_{\mathbb{Q}(t),t}^+(n, r)$.

Let $(i, j), (i', j') \in Y(n, r)$ such that $i \leq j$ and $i' \leq j'$. Then the coefficient of ξ_λ in the expansion of the product $\xi_{i,j}\xi_{i',j'}$ in the basis (7) of $S_{\mathbb{Q}(t),t}^+(n, r)$ is given by

$$(\xi_{i,j}\xi_{i',j'}) (\bar{c}_{l(\lambda),l(\lambda)}) = \sum_{h \in I(n, r)} \xi_{i,j} (\bar{c}_{l(\lambda),h}) \xi_{i',j'} (\bar{c}_{h,l(\lambda)}),$$

where \bar{c}_{ij} denotes the image of c_{ij} under the epimorphism

$$A_{\mathbb{Q}(t),t}(n, r) \rightarrow A_{\mathbb{Q}(t),t}^+(n, r).$$

Thus $\bar{c}_{l(\lambda),h} \neq 0$ and $\bar{c}_{h,l(\lambda)} \neq 0$ imply that $l(\lambda) \leq h \leq l(\lambda)$. Hence

$$(\xi_{i,j}\xi_{i',j'}) (\bar{c}_{l(\lambda),l(\lambda)}) = \xi_{i,j} (\bar{c}_{l(\lambda),l(\lambda)}) \xi_{i',j'} (\bar{c}_{l(\lambda),l(\lambda)}),$$

which is zero, if $i \neq j$ or $i' \neq j'$. This proves that the product $\xi_{i,j}\xi_{i',j'}$ lies in $\mathcal{J}_{\mathbb{Q}(t),t}$, if $\xi_{i,j} \in \mathcal{J}_{\mathbb{Q}(t),t}$ or $\xi_{i',j'} \in \mathcal{J}_{\mathbb{Q}(t),t}$. \square

For any $\lambda \in \Lambda(n, r)$ we can apply the normalised bar construction to $S_{R,q}^+(n, r)$, L , and the rank-one module R_λ .

Denote $B_k(S_{R,q}^+(n, r), \mathcal{J}, R_\lambda)$ by $B_{k,\lambda}^+$ for $k \geq -1$. We get

$$B_{-1,\lambda}^+ = R_\lambda, \quad B_{0,\lambda}^+ = S_{R,q}^+(n, r) \otimes_L R_\lambda,$$

$$B_{k,\lambda}^+ = S_{R,q}^+(n, r) \otimes_L \mathcal{J}^{\otimes k} \otimes_L R_\lambda, \quad k \geq 1.$$

For any $\mu \in \Lambda(n, r)$, $M \in \text{mod-}L$ and $N \in L\text{-mod}$, we have

$$(M \otimes_L R\xi_\mu) \otimes_R (R\xi_\mu \otimes_L N) \cong M \otimes_L R\xi_\mu \otimes_L N.$$

Thus

$$\begin{aligned} M \otimes_L N &\cong M \otimes_L L \otimes_L N \cong \bigoplus_{\mu \in \Lambda} (M \otimes_L R\xi_\mu) \otimes_R (R\xi_\mu \otimes_L N) \\ &\cong \bigoplus_{\mu \in \Lambda} M\xi_\mu \otimes_R \xi_\mu N, \end{aligned}$$

since $M \otimes_L R\xi_\mu \cong M\xi_\mu$ and $R\xi_\mu \otimes_L N \cong \xi_\mu N$. Hence

$$\begin{aligned} B_{0,\lambda}^+ &\cong \bigoplus_{\mu \in \Lambda} S_{R,q}^+(n, r) \xi_\mu \otimes_R \xi_\mu R_\lambda \\ &= S_{R,q}^+(n, r) \xi_\lambda \otimes_R R_\lambda \cong S_{R,q}^+(n, r) \xi_\lambda, \end{aligned}$$

since $\xi_\mu R_\lambda = 0$ unless $\mu = \lambda$. Further

$$\begin{aligned} B_{k,\lambda}^+ &\cong \bigoplus_{\mu^{(1)}, \dots, \mu^{(k+1)} \in \Lambda} S_{R,q}^+(n, r) \xi_{\mu^{(1)}} \otimes_R \xi_{\mu^{(1)}} \mathcal{J} \xi_{\mu^{(2)}} \otimes_R \dots \\ &\quad \otimes_R \xi_{\mu^{(k)}} \mathcal{J} \xi_{\mu^{(k+1)}} \otimes_R \xi_{\mu^{(k+1)}} R_\lambda. \end{aligned}$$

As $\{\xi_{ij} \mid (i, j) \in Y(n, r), i < j\}$ is an R -basis of \mathcal{J} , we get that $\xi_\mu \mathcal{J} \xi_\tau$ is zero, unless $\mu \triangleright \tau$. If $\mu \triangleright \tau$, then $\xi_\mu \mathcal{J} \xi_\tau$ has an R -basis

$$\{\xi_{ij} \mid (i, j) \in Y(n, r), i < j, j \in \tau, i \in \mu\}.$$

Thus for every $k \geq 1$ we can write

$$B_{k,\lambda}^+ \cong \bigoplus_{\mu^{(1)} \triangleright \dots \triangleright \mu^{(k)} \triangleright \lambda} S_{R,q}^+(n,r) \xi_{\mu^{(1)}} \otimes \xi_{\mu^{(1)}} \mathcal{J} \xi_{\mu^{(2)}} \otimes \dots \otimes \xi_{\mu^{(k)}} \mathcal{J} \xi_{\lambda},$$

where \otimes means \otimes_R . Note that $B_{k,\lambda}^+$ is zero for k sufficiently large.

Given $\mu \triangleright \lambda$, we define $\Omega_k^+(\lambda, \mu)$ to be the set of all sequences

$$(i^{(1)}, j^{(1)}), \dots, (i^{(k)}, j^{(k)})$$

of elements in $Y(n, r)$ such that $i^{(1)} \in \mu$, $j^{(k)} \in \lambda$, and

$$i^{(1)} < j^{(1)} \sim i^{(2)} < j^{(2)} \sim i^{(3)} < \dots < j^{(k)},$$

where $j \sim i$ means that i and j have the same content. Then we have isomorphisms of $S_{R,q}^+(n, r)$ -modules

$$B_{k,\lambda}^+ \cong \begin{cases} S_{R,q}^+(n, r) \xi_{\lambda}, & k = 0 \\ \bigoplus_{\mu \triangleright \lambda} (S_{R,q}^+(n, r) \xi_{\mu})^{\#\Omega_k^+(\lambda, \mu)}, & k \geq 1. \end{cases} \quad (8)$$

So we get the following result

Theorem 8.6. *Let $\lambda \in \Lambda(n, r)$. Then the complex $B_{*,\lambda}^+$ is a projective resolution of R_{λ} over $S_{R,q}^+(n, r)$.*

Now consider $\lambda \in \Lambda^+(n, r)$. The *Weyl module* associated with λ is

$$W_{\lambda} = S_{R,q}(n, r) \otimes_{S_{R,q}^+(n, r)} R_{\lambda}.$$

By Theorem 8.4, R_{λ} is an acyclic module for the functor $S_{R,q} \otimes_{S_{R,q}^+(n, r)} -$. Therefore $B_{*,\lambda} := S_{R,q}(n, r) \otimes_{S_{R,q}^+(n, r)} B_{*,\lambda}^+$ is a projective resolution of W_{λ} . Moreover, since

$$S_{R,q}(n, r) \otimes_{S_{R,q}^+(n, r)} S_{R,q}^+(n, r) \cong S_{R,q}(n, r),$$

we get the following theorem.

Theorem 8.7. *Let $\lambda \in \Lambda^+(n, r)$. Define the complex B_{λ} as follows:*

$$B_{-1,\lambda} = W_{\lambda}, \quad B_{0,\lambda} = S_{R,q}(n, r) \xi_{\lambda},$$

and for $k \geq 1$, we set $B_{k,\lambda}$ to be

$$\bigoplus_{\substack{\mu^{(1)} \triangleright \dots \triangleright \mu^{(k)} \triangleright \lambda \\ \mu^{(1)}, \dots, \mu^{(k)} \in \Lambda(n, r)}} S_{R,q}(n, r) \xi_{\mu^{(1)}} \otimes_R \xi_{\mu^{(1)}} \mathcal{J} \xi_{\mu^{(2)}} \otimes_R \dots \otimes_R \xi_{\mu^{(k)}} \mathcal{J} \xi_{\lambda}.$$

Define d_0 to be the canonical projection of $S_{R,q}(n, r) \xi_{\lambda}$ on W_{λ} , and for $k \geq 1$ define $d_k: B_{k,\lambda} \rightarrow B_{k-1,\lambda}$ to be the R -linear extension of the map

$$x_0 \otimes x_1 \cdots \otimes x_k \mapsto \sum_{t=0}^{k-1} (-1)^t x_0 \otimes \cdots \otimes x_t x_{t+1} \otimes \cdots \otimes x_k.$$

Then B_{λ} is a projective resolution of W_{λ} over $S_{R,q}(n, r)$.

We will show now that B_λ is stable under base change. Our resolutions for a moment will get additional indices to emphasize dependence on R and $q \in R$. Let R and R' be commutative rings, $\theta: R \rightarrow R'$ a ring homomorphism, $q \in R$ and $q' := \theta(q) \in R'$ invertible elements. Since $S_{R,q}(n, r) \otimes_R R' \cong S_{R',q}(n, r)$ and $B_{k,\lambda}^R$ are free $S_{R,q}(n, r)$ -modules for $k \geq 0$, we get that $(B_{k,\lambda}^R \otimes_R R', k \geq 0)$ and $(B_{k,\lambda}^{R'}, k \geq 0)$ are isomorphic complexes. Moreover, from the commutative diagram with exact rows

$$\begin{array}{ccccccc} B_{1,\lambda}^R \otimes_R R' & \xrightarrow{d_1} & B_{0,\lambda}^R \otimes_R R' & \xrightarrow{d_0} & W_\lambda^R \otimes_R R' & \longrightarrow & 0 \\ \cong \downarrow & & \cong \downarrow & & \cong \downarrow \exists! & & \\ B_{1,\lambda}^{R'} & \xrightarrow{d_1} & B_{0,\lambda}^{R'} & \xrightarrow{d_0} & W_\lambda^{R'} & \longrightarrow & 0, \end{array}$$

it follows that $B_{*,\lambda}^R \otimes_R R'$ and $B_{*,\lambda}^{R'}$ are isomorphic also in degree -1 .

9 The Hecke algebra and resolutions of co-Specht modules

In this section we will use the notation of [DJ86] but will denote by $\text{lng}(\sigma)$ the length of $\sigma \in \Sigma_r$. The *Hecke algebra* $\mathcal{H} = \mathcal{H}_{R,q}$ associated with Σ_r over R is free as an R -module with basis $\{T_\sigma \mid \sigma \in \Sigma_r\}$, where

$$T_s T_\sigma = \begin{cases} T_{s\sigma}, & \text{if } \text{lng}(s\sigma) = \text{lng}(\sigma) + 1 \\ qT_{s\sigma} + (q-1)T_\sigma, & \text{otherwise,} \end{cases}$$

for $\sigma, s \in \Sigma_r$ with $\text{lng}(s) = 1$.

In [BM12] Boltje and Maisch constructed for every composition λ of r a chain complex \widetilde{C}_*^λ of \mathcal{H} -modules. These complexes are lifting to the q -setting of the corresponding $R\Sigma_r$ -module complexes described in [BH11]. It was proved in [SY12], that \widetilde{C}_*^λ is a permutation resolution of the co-Specht modules $\text{Hom}_R(S^\lambda, R)$ for $q = 1$ and λ a partition of r . In this section we will prove a similar result for any invertible q in R .

Choose any $n \geq r$, and let

$$\begin{aligned} \omega &= (1, \dots, 1, 0, \dots, 0) \in \Lambda(n, r) \\ u &= (1, 2, \dots, r) \in I(n, r). \end{aligned}$$

Then (see [Don98, Section 0.23])

$$\{\xi_{u,u\pi} \mid \pi \in \Sigma_r\}$$

is an R -basis of $\xi_\omega S_{R,q}(n, r) \xi_\omega$ and

$$\begin{aligned} \xi_\omega S_{R,q}(n, r) \xi_\omega &\rightarrow \mathcal{H} \\ \xi_{u,u\pi} &\mapsto T_{\pi^{-1}} \end{aligned}$$

is an isomorphism of R -algebras. Therefore we have the Schur functor

$$\begin{aligned} F: S_{R,q}(n, r)\text{-mod} &\rightarrow \mathcal{H}\text{-mod} \\ V &\mapsto \xi_\omega V. \end{aligned}$$

Applying F to the resolution B_λ of W_λ with $\lambda \in \Lambda^+(n, r)$, we obtain an exact sequence $F(B_\lambda)$. It is our aim to prove that $F(B_\lambda)$ and \widetilde{C}_*^λ are isomorphic chain complexes of \mathcal{H} -modules. This will prove that the complexes \widetilde{C}_*^λ are resolutions of the co-Specht modules $\text{Hom}_R(S^\lambda, R)$ over \mathcal{H} .

We start by reminding to the reader some facts on Hecke algebra. Denote by Σ_λ the standard Young subgroup corresponding to the composition λ . By [DJ86, Lemma 1.1] each right coset of Σ_λ in Σ_r contains a unique element of minimal length, the *distinguished* coset representative of Σ_λ in Σ_r . We denote by D_λ the set of these elements. Given two compositions λ and μ , we also define $D_{\lambda,\mu} = D_\lambda \cap D_\mu^{-1}$. By [DJ86, Lemma 1.6] the set $D_{\lambda,\mu}$ is a system of Σ_λ - Σ_μ double coset representatives in Σ_r .

Recall that, for $\lambda \in \Lambda(n, r)$, we write $l(\lambda)$ for the multi-index $(1^{\lambda_1}, 2^{\lambda_2}, \dots, n^{\lambda_n})$. Then every element of $Y(n, r)$ is of the form $(l(\lambda), j)$ for some $\lambda \in \Lambda(n, r)$ and $j \in I(n, r)$. It is easy to see (cf. [DD91, pp. 24-25]), that for given $\lambda, \mu \in \Lambda(n, r)$, there is a bijective correspondence

$$\{(l(\lambda), j) \in Y(n, r) \mid j \in \mu\} \rightarrow D_{\lambda,\mu}, \quad (9)$$

defined as follows. For a given pair $(l(\lambda), j)$ the set

$$\{\pi \in \Sigma_r \mid l(\mu)\pi^{-1} = j\}$$

is a Σ_μ -orbit, and thus contains a unique distinguished element \bar{d} of D_μ^{-1} . We define d as the representative of $\Sigma_\lambda \bar{d} \Sigma_\mu$ in $D_{\lambda,\mu}$.

For $\lambda \in \Lambda(n, r)$, define $x_\lambda := \sum_{\pi \in \Sigma_\lambda} T_\pi$ and $M^\lambda := x_\lambda \mathcal{H}$. Then $\text{Hom}_{\mathcal{H}}(M^\mu, M^\lambda)$ has an R -basis

$$\{\varphi_d^{\lambda,\mu} \mid d \in D_{\lambda,\mu}\},$$

where

$$\varphi_d^{\lambda,\mu}(x_\mu) = \sum_{\pi \in \Sigma_\lambda d \Sigma_\mu} T_\pi, \quad d \in D_{\lambda,\mu}.$$

Theorem 3.2.5 and Corollary 3.2.6 in [DD91] say that there is an algebra isomorphism

$$S_{R,q}(n, r) \rightarrow \bigoplus_{\mu, \lambda \in \Lambda(n, r)} \text{Hom}_{\mathcal{H}}(M^\mu, M^\lambda) \quad (10)$$

$$\xi_{l(\lambda), j} \mapsto \varphi_d^{\lambda,\mu},$$

where the correspondence $(l(\lambda), j) \mapsto d$ is given by (9).

Denote by $\mathcal{T}(\lambda, \mu)$ the set of all λ -tableaux with content μ and by $\mathcal{T}^{rs}(\lambda, \mu)$ the set of all row semistandard λ -tableaux with content μ . Write

$$T^\lambda = \begin{array}{ccccccc} & 1 & & 2 & \dots & \lambda_1 & \\ & \lambda_1 + 1 & & \lambda_1 + 2 & \dots & \dots & \dots & \lambda_1 + \lambda_2 \\ & \dots & & & & & & \\ \lambda_1 + \dots + \lambda_{n-1} + 1 & & \dots & & \dots & \dots & & r \end{array}$$

and for each $i \in I(n, r)$, let T_i^λ be the λ -tableaux

$$T_i^\lambda = \begin{array}{ccccccc} & i_1 & & i_2 & \dots & i_{\lambda_1} & \\ & i_{\lambda_1+1} & & i_{\lambda_1+2} & \dots & \dots & \dots & i_{\lambda_1+\lambda_2} \\ & \dots & & & & & & \\ i_{\lambda_1+\dots+\lambda_{n-1}+1} & & \dots & & \dots & \dots & & i_r \end{array}$$

Recall that $(i, j) \in Y(n, r)$ if and only if $i_1 \leq i_2 \leq \dots \leq i_r$ and $j_\nu \leq j_{\nu+1}$ if $i_\nu = i_{\nu+1}$, $1 \leq \nu \leq r-1$. Therefore there is a bijective correspondence

$$\begin{aligned} \{(l(\lambda), j) \in Y(n, r) \mid j \in \mu\} &\rightarrow \mathcal{T}^{rs}(\lambda, \mu) \\ (l(\lambda), j) &\mapsto T_j^\lambda \end{aligned}$$

that in combination with (9) induces the bijection

$$D_{\lambda, \mu} \leftrightarrow \mathcal{T}^{rs}(\lambda, \mu). \quad (11)$$

Boltje and Maisch say that a λ -tableaux in $\mathcal{T}(\lambda, \mu)$ is *ascending* if, for every $a \in \mathbb{N}$, the a th row of this tableau contains only entries which are greater than or equal to a . They denote the set of all ascending elements of $\mathcal{T}^{rs}(\lambda, \mu)$ by $\mathcal{T}^\wedge(\lambda, \mu)$. One has $\mathcal{T}^\wedge(\lambda, \mu) \neq \emptyset$ if and only if $\mu \leq \lambda$, if and only if $T_{l(\mu)}^\lambda \in \mathcal{T}^\wedge(\lambda, \mu)$. Notice that for $j \in I(n, r)$, the λ -tableau T_j^λ is ascending if and only if $l(\lambda) \leq j$. Therefore we have a bijective correspondence

$$\begin{aligned} Y(\lambda, \mu)^\wedge := \{(l(\lambda), j) \in Y(n, r) \mid j \in \mu, l(\lambda) \leq j\} &\rightarrow \mathcal{T}^\wedge(\lambda, \mu) \\ (l(\lambda), j) &\mapsto T_j^\lambda. \end{aligned}$$

Denote by $D_{\lambda, \mu}^\wedge$ the image of $Y(\lambda, \mu)^\wedge$ in $D_{\lambda, \mu}$ under the correspondence (9). Boltje and Maisch define for each $\mu \leq \lambda$

$$\mathrm{Hom}_{\mathcal{H}}^\wedge(M^\mu, M^\lambda) := \bigoplus_{d \in D_{\lambda, \mu}^\wedge} R\varphi_d^{\lambda, \mu} \subset \mathrm{Hom}_{\mathcal{H}}(M^\mu, M^\lambda).$$

Then under the isomorphism (10), $\mathrm{Hom}_{\mathcal{H}}^\wedge(M^\mu, M^\lambda)$ corresponds to

$$\bigoplus_{(l(\lambda), j) \in I^2(\lambda, \mu)^\wedge} R\xi_{l(\lambda), j}.$$

But, since

$$\{\xi_{l(\lambda), j} \mid (l(\lambda), j) \in Y(n, r), l(\lambda) \leq j, \lambda \in \Lambda\}$$

is an R -basis of $S_{R,q}^+(n, r)$ and for any $\nu, \tau \in \Lambda(n, r)$

$$\xi_\nu \xi_{ij} \xi_\tau = \begin{cases} \xi_{ij}, & \text{if } i \in \nu, j \in \tau \\ 0, & \text{otherwise,} \end{cases}$$

we get that $\text{Hom}_{\mathcal{H}}^\wedge(M^\mu, M^\lambda)$ corresponds to $\xi_\lambda S_q^+(n, r) \xi_\mu$. We saw previously that $S_{R,q}^+(n, r) = L \oplus \mathcal{J}$. But if $\lambda \triangleright \mu$, we have $\xi_\lambda L \xi_\mu = 0$. Hence $\text{Hom}_{\mathcal{H}}^\wedge(M^\mu, M^\lambda)$ corresponds to $\xi_\lambda \mathcal{J} \xi_\mu$ if $\lambda \triangleright \mu$.

Next we define the Boltje-Maisch complex $\widetilde{\mathcal{C}}_*^\lambda$. We will restrict ourselves to the case when λ is a partition of r . For every right \mathcal{H} -module N the R -module $\text{Hom}_R(N, R)$ has the structure of a left \mathcal{H} -module given by

$$(h\varepsilon)(n) := \varepsilon(nh),$$

where $h \in \mathcal{H}$, $\varepsilon \in \text{Hom}_R(N, R)$, and $n \in N$. So given an R -module N' , the R -module $\text{Hom}_R(N, R) \otimes_R N'$ can be viewed as an \mathcal{H} -module via

$$h(\varepsilon \otimes n') = (h\varepsilon) \otimes n',$$

where $h \in \mathcal{H}$, $\varepsilon \in \text{Hom}_R(N, R)$, and $n' \in N'$.

For each $\lambda \in \Lambda^+(n, r)$, Boltje and Maisch define a complex

$$\widetilde{\mathcal{C}}_*^\lambda: 0 \rightarrow C_{a(\lambda)}^\lambda \xrightarrow{d_{a(\lambda)}^\lambda} C_{a(\lambda)-1}^\lambda \xrightarrow{d_{a(\lambda)-1}^\lambda} \dots \xrightarrow{d_1^\lambda} C_0^\lambda \xrightarrow{d_0^\lambda} C_{-1}^\lambda \rightarrow 0$$

in the following way:

$$C_{-1}^\lambda = \text{Hom}_R(S^\lambda, R),$$

$$C_k^\lambda = \bigoplus_{\substack{\mu^{(1)} \triangleright \dots \triangleright \mu^{(k)} \triangleright \lambda \\ \mu^{(1)}, \dots, \mu^{(k)} \in \Lambda(n, r)}} \text{Hom}_R(M^{\mu^{(1)}}, R) \otimes_R \text{Hom}_{\mathcal{H}}^\wedge(M^{\mu^{(2)}}, M^{\mu^{(1)}}) \\ \otimes_R \dots \otimes_R \text{Hom}_{\mathcal{H}}^\wedge(M^\lambda, M^{\mu^{(k)}}).$$

The differential $d_k^\lambda: C_k^\lambda \rightarrow C_{k-1}^\lambda$ is given by the sum $\sum_{t=0}^{k-1} (-1)^t d_{kt}$, where for $k \geq 1$ and $1 \leq t \leq k-1$, we set

$$\begin{aligned} d_{k0}(\varepsilon \otimes \phi_1 \otimes \dots \otimes \phi_k) &= \varepsilon \phi_1 \otimes \phi_2 \otimes \dots \otimes \phi_k, \\ d_{kt}(\varepsilon \otimes \phi_1 \otimes \dots \otimes \phi_k) &= \varepsilon \otimes \phi_1 \otimes \dots \otimes \phi_t \phi_{t+1} \otimes \dots \otimes \phi_k, \end{aligned} \quad (12)$$

and $d_0: \text{Hom}_R(M^\lambda, R) \rightarrow \text{Hom}_R(S^\lambda, R)$ is defined to be the restriction on S^λ .

Let us consider the resolution B_λ of W_λ . Applying the Schur functor to B_λ we obtain the exact sequence $F(B_\lambda)$, where

$$F(B_\lambda)_{-1} = \xi_\omega W_\lambda, \quad F(B_\lambda)_0 = \xi_\omega S_{R,q}(n, r) \xi_\lambda,$$

and for $k \geq 1$ the \mathcal{H} -module $F(B_\lambda)_k$ is given by

$$\bigoplus_{\substack{\mu^{(1)} \triangleright \dots \triangleright \mu^{(k)} \triangleright \lambda \\ \mu^{(1)}, \dots, \mu^{(k)} \in \Lambda(n, r)}} \xi_\omega S_q(n, r) \xi_{\mu^{(1)}} \otimes_R \xi_{\mu^{(1)}} \mathcal{J} \xi_{\mu^{(2)}} \otimes_R \dots \otimes_R \xi_{\mu^{(k)}} \mathcal{J} \xi_\lambda.$$

Notice that, for $\mu \in \Lambda(n, r)$ the subspace $\xi_\omega S_q(n, r)\xi_\mu$ corresponds under (10) to $\text{Hom}_{\mathcal{H}}(M^\mu, M^\omega)$. But $M^\omega = x_\omega \mathcal{H} = \mathcal{H}$, since Σ_ω is the trivial group and $x_\omega = \sum_{\pi \in \Sigma_\omega} T_\pi = T_{\text{id}}$. Thus $\xi_\omega S_{R, q}(n, r)\xi_\mu$ corresponds under (10) to $\text{Hom}_{\mathcal{H}}(M^\mu, \mathcal{H})$. Here we have that $\text{Hom}_{\mathcal{H}}(M^\mu, \mathcal{H})$ is a \mathcal{H} -module by

$$(h\psi)(m) = h\psi(m),$$

where $h \in \mathcal{H}$, $m \in M^\mu$, and $\psi \in \text{Hom}_{\mathcal{H}}(M^\mu, \mathcal{H})$.

Thus we can write

$$F(B_\lambda)_0 = \text{Hom}_{\mathcal{H}}(M^\lambda, \mathcal{H}),$$

$$F(B_\lambda)_k = \bigoplus_{\substack{\mu^{(1)} \triangleright \dots \triangleright \mu^{(k)} \triangleright \lambda \\ \mu^{(1)}, \dots, \mu^{(k)} \in \Lambda(n, r)}} \text{Hom}_{\mathcal{H}}(M^{\mu^{(1)}}, \mathcal{H}) \otimes_R \text{Hom}_{\mathcal{H}}^\wedge(M^{\mu^{(2)}}, M^{\mu^{(1)}}) \\ \otimes_R \dots \otimes_R \text{Hom}_{\mathcal{H}}^\wedge(M^\lambda, M^{\mu^{(k)}}),$$

and the differentials d_k in $F(B_\lambda)$ are the sums $\sum_{t=0}^{k-1} (-1)^t d_{kt}$, where the maps d_{kt} are defined analogously to (12).

We will prove that $F(B_\lambda)$ is isomorphic to the complex $\widetilde{\mathcal{C}}_*^\lambda$ in non-negative degrees. Since $F(B_\lambda)$ is exact and $\widetilde{\mathcal{C}}_*^\lambda$ is exact in the degrees 0 and -1 by [BM12, Theorems 4.2 and 4.4], the isomorphism in degree -1 will follow.

To prove that $F(B_\lambda)_k \cong \mathcal{C}_k^\lambda$ for $k \geq 0$, we start by showing that there is an isomorphisms of \mathcal{H} -modules

$$\mathfrak{F}_\mu : \text{Hom}_{\mathcal{H}}(M^\mu, \mathcal{H}) \rightarrow \text{Hom}_R(M^\mu, R)$$

such that for all $\nu \in \Lambda(n, r)$, $\psi \in \text{Hom}_{\mathcal{H}}(M^\mu, \mathcal{H})$, $\varphi \in \text{Hom}_{\mathcal{H}}(M^\nu, M^\mu)$, we have $\mathfrak{F}_\nu(\psi\varphi) = \mathfrak{F}_\mu(\psi)\varphi$.

We will prove this in a more general setting. Let $*$: $\mathcal{H} \rightarrow \mathcal{H}$ be the anti-automorphism of \mathcal{H} given by $T_\pi \mapsto T_\pi^* = T_{\pi^{-1}}$. Let M be any right \mathcal{H} -module. By [DJ86, Theorem 2.6] there is an isomorphism of R -modules

$$\text{Hom}_R(M, R) \rightarrow \text{Hom}_{\mathcal{H}}(M, \mathcal{H}) \\ \varphi \mapsto \hat{\varphi},$$

where

$$\hat{\varphi}(m) := \sum_{\sigma \in \Sigma_r} q^{-\text{lng}(\sigma)} \varphi(m T_\sigma^*) T_\sigma.$$

The inverse of this isomorphism is the map

$$\text{Hom}_{\mathcal{H}}(M, \mathcal{H}) \rightarrow \text{Hom}_R(M, R) \\ \psi \mapsto \tilde{\psi},$$

where $\tilde{\psi}(m)$ is the coefficient of T_{id} in the expansion

$$\psi(m) = \sum_{\sigma \in \Sigma_r} a_\sigma T_\sigma, \quad a_\sigma \in R.$$

Consider the symmetric associative bilinear form $f: \mathcal{H} \otimes \mathcal{H} \rightarrow R$ ([DJ86, Lemma 2.2 and proof of Theorem 2.3]) given by

$$f(T_\sigma, T_\pi) = \begin{cases} q^{\text{lng}(\sigma)}, & \text{if } \sigma = \pi^{-1} \\ 0, & \text{otherwise.} \end{cases}$$

Note that we have

$$f\left(\sum_{\sigma \in \Sigma_r} a_\sigma T_\sigma, T_{\text{id}}\right) = a_{\text{id}}(T_{\text{id}}, T_{\text{id}}) = a_{\text{id}}.$$

Thus for $m \in M$ we get $\tilde{\psi}(m) = f(\psi(m), T_{\text{id}})$. We will prove that $\psi \rightarrow \tilde{\psi}$ is an \mathcal{H} -module homomorphism. Recall that $\text{Hom}_R(M, R)$ is a left \mathcal{H} -module by $(h\varphi)(m) = \varphi(mh)$, where $h \in \mathcal{H}$, $m \in M$, $\varphi \in \text{Hom}_R(M, R)$, and $\text{Hom}_{\mathcal{H}}(M, \mathcal{H})$ is a left \mathcal{H} -module by $(h\psi)(m) = h\psi(m)$, where $h \in \mathcal{H}$, $\psi \in \text{Hom}_{\mathcal{H}}(M, \mathcal{H})$, $m \in M$.

Proposition 9.1. *The map*

$$\begin{aligned} \mathfrak{F}_M: \text{Hom}_{\mathcal{H}}(M, \mathcal{H}) &\rightarrow \text{Hom}_R(M, R) \\ \psi &\mapsto \tilde{\psi} \end{aligned} \tag{13}$$

where $\tilde{\psi}(m) = f(\psi(m), T_{\text{id}})$ for $m \in M$, is an isomorphism of \mathcal{H} -modules.

Proof. Given $h \in \mathcal{H}_{r,q}$, $\psi \in \text{Hom}_{\mathcal{H}}(M, \mathcal{H})$, and $m \in M$, we have

$$\begin{aligned} \widetilde{h\psi}(m) &= f((h\psi)(m), T_{\text{id}}) = f(h\psi(m), T_{\text{id}}) = f(h, \psi(m)T_{\text{id}}) \\ &= f(h, \psi(m)) = f(\psi(m), h) = f(T_{\text{id}}, \psi(m)h) = f(T_{\text{id}}, \psi(mh)) \\ &= f(\psi(mh), T_{\text{id}}) = \tilde{\psi}(mh) = (h\tilde{\psi})(m). \end{aligned}$$

□

Proposition 9.2. *Let M and N be right \mathcal{H} -modules. Then the following diagram is commutative*

$$\begin{array}{ccc} \text{Hom}_{\mathcal{H}}(M, \mathcal{H}) \otimes \text{Hom}_{\mathcal{H}}(N, M) & \xrightarrow{\mathfrak{F}_M \otimes \text{id}} & \text{Hom}_R(M, R) \otimes \text{Hom}_{\mathcal{H}}(N, M) \\ \circ \downarrow & & \downarrow \circ \\ \text{Hom}_{\mathcal{H}}(N, \mathcal{H}) & \xrightarrow{\mathfrak{F}_N} & \text{Hom}_R(N, R). \end{array}$$

Proof. Let $\varphi: N \rightarrow M$, $\psi: M \rightarrow \mathcal{H}$ be homomorphisms of right \mathcal{H} -modules. Then for all $n \in N$

$$\widetilde{\psi\varphi}(n) = f(\psi\varphi(n), T_{\text{id}}) = f(\psi(\varphi(n)), T_{\text{id}}) = \tilde{\psi}(\varphi(n)) = \tilde{\psi}\varphi(n).$$

Thus $\widetilde{\psi\varphi} = \tilde{\psi}\varphi$. □

Returning to our setting we will abbreviate \mathfrak{F}_{M^μ} to \mathfrak{F}_μ . For each $k \geq 0$ define the map $\tau_k: F(B_\lambda)_k \rightarrow \mathcal{C}_{k,\lambda}$ to be the direct sum

$$\tau_k := \bigoplus_{\substack{\mu^{(1)} \triangleright \dots \triangleright \mu^{(k)} \triangleright \lambda \\ \mu^{(1)}, \dots, \mu^{(k)} \in \Lambda(n,r)}} \mathfrak{F}_{\mu^{(1)}} \otimes \text{id} \otimes \dots \otimes \text{id}.$$

Then τ_k is an isomorphism of \mathcal{H} -modules for $k \geq 0$. From Proposition 9.2, we get that for every $k \geq 1$

$$d_{k,0}\tau_k = \tau_{k-1}d_{k,0}.$$

Moreover it is obvious that for all $k \geq 1$ and $1 \leq t \leq k$

$$d_{k,t}\tau_k = \tau_{k-1}d_{k,t}.$$

Thus for all $k \geq 1$ we have $d_k\tau_k = \tau_{k-1}d_k$. This shows that $\tau = (\tau_k)_{k \geq 1}$ is a chain transformation between the truncated complexes $F(B_\lambda)_{\geq 0}$ and $\tilde{\mathcal{C}}_{\geq 0,\lambda}$. Since every τ_k is an isomorphism of \mathcal{H} -modules, we get that $F(B_\lambda)$ and $\tilde{\mathcal{C}}_{*,\lambda}$ are isomorphic in non-negative degrees as promised. The existence of an isomorphism in degree -1 follows from the commutative diagram with exact rows

$$\begin{array}{ccccccc} F(B_\lambda)_1 & \xrightarrow{d_1} & F(B_\lambda)_0 & \xrightarrow{d_0} & F(W_\lambda) & \longrightarrow & 0 \\ \tau_1 \downarrow & & \tau_0 \downarrow & & \downarrow \exists! & & \\ \mathcal{C}_1^\lambda & \xrightarrow{d_1} & \mathcal{C}_0^\lambda & \xrightarrow{d_0} & S^\lambda & \longrightarrow & 0. \end{array}$$

Thus we proved

Theorem 9.3. *Let λ be a partition of r . Then the complexes $\tilde{\mathcal{C}}_*^\lambda$ and $F(B_\lambda)$ are isomorphic. In particular, $\tilde{\mathcal{C}}_*^\lambda$ is an exact complex.*

References

- [BH11] R. Boltje and R. Hartmann, *Permutation resolutions for Specht modules*, J. Algebraic Combin. **34** (2011), no. 1, 141–162. MR 2805203 (2012g:20024)
- [BM12] R. Boltje and F. Maisch, *Permutation resolutions for Specht modules of Hecke algebras*, J. Algebra **365** (2012), 12–28. MR 2928450
- [DD91] R. Dipper and S. Donkin, *Quantum GL_n* , Proc. London Math. Soc. (3) **63** (1991), no. 1, 165–211. MR 1105721 (92g:16055)
- [DJ86] R. Dipper and G. James, *Representations of Hecke algebras of general linear groups*, Proc. London Math. Soc. (3) **52** (1986), no. 1, 20–52. MR 812444 (88b:20065)
- [Don80] S. Donkin, *Hopf complements and injective comodules for algebraic groups*, Proc. London Math. Soc. (3) **40** (1980), no. 2, 298–319. MR 566493 (81f:20053)

- [Don86] ———, *On Schur algebras and related algebras. I*, J. Algebra **104** (1986), no. 2, 310–328. MR 866778 (89b:20084a)
- [Don96] ———, *Standard homological properties for quantum GL_n* , J. Algebra **181** (1996), no. 1, 235–266. MR 1382035 (97b:20065)
- [Don98] ———, *The q -Schur algebra*, London Mathematical Society Lecture Note Series, vol. 253, Cambridge University Press, Cambridge, 1998. MR 1707336 (2001h:20072)
- [Gre76] J. A. Green, *Locally finite representations*, J. Algebra **41** (1976), no. 1, 137–171. MR 0412221 (54 #348)
- [Gre07] ———, *Polynomial representations of GL_n* , augmented ed., Lecture Notes in Mathematics, vol. 830, Springer, Berlin, 2007, With an appendix on Schensted correspondence and Littelmann paths by K. Erdmann, Green and M. Schocker. MR 2349209 (2009b:20084)
- [Jan03] Jens Carsten Jantzen, *Representations of algebraic groups*, second ed., Mathematical Surveys and Monographs, vol. 107, American Mathematical Society, Providence, RI, 2003. MR 2015057 (2004h:20061)
- [Mat70] Hideyuki Matsumura, *Commutative algebra*, W. A. Benjamin, Inc., New York, 1970. MR 0266911 (42 #1813)
- [ML63] Saunders Mac Lane, *Homology*, Die Grundlehren der mathematischen Wissenschaften, Bd. 114, Academic Press Inc., Publishers, New York, 1963. MR 0156879 (28 #122)
- [SY12] A. P. Santana and I. Yudin, *Characteristic-free resolutions of Weyl and Specht modules*, Adv. Math. **229** (2012), no. 4, 2578–2601. MR 2880232
- [vdK89] Wilberd van der Kallen, *Longest weight vectors and excellent filtrations*, Math. Z. **201** (1989), no. 1, 19–31. MR 990185 (91a:20043)
- [Wei94] Charles A. Weibel, *An introduction to homological algebra*, Cambridge Studies in Advanced Mathematics, vol. 38, Cambridge University Press, Cambridge, 1994. MR 1269324 (95f:18001)
- [Woo92] D. Woodcock, *On Schur algebras for the group of lower triangular matrices*, Tech. report, University of Bielefeld, 1992.
- [Woo94a] D. J. Woodcock, *Borel Schur algebras*, Comm. Algebra **22** (1994), no. 5, 1703–1721. MR 1264736 (95e:20060)
- [Woo94b] ———, *A vanishing theorem for Schur modules*, J. Algebra **165** (1994), no. 3, 483–506. MR 1275916 (95d:20076)
- [Woo97] D. Woodcock, *Schur algebras and global bases: new proofs of old vanishing theorems*, J. Algebra **191** (1997), no. 1, 331–370. MR 1444503 (98d:17027)