





# Spectral Problems and Orthogonal Polynomials on the Unit Circle



Kenier Castillo Rodríguez

Department of Mathematics

Carlos III University of Madrid

A thesis submitted for the degree of  
*Doctor of Philosophy in Mathematics*

September 2012

© 2012  
Carlos III University of Madrid  
All Rights Reserved

---

# Spectral Problems and Orthogonal Polynomials on the Unit Circle <sup>1</sup>

Advisor:

Francisco Marcellán Español  
Full Professor  
*Carlos III University of Madrid*

Approved by <sup>2</sup>:

Walter Van Assche  
Full Professor  
Committee Chair  
*Catholic University of Leuven*

José M. Rodríguez  
Full Professor  
Committee Member  
*Carlos III University of Madrid*

Alicia Cachafeiro  
Full Professor  
Committee Member  
*University of Vigo*

Guillermo López Lagomasino  
Full Professor  
Committee Secretary  
*Carlos III University of Madrid*

Dimitar K. Dimitrov  
Full Professor  
Committee Member  
*State University of São Paulo*

Calification: Cum Laude  
September 25, 2012

---

<sup>1</sup>This work was supported by FPU Research Fellowships, Spanish Ministry of Education, Re. AP2008-00471 and Spanish Ministry of Science and Innovation, grant MTM2009-12740-C03-01.

<sup>2</sup>External examiners' reports by A. Sri Ranga (Full Professor, State University of São Paulo) and A. Zhedanov (Senior Researcher, Donetsk Institute of Physics and Technology).

---

To Elena, whose love makes it all possible

---



## Acknowledgements

I would like to express my deepest gratitude to my advisor F. (Paco) Marcellán for his patience and guidance during my doctoral studies. Additionally, my sincere thanks to L. E. Garza and F. (Fe) R. Rafaeli who helped me with ideas and suggestions during the last years.

My appreciation to my friends and colleagues from State University of São Paulo, E. X. L. de Andrade, C. F. Bracciali, D. K. Dimitrov, R. L. Lamblém, M. V. Mello, and A. Sri Ranga, as well as from Catholic University of Leuven, M. Humet and M. Van Barel, for their invaluable help.

I am grateful to my friends from Carlos III University of Madrid, J. Borrego, J. Grujic, G. Miritello, and J. M. Pérez-Pardo, for the many good times shared. I also am indebted to F. M. Dopico, A. G. García, E. Moro, and J. M. Rodríguez by their teachings and example.

Finally, I wish to thank my entire extended family, without their love and support this work would not have been possible.

— K. Castillo  
Toledo, Spain. May 6, 2012

---

## Abstract

The main purpose of the work presented here is to study transformations of sequences of orthogonal polynomials associated with a hermitian linear functional  $\mathcal{L}$ , using spectral transformations of the corresponding  $C$ -function  $F$ . We show that a rational spectral transformation of  $F$  is given by a finite composition of four canonical spectral transformations. In addition to the canonical spectral transformations, we deal with two new examples of linear spectral transformations. First, we analyze a spectral transformation of  $\mathcal{L}$  such that the corresponding moment matrix is the result of the addition of a constant on the main diagonal or on two symmetric sub-diagonals of the initial moment matrix. Next, we introduce a spectral transformation of  $\mathcal{L}$  by the addition of the first derivative of a complex Dirac linear functional when its support is a point on the unit circle or two points symmetric with respect to the unit circle. In this case, outer relative asymptotics for the new sequences of orthogonal polynomials in terms of the original ones are obtained. Necessary and sufficient conditions for the quasi-definiteness of the new linear functionals are given. The relation between the corresponding sequence of orthogonal polynomials in terms of the original one is presented. We also consider polynomials which satisfy the same recurrence relation as the polynomials orthogonal with respect to the linear functional  $\mathcal{L}$ , with the restriction that the Verblunsky coefficients are in modulus greater than one. With positive or alternating positive-negative values for Verblunsky coefficients, zeros, quadrature rules, integral representation, and associated moment problem are analyzed. We also investigate the location, monotonicity, and asymptotics of the zeros of polynomials orthogonal with respect to a discrete Sobolev inner product for measures supported on the real line and on the unit circle.

**Keywords:** Orthogonal polynomials on the real line; orthogonal polynomials on the unit circle; Szegő polynomials on the real line; Hankel matrices; Toeplitz matrices; discrete Sobolev orthogonal polynomials on the real line; discrete Sobolev orthogonal polynomials on the unit circle; outer relative asymptotics; zeros;  $\mathcal{S}$ -functions;  $\mathcal{C}$ -functions; rational spectral transformations; canonical spectral transformations.

**2010 MSC:** 42C05, 33C45

# Contents

<b>Contents</b>	<b>ix</b>
<b>List of Figures</b>	<b>xiii</b>
<b>List of Tables</b>	<b>xv</b>
<b>Nomenclature</b>	<b>xix</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Motivation and main objectives . . . . .	1
1.2 Overview of the text . . . . .	7
<b>2 Orthogonal polynomials</b>	<b>11</b>
2.1 Orthogonal polynomials on the real line . . . . .	12
2.1.1 Classical orthogonal polynomials . . . . .	20
2.1.2 $\mathcal{S}$ -functions and rational spectral transformations . . . . .	23
2.2 Orthogonal polynomials on the unit circle . . . . .	29
2.2.1 $\mathcal{C}$ -functions and rational spectral transformations . . . . .	36
2.2.2 Connection with orthogonal polynomials on $[-1, 1]$ . . . . .	40
<b>3 On special classes of Szegő polynomials</b>	<b>43</b>
3.1 Special class of Szegő polynomials . . . . .	44
3.1.1 Szegő polynomials from continued fractions . . . . .	45
3.1.2 Szegő polynomials from series expansions . . . . .	49
3.1.3 Polynomials of second kind . . . . .	52
3.1.4 Para-orthogonal polynomials . . . . .	53
3.2 Polynomials with real zeros . . . . .	57

## CONTENTS

---

3.2.1	Associated moment problem . . . . .	65
<b>4</b>	<b>Spectral transformations of moment matrices</b>	<b>71</b>
4.1	Hankel matrices . . . . .	72
4.1.1	Perturbation on the anti-diagonals of a Hankel matrix . . . . .	73
4.1.2	Zeros . . . . .	76
4.2	Toeplitz matrices . . . . .	79
4.2.1	Diagonal perturbation of a Toeplitz matrix . . . . .	80
4.2.2	General perturbation of a Toeplitz matrix . . . . .	87
<b>5</b>	<b>Spectral transformations associated with mass points</b>	<b>97</b>
5.1	Adding the derivative of a Dirac's delta . . . . .	98
5.1.1	Mass point on the unit circle . . . . .	99
5.1.1.1	Outer relative asymptotics . . . . .	102
5.1.2	Mass points outside the unit circle . . . . .	105
5.1.2.1	Outer relative asymptotics . . . . .	107
5.1.3	$\mathcal{C}$ -functions and linear spectral transformations . . . . .	109
5.2	Non-standard inner products . . . . .	111
5.2.1	Outer relative asymptotics . . . . .	112
5.2.2	Zeros . . . . .	116
<b>6</b>	<b>Generators of rational spectral transformations</b>	<b>127</b>
6.1	Hermitian polynomial transformation . . . . .	128
6.2	Laurent polynomial transformation: Direct problem . . . . .	130
6.2.1	Regularity conditions and Verblunsky coefficients . . . . .	130
6.2.2	$\mathcal{C}$ -functions . . . . .	132
6.3	Laurent polynomial transformations: Inverse problem . . . . .	133
6.3.1	Regularity conditions and Verblunsky coefficients . . . . .	136
6.3.2	$\mathcal{C}$ -functions . . . . .	144
6.4	Rational spectral transformations . . . . .	147
6.4.1	Generator system for rational spectral transformations . . . . .	150
<b>7</b>	<b>Conclusions and open problems</b>	<b>153</b>
7.1	Conclusions . . . . .	153
7.2	Open problems . . . . .	155

## CONTENTS

---

<b>A</b>	<b>Discrete Sobolev orthogonal polynomials</b>	<b>161</b>
A.1	Monotonicity and asymptotics of zeros . . . . .	161
A.1.1	Jacobi polynomials . . . . .	167
A.1.2	Laguerre polynomials . . . . .	169
<b>B</b>	<b>Uvarov spectral transformation</b>	<b>171</b>
B.1	Mass points on the unit circle . . . . .	171
B.2	Mass points outside the unit circle . . . . .	174
	<b>Bibliography</b>	<b>177</b>

## **CONTENTS**

---



# List of Figures

1.1	A model for 1-dimensional lattice . . . . .	2
4.1	Verblunsky coefficients associated with sub-diagonal perturbation: Bernstein-Szegő and Chebyshev polynomials . . . . .	86
5.1	Zeros of Sobolev orthogonal polynomials with variation on $n$ : Lebesgue polynomials . . . . .	123
5.2	Zeros of Sobolev orthogonal polynomials with variation on $n$ : Bernstein-Szegő polynomials . . . . .	124
5.3	Zeros of Sobolev orthogonal polynomials with variation on $\lambda$ : Lebesgue and Bernstein-Szegő polynomials . . . . .	125
6.1	Verblunsky coefficients associated with the inverse problem: Chebyshev polynomials . . . . .	142
6.2	Verblunsky coefficients associated with the inverse problem: Geronimus polynomials . . . . .	144
7.1	Zeros of the Szegő-type polynomials ${}_2F_1(-100, b + 1; 2b + 2; 1 - z)$ . . . . .	156
A.1	Zeros of Sobolev orthogonal polynomials: Jacobi polynomials . . . . .	168
A.2	Zeros of Sobolev orthogonal polynomials: Laguerre polynomials . . . . .	170

## **LIST OF FIGURES**

---

# List of Tables

4.1	Zeros of polynomials associated with anti-diagonal perturbation, $n = 2$	79
4.2	Zeros of polynomials associated with anti-diagonal perturbation, $n = 3$	79
5.1	Critical values of $\alpha$ for Sobolev orthogonal polynomials: Lebesgue polynomials . . . . .	123
A.1	Zeros for Sobolev orthogonal polynomials with $c = 1$ : Jacobi polynomials	168
A.2	Zeros for Sobolev orthogonal polynomials with $c = 2$ : Jacobi polynomials	169
A.3	Zeros for Sobolev orthogonal polynomials with $c = 0$ : Laguerre polynomials . . . . .	170
A.4	Zeros for Sobolev orthogonal polynomials with $c = -2$ : Laguerre polynomials . . . . .	170

## **LIST OF TABLES**

---

# Nomenclature

## Roman Symbols

$a_n$  Jacobi parameter

$b_n$  Jacobi parameter

$\mathbb{C}$  complex numbers

$\mathbb{C}_+$   $\{z \in \mathbb{C}; \Im z > 0\}$

$c_n$  moments associated with  $\mathcal{L}$  or  $d\sigma$

$\mathbb{D}$  unit disc,  $\{z \in \mathbb{C}; |z| < 1\}$

$\det$  determinant

$\deg$  degree of a polynomial

$F(z)$   $C$ -function

**H** Hankel matrix

$H_n$  Hermite polynomial

$\Im$  imaginary part

**I** identity matrix

$I$  support of the measure  $\mu$ ,  $\text{supp}(\mu) = \{x \in \mathbb{R}; \mu(x - \epsilon, x + \epsilon) > 0 \text{ for every } \epsilon > 0\}$

**J** Jacobi matrix

$K_n$  kernel polynomial

## LIST OF TABLES

---

$\mathbf{k}_n$	$\langle \Phi_n, \Phi_n \rangle_{\mathcal{L}} = \ \Phi_n\ _{\sigma}^2$
$\mathcal{L}$	hermitian linear functional
$L_n^{(\alpha)}$	Laguerre polynomial
$\mathcal{M}$	linear functional
$\mathcal{N}$	Nevai class
$P_n$	monic orthogonal polynomial on the real line
$P_n^{(\alpha, \beta)}$	Jacobi polynomial
$p_n$	orthonormal polynomial on the real line
$\mathbb{P}$	linear space of polynomials with complex coefficients, $\text{span}\{z^k\}_{k \in \mathbb{Z}_+}$
$\mathbb{P}_n$	linear space of polynomials with complex coefficients and degree at most $n$
$\Re$	real part
$\mathbb{R}$	real line
$\mathbb{R}_+$	positive half of the real line, $\{x \in \mathbb{R}; x > 0\}$
$\text{sgn}(x)$	$ x ^{-1}x, x \neq 0$
$\mathcal{S}$	$\mathcal{S}$ -function
$\mathcal{S}$	Szegő class
$\mathbf{T}$	Toeplitz matrix
$\mathbb{T}$	unit circle $\{z \in \mathbb{C};  z  = 1\}$
$T_n$	Chebyshev polynomials of the first kind
$\mathbf{v}^H$	transpose conjugate of $\mathbf{v}$
$\mathbf{v}^T$	transpose of $\mathbf{v}$
$\mathbb{Z}$	integers $\{0, \pm 1, \pm 2, \dots\}$
$\mathbb{Z}_+$	non-negative integers $\{0, 1, 2, \dots\}$

$\mathbb{Z}_-$  negative integers  $\{0, -1, -2, \dots\}$

**Greek Symbols**

$\delta_{n,m}$  Kronecker delta

$\mu$  measure supported on the real line

$\sigma$  measure supported on the unit circle

$\kappa_n$  leading coefficient of  $\phi_n(z) = \kappa_n z^n + \dots$ ,  $\kappa_n = \|\Phi_n\|_\sigma^{-1}$

$\Lambda$  linear space of Laurent polynomials with complex coefficients,  $\text{span}\{z^k\}_{k \in \mathbb{Z}}$

$\mu_n$  moments associated with  $\mathcal{M}$  or  $d\mu$

$\Omega_n$  second kind orthogonal polynomial on the unit circle

$\Phi_n$  monic orthogonal polynomial on the unit circle

$\phi_n$  orthonormal polynomial on the unit circle

$\Psi_n$  perturbed monic orthogonal polynomial on the unit circle

$\psi_n$  perturbed orthonormal polynomial on the unit circle

$\Phi_n^*(z)$  reverse polynomial  $z^n \overline{\Phi_n(z)}$

$(\Phi_n(z))^* \overline{\Phi_n(z^{-1})}$

**Other Symbols**

$(x)_n^+$  Pochhammer's symbol for the rising factorial,  $x(x+1)\cdots(x+n-1)$ ,  $n \geq 1$

## **LIST OF TABLES**

---



# Chapter 1

## Introduction

I tend to write about what interests me, in the hope that others will also be interested.

— J. Milnor. *Interview with John Milnor* [163]

Orthogonal polynomials have very useful properties in the solution of mathematical and physical problems. Their relations with moment problems [99; 176], rational approximation [24; 149], operator theory [102; 109], analytic functions (de Branges' proof [56] of the Bieberbach conjecture), interpolation, quadrature [46; 73; 141; 188], electrostatics [93], statistical quantum mechanics [180], special functions [11], number theory [15] (irrationality [16] and transcendence [55]), graph theory [27], combinatorics, random matrices [57], stochastic process [172], data sorting and compression [63], computer tomography [121], and their role in the spectral theory of linear differential operators and Sturm-Liouville problems [148], as well as their applications in the theory of integrable systems [69; 84; 143; 144] constitute some illustrative samples of their impact.

### 1.1 Motivation and main objectives

Let consider the classical mechanical problem of a 1-dimensional chain of particles with neighbor interactions. Assume that the system is homogeneous (contains no impurities) and that the mass of each particle is  $m$ . We denote by  $y_n$  the displacement of the  $n$ -th particle, and by  $\varphi(y_{n+1} - y_n)$  the interaction potential between neighboring

## 1. INTRODUCTION

---

particles. We can consider this system as a chain of infinitely many particles joined together with non-linear springs; see Figure 1.1.




Figure 1.1: A model for 1-dimensional lattice

Therefore, if

$$F(r) = -\frac{d}{dr}\varphi(r) = -\varphi'(r)$$

is the force of the spring when it is stretched by the amount  $r$ , and  $r_n = y_{n+1} - y_n$  is the mutual displacement, then by Newton's law, the equation that governs the evolution is

$$my_n'' = \varphi'(y_{n+1} - y_n) - \varphi'(y_n - y_{n-1}).$$

If  $F(r)$  is proportional to  $r$ , that is, when  $F(r)$  obeys Hooke's law, the spring is linear and the potential can be written as  $\varphi(r) = \frac{\kappa}{2}r^2$ . Thus, the equation of motion is

$$my_n'' = \kappa(y_{n-1} - 2y_n + y_{n+1}),$$

and the solutions  $y_n^{(l)}$  are given by a linear superposition of the normal modes. In particular, when the particles  $n = 0$  and  $n = N + 1$  are fixed,

$$y_n^{(l)} = C_n \sin\left(\frac{\pi l}{N+1}\right) \cos(\omega_l t + \delta_l), \quad l = 1, 2, \dots, N,$$

where  $\omega_l = 2\sqrt{\kappa/m} \sin(\pi l / (2N+2))$ , the amplitude  $C_n$  of each mode is a constant determined by the initial conditions. In this case there is no transfer of energy between the modes. Therefore, the linear lattice is non-ergodic, and cannot be an object of statistical mechanics unless some modification is made. In the early 1950s, the general belief was that if a non-linearity is introduced in the model, then the energy flows between the different modes, eventually leading to a stable state of statistical equilibrium [68]. This phenomenon was explained by the connection to solitons <sup>1</sup>.

---

<sup>1</sup>In mathematics and physics, a soliton is a self-reinforcing solitary wave (a wave packet or pulse) that maintains its shape while it travels at constant speed. Solitons are caused by a cancellation of non-linear and dispersive effects in the medium.

---

There are non-linear lattices which admit periodic behavior at least when the energy is not too high. Lattices with exponential interaction have the desired properties. The Toda lattice [191] is given by setting

$$\varphi(r) = e^{-r} + r - 1.$$

Flaschka [69] (see also [143; 144]) proved complete integrability for the Toda lattice by recasting it as a Lax equation for Jacobi matrices. Later, Van Moerbeke [142], following a similar work [138] on Hill's equation [122], used the Jacobi matrices to define the Toda hierarchy for the periodic Toda lattices, and to find the corresponding Lax pairs.

Flaschka's change of variable is given by

$$a_n = \frac{1}{2}e^{-(y_{n+1}-y_n)/2}, \quad b_n = \frac{1}{2}y'_n.$$

Hence the new variables obey the evolution equations

$$a'_n = a_n(b_{n+1} - b_n), \tag{1.1}$$

$$b'_n = 2(a_n^2 - a_{n-1}^2), \quad a_{-1} = 0, \quad n \geq 0, \tag{1.2}$$

with initial data  $b_n = b_n(0) = \overline{b_n(0)}$ ,  $a_n = a_n(0) > 0$ , which we suppose uniformly bounded<sup>1</sup>. Set  $\mathbf{J}_t$  to be the semi-infinite Jacobi matrix associated with the system (1.1)-(1.2), that is

$$\mathbf{J}_t = \begin{bmatrix} b_0(t) & a_0(t) & 0 & 0 & \cdots \\ a_0(t) & b_1(t) & a_1(t) & 0 & \cdots \\ 0 & a_1(t) & b_2(t) & a_2(t) & \ddots \\ 0 & 0 & a_2(t) & b_3(t) & \ddots \\ \vdots & \vdots & \ddots & \ddots & \ddots \end{bmatrix}.$$

We use the notation  $\mathbf{J}_0 = \mathbf{J}_\mu$ , which is the matrix of the operator of multiplication by  $x$  in the basis of orthonormal polynomials on the real line. Favard's theorem says that, given any Jacobi matrix  $\widetilde{\mathbf{J}}$ , there exists a measure  $\mu$  on the real line for which  $\widetilde{\mathbf{J}} = \mathbf{J}_\mu$ . In general,  $\mu$  is not unique.

Flaschka's main observation is that the equations (1.1)-(1.2) can be reformulated in

---

<sup>1</sup>  $\sup_n (|a_n| + |b_n|) < \infty$ .

## 1. INTRODUCTION

---

terms of the Jacobi matrix  $\mathbf{J}_t$  as Lax pairs

$$\mathbf{J}'_t = [\mathbf{A}, \mathbf{J}_t] = \mathbf{A}\mathbf{J}_t - \mathbf{J}_t\mathbf{A},$$

with

$$\mathbf{A} = \begin{bmatrix} 0 & a_0(t) & 0 & 0 & \cdots \\ -a_0(t) & 0 & a_1(t) & 0 & \cdots \\ 0 & -a_1(t) & 0 & a_2(t) & \ddots \\ 0 & 0 & -a_2(t) & 0 & \ddots \\ \vdots & \vdots & \ddots & \ddots & \ddots \end{bmatrix} = (\mathbf{J}_t)_+ - (\mathbf{J}_t)_-,$$

where we use the standard notation  $(\mathbf{J}_t)_+$  for the upper-triangular, and  $(\mathbf{J}_t)_-$  for the lower-triangular projection of the matrix  $\mathbf{J}_t$ . At the same time, the corresponding orthogonality measure  $d\mu(\cdot, t)$  goes through a simple spectral transformation,

$$d\mu(x, t) = e^{-tx} d\mu(x, 0), \quad t > 0. \quad (1.3)$$

Notice that spectral transformations of orthogonal polynomials on the real line play a central role in the solution of the problem, see also [34]. Indeed, the solution of Toda lattice is a combination of the inverse spectral problem from  $\{a_n\}_{n \geq 0}$ ,  $\{b_n\}_{n \geq 0}$  associated with the measure  $d\mu = d\mu(\cdot, 0)$ , the spectral transformation (1.3), and the direct spectral problem from  $\{a_n(t)\}_{n \geq 0}$ ,  $\{b_n(t)\}_{n \geq 0}$  associated with the measure  $d\mu(\cdot, t)$ .

Given the infinity matrix  $\mathbf{J}_\mu$ , which is a bounded self-adjoint operator in  $\ell^2(\mathbb{Z}_+)$ , we can define [80; 177] the so-called  $\mathcal{S}$ -function by

$$S(x) = \langle \mathbf{e}_1, (\mathbf{J}_\mu - x)^{-1} \mathbf{e}_1 \rangle,$$

where  $\{\mathbf{e}_i\}_{i \geq 0} = \{(\delta_{i,j})_{j \geq 0}\}_{i \geq 0}$  is a vector basis in  $\ell^2(\mathbb{Z}_+)$ . In terms of the spectral measure  $\mu$  associated with  $\mathbf{J}_\mu$ ,

$$S(x) = \int_I \frac{d\mu(y)}{x-y}. \quad (1.4)$$

In many problems, (1.4) has more simple analytical and transformation properties than the measure  $\mu$ , and, hence,  $S$  is often much more convenient for analysis. Recently, Peherstorfer, Spiridonov, and Zhedanov [154] established a correspondence between the Toda lattice and differential equations for (1.4), using an alternative approach proposed in [153]. If the coefficients  $\{a_n(t)\}_{n \geq 0}$ ,  $\{b_n(t)\}_{n \geq 0}$  satisfy the system

---

of equations (1.1)-(1.2) with  $a_0(t)$  and  $b_0(t)$  taken as arbitrary initial functions of time, then the corresponding  $\mathcal{S}$ -function  $S(\cdot, t)$  satisfies the Riccati equation

$$\frac{\partial}{\partial t}S(x, t) = -1 + (x - b_0(t))S(x, t) - a_0^2(t)S^2(x, t).$$

Usually, the Toda lattice is studied using matrix spectral functions [150].

The problem of classifying all possible spectral transformations of orthogonal polynomials corresponding to a rational spectral transformation of the  $\mathcal{S}$ -function  $S$ , i.e.,

$$S_R(x) = \frac{a(x)S(x) + b(x)}{c(x)S(x) + d(x)}, \quad a(x)d(x) - b(x)c(x) \neq 0, \quad (1.5)$$

where  $a$ ,  $b$ ,  $c$ , and  $d$  are coprime polynomials, in other words, the description of a generator system of the set of rational spectral transformations, was raised by Marcellán, Dehesa, and Ronveaux [123] in 1990. Two years later, Peherstorfer [151] analyzed a particular class of rational spectral transformations. Indeed, he deduced the relation between the corresponding linear functionals. In 1997, Zhedanov [202] proved that the linear spectral transformation, i.e.,

$$S_L(x) = \frac{a(x)S(x) + b(x)}{d(x)}, \quad (1.6)$$

where  $a$ ,  $b$ , and  $d$  are polynomials that provide a true asymptotic behavior to (1.5) can be represented as a finite composition of Christoffel [188] and Geronimus [74; 75] spectral transformations, and also that any rational spectral transformation can be obtained as a finite composition of linear and associated elementary transformations [202]. Notice that (1.6) is not contained into (1.5), to achieve this we need to relax the conditions imposed on (1.5). We recall that the problem of classifying general rational spectral transformation is not yet solved. Here a natural question arises. What we can say about the generator system for rational spectral transformations of  $C$ -functions in the theory of orthogonal polynomials on the unit circle? This work is organized around this question.

Surprisingly, the theory of orthogonal polynomials with respect to non-trivial probability measures supported on the unit circle had not been so popular until the mid-1980's. The monographs by Szegő [87; 188], Freud [71], and Geronimus [78] were the main (and the few) major contributions to the subject, despite the fact that people working in linear prediction theory and digital signal processing used as a basic background orthogonal polynomials on the unit circle; see [59] and references therein. The recent

## 1. INTRODUCTION

---

monograph by Simon [178] constitutes an updated overview of the most remarkable directions of research in the theory, both from a theoretical approach (extensions of the Szegő theory from an analytic point of view), as well as from their applications in the spectral analysis of unitary operators, GGT [77; 86; 190] and CMV [29] matrix representations of the multiplication operator, quadrature formulas, and integrable systems (Ablowitz-Ladik systems [146], which include Schur flows as particular case), among others. Many concepts developed on orthogonal polynomials on the real line have analogues in this theory.

The Schur flow equation – which can be naturally called Toda lattice for the unit circle – is given by

$$\alpha'_n = (1 - |\alpha_n|^2)(\alpha_{n+1} - \alpha_{n-1}), \quad \alpha_{-1} = 0, \quad n \geq 0, \quad (1.7)$$

where  $\{\alpha_n\}_{n \geq 0}$  is a complex function sequence with  $|\alpha_n| < 1$ , initially occurred in [1; 2], under the name of discrete modified KdV equation, as a spatial discretization of the modified Korteweg-de Vries equation [108]

$$\frac{\partial}{\partial t} f(x, t) = 6f^2(x, t) \frac{\partial}{\partial x} f(x, t) - \frac{\partial^3}{\partial x^3} f(x, t).$$

In a very recent work [84], Golinskii proved that the solution of the system (1.7) reduces to the combination of the direct and the inverse spectral problems related by means of the Bessel transformation

$$d\sigma(z, t) = C(t)e^{t(z+z^{-1})} d\sigma(z, 0), \quad t > 0, \quad (1.8)$$

where  $\sigma$  is a non-trivial probability measure supported on the unit circle and  $C(t)$  is a normalization factor. Additionally, using CMV matrices the Lax pair for this system is found, and the dynamics of the corresponding spectral measures are described.

Given (1.4), the natural 'S-function' in the theory of orthogonal polynomials on the unit circle is the C-function  $F$  [177] given by

$$F(z) = \int_{-\pi}^{\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} d\sigma(\theta), \quad (1.9)$$

where  $\sigma$  is a non-trivial probability measure supported on  $[-\pi, \pi]$ . The Cauchy kernel

---

has the Poisson kernel as its real part, and this is positive, so

$$\Re F(z) > 0, \quad |z| < 1, \quad F(0) = 1.$$

Hence, (1.9) is the function introduced by Carathéodory in [31].

In this work – in the more general framework of hermitian linear functionals which are not necessarily positive definite – we consider, sequences of orthogonal polynomials deduced from spectral transformations of the corresponding  $C$ -function  $F$ . Our aim is to obtain and analyze the generator system of rational spectral transformations for non-trivial  $C$ -functions given by

$$\tilde{F}(z) = \frac{A(z)F(z) + B(z)}{C(z)F(z) + D(z)}, \quad A(z)D(z) - B(z)C(z) \neq 0, \quad (1.10)$$

where  $A$ ,  $B$ ,  $C$ , and  $D$  are coprime polynomials or Laurent polynomials. This result can be considered as a 'unit circle analogue' of the well known result by Zhedanov for orthogonal polynomials on the real line [202]. Furthermore, we introduce and study relevant examples of general linear spectral transformations associated with the addition of Lebesgue measure and derivatives of complex Dirac's deltas. However, in this, as well as in all research directions, more problems related to our original problem have arisen; some have been solved, and the rest are in Chapter 7 as a part of the open problems formulated therein.

## 1.2 Overview of the text

The original contributions of this work appear in twelve articles whose content is distributed as follows.

Chapter 3 develops the results of [41; 42]. Chapter 4 corresponds to [33; 36; 38]. The results of Chapter 5 are contained in [32; 37; 39; 40]. In Chapter 6 are included the results of [35; 43]. Finally, the Appendixes A and B contain results discussed in [37; 44].

Chapter 2 is meant for non-experts and therefore it contains some introductory and background material. We give a brief outline of orthogonal polynomials on the real line and on the unit circle, respectively. However, proofs of statements are not given. The emphasis is focussed on spectral transformation of the corresponding  $\mathcal{S}$ -functions and  $C$ -functions. This chapter could be omitted without destroying the unity

## 1. INTRODUCTION

---

or completeness of the work. The original content of this work appears in the next chapters. Let us describe briefly our main contributions.

In Chapter 3 we study the sequence of polynomials  $\{\Phi_n\}_{n \geq 0}$  which satisfies the following recurrence relation

$$\Phi_{n+1}(z) = z\Phi_n(z) + (-1)^{n+1}\alpha_{n+1}\Phi_n^*(z), \quad \alpha_{n+1} \in \mathbb{C}, \quad n \geq 0,$$

with the restriction  $|\alpha_{n+1}| > 1$ . The analysis of Perron-Carathéodory continued fractions shows that these polynomials satisfy the Szegő orthogonality with respect to a hermitian linear functional  $\mathcal{L}$  in  $\mathbb{P}$ , which satisfies a special quasi-definite condition. In two particular cases,  $\alpha_n > 0$  and  $(-1)^n\alpha_n > 0$ , respectively, zeros of the sequence of polynomials  $\{\Phi_n\}_{n \geq 0}$  (real Szegő polynomials [194]) and associated quadrature rules are also studied. As a consequence of this study, we solve the following moment problem. Given a sequence  $\{\mu_n\}_{n=0}^{\infty}$  of real numbers, we find necessary and sufficient conditions for the existence and uniqueness of a measure  $\mu$  supported on  $(1, \infty)$ , such that

$$\mu_n = \int_1^{\infty} T_n(x) d\mu(x), \quad n \geq 0. \quad (1.11)$$

Here  $\{T_n\}_{n \geq 0}$  are the Chebyshev polynomials of the first kind.

It is very well known that the Gram matrix of the bilinear form in  $\mathbb{P}$  associated with a linear functional  $\mathcal{L}$  in  $\Lambda$ , in terms of the canonical basis  $\{z^n\}_{n \geq 0}$ , is a Toeplitz matrix. In Chapter 4 we analyze two linear spectral transformations of  $\mathcal{L}$  such that the corresponding Toeplitz matrix is the result of the addition of a constant in the main diagonal, i.e.,

$$\langle f, g \rangle_{\mathcal{L}_0} = \langle f, g \rangle_{\mathcal{L}} + m \int_{\mathbb{T}} f(z) \overline{g(z)} \frac{dz}{2\pi iz}, \quad f, g \in \mathbb{P}, \quad m \in \mathbb{R},$$

or in two symmetric sub-diagonals, i.e.,

$$\langle f, g \rangle_{\mathcal{L}_j} = \langle f, g \rangle_{\mathcal{L}} + \mathbf{m} \int_{\mathbb{T}} z^j f(z) \overline{g(z)} \frac{dz}{2\pi iz} + \overline{\mathbf{m}} \int_{\mathbb{T}} z^{-j} f(z) \overline{g(z)} \frac{dz}{2\pi iz}, \quad \mathbf{m} \in \mathbb{C}, \quad j > 0,$$

of the initial Toeplitz matrix. We focus our attention on the analysis of the quasi-definite character of the perturbed linear functional, as well as in the explicit expressions of the new orthogonal polynomial sequence in terms of the first one. These transformations are known as local spectral transformations of the corresponding  $\mathcal{C}$ -function (1.9); see Chapter 6. Analogous transformations for orthogonal polynomials



---

on the real line, i.e., perturbations on the anti-diagonals of the corresponding Hankel matrix, are also considered. We define the modification of a quasi-definite functional  $\mathcal{M}$  by the addition of derivatives of a real Dirac's delta linear functional, whose action results in such a perturbation, i.e.,

$$\langle \mathcal{M}_j, p \rangle = \langle \mathcal{M}, p \rangle + mp^{(j)}(a), \quad p \in \mathbb{P}, \quad m, a \in \mathbb{R}, \quad j \geq 0.$$

We establish necessary and sufficient conditions in order to preserve the quasi-definite character. A relation between the corresponding sequences of orthogonal polynomials is obtained, as well as the asymptotic behavior of their zeros. We also determine the relation between such perturbations and the so-called canonical linear spectral transformations. In the first part of Chapter 5 we deal with a new example of linear spectral transformation associated with the influence of complex Dirac's deltas and their derivatives on the quasi-definiteness and the sequence of orthogonal polynomials associated with  $\mathcal{L}$ . This problem is related to the inverse polynomial modification [30], which is one of the generators of linear spectral transformations for the  $\mathcal{C}$ -function (1.9), as we see in Chapter 6. We analyze the regularity conditions of a modification of the quasi-definite linear functional  $\mathcal{L}$  by the addition of the first derivative of the complex Dirac's linear functional when its support is a point on the unit circle, i.e.,

$$\langle f, g \rangle_{\mathcal{L}_1} = \langle f, g \rangle_{\mathcal{L}} - im \left( \alpha f'(\alpha) \overline{g(\alpha)} - \bar{\alpha} f(\alpha) \overline{g'(\alpha)} \right), \quad m \in \mathbb{R}, \quad |\alpha| = 1,$$

or two symmetric points with respect to the unit circle, i.e.,

$$\begin{aligned} \langle f, g \rangle_{\mathcal{L}_2} = & \langle f, g \rangle_{\mathcal{L}} + im \left( \alpha^{-1} f(\alpha) \overline{g'(\bar{\alpha}^{-1})} - \alpha f'(\alpha) \overline{g(\bar{\alpha}^{-1})} \right) \\ & + i\bar{m} \left( \bar{\alpha} f(\bar{\alpha}^{-1}) \overline{g'(\alpha)} - \bar{\alpha}^{-1} p'(\bar{\alpha}^{-1}) \overline{q(\alpha)} \right), \quad m, \alpha \in \mathbb{C}, \quad |\alpha| \neq 0, 1. \end{aligned}$$

Outer relative asymptotics for the new sequence of monic orthogonal polynomials in terms of the original ones are obtained.

In the second part of Chapter 5 we assume  $\mathcal{L}$  is a positive definite linear functional associated with a positive measure  $\sigma$ . We study the relative asymptotics of the discrete Sobolev orthogonal polynomials. We focus our attention on the behavior of the zeros

## 1. INTRODUCTION

---

with respect to the particular case,

$$\langle f, g \rangle_{S_1} = \int_{\mathbb{T}} f(z) \overline{g(z)} d\sigma(z) + \lambda f^{(j)}(\alpha) \overline{g^{(j)}(\alpha)}, \quad \alpha \in \mathbb{C}, \quad \lambda \in \mathbb{R}_+, \quad j \geq 0.$$

In Chapter 6 we obtain and study the set of generators for rational spectral transformations, which are related with the direct polynomial modification, i.e.,

$$\langle \mathcal{L}_R, f \rangle = \left\langle \mathcal{L}, \left( z - \alpha + z^{-1} - \bar{\alpha} \right) f(z) \right\rangle, \quad f \in \Lambda, \quad \alpha \in \mathbb{C},$$

and the inverse of a polynomial modification, i.e.,

$$\left\langle \mathcal{L}_{R^{(-1)}}, \left( z - \alpha + z^{-1} - \bar{\alpha} \right) f(z) \right\rangle = \langle \mathcal{L}, f \rangle, \quad f \in \Lambda, \quad \alpha \in \mathbb{C},$$

as well as with the  $\pm k$  associated polynomials [152]. We deduce the relation between the corresponding  $C$ -functions and we study the regularity of the new linear functionals. We classify the spectral transformations of a  $C$ -function in terms of the moments associated with the linear functional  $\mathcal{L}$ . We also characterize the polynomial coefficients of a generic rational spectral transformation.

In Chapter 7 some concluding remarks which include indications of the direction of future work are presented. Finally, in Appendix A we consider the discrete Sobolev inner product associated with measures supported on the interval  $(a, b) \subseteq \mathbb{R}$  (not necessarily bounded), i.e.,

$$\langle p, q \rangle_{D_1} = \int_a^b p(x)q(x)d\mu(x) + \lambda p^{(j)}(\alpha)q^{(j)}(\alpha), \quad \alpha \notin (a, b), \quad \lambda > 0, \quad j \geq 0,$$

generalizing some known results concerning the asymptotic behavior of the zeros of the corresponding sequence of orthogonal polynomials. We also provide some numerical examples to illustrate the behavior of the zeros. Moreover, in Appendix B, the Uvarov perturbation of a quasi-definite linear functional by the addition of Dirac's linear functionals supported on  $r$  different points is studied.

## Chapter 2

# Orthogonal polynomials

What is true for OPRL<sup>1</sup> is even more true for orthogonal polynomials on the unit circle (OPUC).

— B. Simon. OPUC on one foot. *Boll. Amer. Math. Soc.*, 42:431-460, 2005

Orthogonal polynomials on the real line have attracted the interest of researchers for a long time. This subject is a classical one whose origins can be traced to Legendre's work [114] on planetary motion. The study of the algebraic and analytic properties of orthogonal polynomials in the complex plane was initiated by Szegő in [187], and later continued by Szegő himself and several authors as Geronimus, Keldysh, Korovkin, Lavrentiev, and Smirnov. An overview of the developments until 1964, with more than 50 references on this subject, is due to Suetin [184]. The complex analogue of the theory of orthogonal polynomials on the real line is naturally played by orthogonal polynomials on the unit circle. Following the works of Stieltjes, Hamburger, Toeplitz and others, Szegő investigated orthogonality on the unit circle in a series of papers around 1920 [185; 186], where he introduced orthogonal polynomials, known in the literature as Szegő polynomials.

In this chapter we present a short introduction to the theory of orthogonal polynomials on the real line (especially for comparison purposes) and orthogonal polynomials on the unit circle. We discuss recurrence relations, reproducing kernel, associated moment problems, distribution of their zeros, quadrature rules, among other results that we need in the sequel. We also consider transformations of orthogonal polynomials

---

<sup>1</sup>Orthogonal Polynomials on the Real Line.

## 2. ORTHOGONAL POLYNOMIALS

---

using spectral transformations of the corresponding  $\mathcal{S}$ -functions and  $\mathcal{C}$ -functions, respectively. Finally, we establish the connection between measures on a bounded interval and on the unit circle by the so-called Szegő transformation. Most of the material is classical and available in different monographs as [46], [71], [188], [78; 79], and the very recent monographs by Simon [178; 180]. Therefore formal theorems and proofs are not given.

### 2.1 Orthogonal polynomials on the real line

#### Definition

Let  $\mathcal{M}$  be a linear functional in the linear space  $\mathbb{P}$  of the polynomials with complex coefficients. We define the moment of order  $n$  associated with  $\mathcal{M}$  as the complex number

$$\mu_n = \langle \mathcal{M}, x^n \rangle, \quad n \geq 0. \quad (2.1)$$

The Gram matrix associated with the canonical basis  $\{x^n\}_{n \geq 0}$  of  $\mathbb{P}$  is given by

$$\mathbf{H} = \left[ \langle \mathcal{M}, x^{i+j} \rangle \right]_{i,j \geq 0} = \begin{bmatrix} \mu_0 & \mu_1 & \dots & \mu_n & \dots \\ \mu_1 & \mu_2 & \dots & \mu_{n+1} & \dots \\ \vdots & \vdots & \ddots & \vdots & \\ \mu_n & \mu_{n+1} & \dots & \mu_{2n} & \dots \\ \vdots & \vdots & & \vdots & \ddots \end{bmatrix}. \quad (2.2)$$

The matrices of this type, with constant values along anti-diagonals, are known as Hankel matrices [92].

The moment functional (2.1) is said to be quasi-definite if the moment matrix  $\mathbf{H}$  is strongly regular, or, equivalently, if the determinants of the principal leading submatrices  $\mathbf{H}_n$  of order  $(n+1) \times (n+1)$  are all different from 0 for every  $n \geq 0$ . In this case there exists a unique (up to an arbitrary non-zero factor) sequence  $\{P_n\}_{n \geq 0}$  of monic orthogonal polynomials with respect to  $\mathcal{M}$ . We define the orthogonal monic polynomial,  $P_n$ , of degree  $n$ , by

$$\langle \mathcal{M}, P_n P_m \rangle = \gamma_n^{-2} \delta_{n,m}, \quad \gamma_n \neq 0.$$

---

### Three-term recurrence relation

One of the most important characteristics of orthogonal polynomials on the real line is the fact that any three consecutive polynomials are connected by a simple relation which we can derive in a straightforward way. Indeed, let consider the polynomial  $P_{n+1}(x) - xP_n(x)$ , which is of degree at most  $n$ . Since  $\{P_k\}_{k=0}^n$  is a basis for the linear space  $\mathbb{P}_n$ , we can write

$$xP_n(x) = P_{n+1}(x) + \sum_{k=0}^n \lambda_{n,k} P_k(x), \quad \lambda_{n,k} = \frac{\langle \mathcal{M}, xP_n P_k \rangle}{\langle \mathcal{M}, P_k^2 \rangle}.$$

As  $P_n$  is orthogonal to every polynomial of degree at most  $n-1$ , we have  $\lambda_{n,0} = \lambda_{n,1} = \dots = \lambda_{n,n-2} = 0$  and

$$\lambda_{n,n-1} = \frac{\langle \mathcal{M}, P_n^2 \rangle}{\langle \mathcal{M}, P_{n-1}^2 \rangle}, \quad \lambda_{n,n} = \frac{\langle \mathcal{M}, xP_n^2 \rangle}{\langle \mathcal{M}, P_n^2 \rangle}.$$

We can thus find suitable complex numbers  $b_0, b_1, \dots$  and  $d_1, d_2, \dots$ , such that

$$xP_n(x) = P_{n+1}(x) + b_n P_n(x) + d_n P_{n-1}(x), \quad d_n \neq 0, \quad n \geq 0. \quad (2.3)$$

This three-term recurrence relation holds if we set  $P_{-1} = 0$  and  $P_0 = 1$  as initial conditions.

We next take up the important converse of the previous result. Let  $\{b_n\}_{n \geq 0}$  and  $\{d_n\}_{n \geq 1}$  be arbitrary sequences of complex numbers with  $d_n \neq 0$ , and let  $\{P_n\}_{n \geq 0}$  be defined by the recurrence relation (2.3). Then, there is a unique functional  $\mathcal{M}$  such that  $\langle \mathcal{M}, 1 \rangle = d_1$ , and  $\{P_n\}_{n \geq 0}$  is the sequence of monic orthogonal polynomials with respect to  $\mathcal{M}$ . We refer to this result as Favard's theorem [64].

### Jacobi matrices

We can write the three-term recurrence relation (2.3) in matrix form,

$$x\mathbf{P}(x) = \mathbf{J}\mathbf{P}(x), \quad \mathbf{P} = [P_0, P_1, \dots]^T,$$

where the semi-infinite tridiagonal matrix  $\mathbf{J}$  is defined by

## 2. ORTHOGONAL POLYNOMIALS

---

$$\mathbf{J} = \begin{bmatrix} b_0 & 1 & 0 & 0 & \cdots \\ d_1 & b_1 & 1 & 0 & \cdots \\ 0 & d_2 & b_2 & 1 & \ddots \\ 0 & 0 & d_3 & b_3 & \ddots \\ \vdots & \vdots & \ddots & \ddots & \ddots \end{bmatrix}.$$

$\mathbf{J}$  is said to be the monic Jacobi matrix [94] associated with the linear functional  $\mathcal{M}$ . A useful property of the matrix  $\mathbf{J}$  is that the eigenvalues of its  $n \times n$  leading principal submatrices  $\mathbf{J}_n$  are the zeros of the polynomial  $P_n$ . Indeed,  $P_n$  is the characteristic polynomial of  $\mathbf{J}_n$ ,

$$P_n(x) = \det(x\mathbf{I}_n - \mathbf{J}_n),$$

where  $\mathbf{I}_n$  is the  $n \times n$  identity matrix.

### Integral representation

We can say that  $\mathcal{M}$  is positive definite if and only if its moments are all real and  $\det \mathbf{H}_n > 0$ ,  $n \geq 0$ . In this case there exists a unique sequence of orthonormal polynomials  $\{p_n\}_{n \geq 0}$  with respect to  $\mathcal{M}$ , i.e., the following condition is satisfied,

$$\langle \mathcal{M}, p_n p_m \rangle = \delta_{n,m},$$

where

$$p_n(x) = \gamma_n x^n + \delta_n x^{n-1} + (\text{lower degree terms}), \quad \gamma_n > 0, \quad n \geq 0.$$

From the Riesz representation theorem [164; 170], we know that every positive definite linear functional  $\mathcal{M}$  has an integral representation (not necessarily unique)

$$\langle \mathcal{M}, x^n \rangle = \int_I x^n d\mu(x), \quad (2.4)$$

where  $\mu$  denotes a non-trivial positive Borel measure supported on some infinite subset  $I$  of the real line. For orthonormal polynomials, (2.3) becomes

$$x p_n(x) = a_{n+1} p_{n+1}(x) + b_n p_n(x) + a_n p_{n-1}(x), \quad a_n^2 = d_n, \quad n \geq 0, \quad (2.5)$$

---

with initial conditions  $p_{-1} = 0$ ,  $p_0 = \mu_0^{-1/2}$ , and the recurrence coefficients are given by

$$a_n = \int_I x p_{n-1}(x) p_n(x) d\mu(x) = \frac{\gamma_{n-1}}{\gamma_n} > 0,$$

$$b_n = \int_I x p_n^2(x) d\mu(x) = \frac{\delta_n}{\gamma_n} - \frac{\delta_{n+1}}{\gamma_{n+1}}.$$

Therefore,

$$p_n(x) = (a_n a_{n-1} \cdots a_1)^{-1} P_n(x) = \gamma_n P_n(x)$$

and the associated Jacobi matrix is

$$\mathbf{J}_\mu = \begin{bmatrix} b_0 & a_1 & 0 & 0 & \cdots \\ a_1 & b_1 & a_2 & 0 & \cdots \\ 0 & a_2 & b_2 & a_3 & \ddots \\ 0 & 0 & a_3 & b_3 & \ddots \\ \vdots & \vdots & \ddots & \ddots & \ddots \end{bmatrix}.$$

There are explicit formulas for orthogonal polynomials in terms of determinants. The orthonormal polynomial of degree  $n$  is given by Heine's formula [90]

$$p_n(x) = \frac{1}{\sqrt{\det \mathbf{H}_n \det \mathbf{H}_{n-1}}} \begin{vmatrix} \mu_0 & \mu_1 & \mu_2 & \cdots & \mu_n \\ \mu_1 & \mu_2 & \mu_3 & \cdots & \mu_{n+1} \\ \vdots & \vdots & \cdots & \vdots & \vdots \\ \mu_{n-1} & \mu_n & \mu_{n+1} & \cdots & \mu_{2n-1} \\ 1 & x & x^2 & \cdots & x^n \end{vmatrix}, \quad (2.6)$$

where the leading coefficient  $\gamma_n$  is the ratio of two Hankel determinants,

$$\gamma_n = \sqrt{\frac{\det \mathbf{H}_{n-1}}{\det \mathbf{H}_n}}.$$

In the positive definite case, Favard's theorem can be rephrased as follows. If  $d_{n+1} > 0$  and  $b_n \in \mathbb{R}$ ,  $n \geq 0$ , there is a non-trivial positive Borel measure  $\mu$  for which the Jacobi matrix is  $\mathbf{J}_\mu$ ; equivalently, the corresponding sequence of orthogonal polynomials obeys (2.5). In general this measure is not unique, but a sufficient condition for uniqueness is that the moment problem is determinate.

## 2. ORTHOGONAL POLYNOMIALS

---

### Moment problem

Moment problems occur in different mathematical contexts like probability theory, mathematical physics, statistical mechanics, potential theory, constructive analysis or dynamical systems. An excellent account of the history of moment problems is given in [107]. In its simplest terms, a moment problem is related to the existence of a measure  $\mu$  defined on an interval  $I \subseteq \mathbb{R}$  for which all the moments

$$\mu_n = \int_I x^n d\mu(x), \quad n \geq 0, \quad (2.7)$$

exist. If the solution to the moment problem is unique, it is called determinate. Otherwise, the moment problem is said to be indeterminate. The monographs [3] and [175] are the classical sources on moment problems; see also [176] from a different point of view using methods from the theory of finite difference operators.

There are many variations of a moment problem, depending on the interval  $I$ . In all of them, as suggested above, there are two questions to be answered, namely existence and uniqueness. Three particular cases of the general moment problem have come to be called classical moment problems, although strictly the term describes a much wider class. These are the following:

- i)* The Hamburger moment problem, where the measure is supported on  $(-\infty, \infty)$ .
- ii)* The Stieltjes moment problem, where the measure is supported on  $(0, \infty)$ .
- iii)* The Hausdorff moment problem, where the measure is supported on  $(0, 1)$ .

The Hausdorff moment problem is always determinate [89]. Stieltjes, in his memoir [182] introduced and solved the moment problem which was named after him by making extensive use of continued fractions. The necessary and sufficient conditions for determinacy of this moment problem are given by  $\det \mathbf{H}_n > 0$ ,  $\det \mathbf{H}_n^{(1)} > 0$ ,  $n \geq 0$ , where

$$\mathbf{H}_n^{(1)} = \begin{bmatrix} \mu_1 & \mu_2 & \dots & \mu_{n+1} \\ \mu_2 & \mu_3 & \dots & \mu_{n+2} \\ \vdots & \vdots & \ddots & \\ \mu_{n+1} & \mu_{n+2} & \dots & \mu_{2n+1} \end{bmatrix}.$$

In [88], Hamburger solved the moment problem on the whole real line, showing that it was not just a trivial extension of Stieltjes' work. The Hamburger moment problem is determined if and only if  $\det \mathbf{H}_n > 0$ ,  $n \geq 0$ .



---

More recent variations of these problems are the strong moment problems. In these cases, the sequence  $\{\mu_n\}_{n \geq 0}$  is replaced by the bilateral sequence  $\{\mu_n\}_{n \in \mathbb{Z}}$  of real numbers and the moment problem can be stated as follows. Given such a sequence  $\{\mu_n\}_{n \in \mathbb{Z}}$  of real numbers, find a measure  $\mu$  such that

$$\mu_n = \int_I x^n d\mu(x), \quad n \in \mathbb{Z}.$$

The strong Stieltjes and strong Hamburger moment problems can be formulated in the same way as the classical problems. The necessary and sufficient conditions are also given in terms of Hankel determinants involving the moments. Jones, Thron, and Waadeland [101] proposed and solved the strong Stieltjes moment problem, while Jones, Njåstad, and Thron [96] solved the strong Hamburger moment problem. In both cases, a central role was played by continued fractions.

### Continued fractions

As Brezinski [22] points out, continued fractions were used implicitly for many centuries before their real discovery. An excellent text on the arithmetical and metrical properties of regular continued fractions is the classical work of Khintchine [106], which is the starting point for the most recent book by Rocket and Szűsz [167]. In addition to these texts, the analytic theory of continued fractions is very well covered in [100; 119; 120; 196].

A continued fraction is a finite or infinite expansion of the form

$$q_0 + \frac{r_1}{r_2} = q_0 + \frac{r_1}{q_1 + \frac{r_2}{q_2 + \frac{r_3}{q_3 + \dots}}}, \quad (2.8)$$

where  $\{r_n\}_{n \geq 0}$  and  $\{q_n\}_{n \geq 0}$  are real or complex numbers, or functions of real or complex variables. The finite continued fraction,

$$\frac{R_n}{Q_n} = q_0 + \frac{r_1}{q_1} + \frac{r_2}{q_2} + \frac{r_3}{q_3} + \dots + \frac{r_n}{q_n},$$

obtained by truncation of (2.8), is called the  $n$ -th approximate of the continued fraction (2.8). The limit of  $R_n/Q_n$  when  $n$  tends to infinity is the value of the continued fraction.

## 2. ORTHOGONAL POLYNOMIALS

---

The numerators  $R_n$  and denominators  $Q_n$  satisfy, respectively, the Wallis recurrence relations [197]

$$R_{n+1} = q_{n+1}R_n + r_{n+1}R_{n-1}, \quad n \geq 1, \quad (2.9)$$

$$Q_{n+1} = q_{n+1}Q_n + r_{n+1}Q_{n-1}, \quad n \geq 1, \quad (2.10)$$

with  $R_0 = q_0$ ,  $Q_0 = 1$ ,  $R_1 = q_0q_1 + r_1$ , and  $Q_1 = q_1$ . These formulas lead directly to the connection between orthogonal polynomials and continued fractions. If we consider the following continued fraction

$$\cfrac{1}{x-b_0} - \cfrac{a_1^2}{x-b_1} - \cfrac{a_2^2}{x-b_2} - \cfrac{a_3^2}{x-b_3} - \dots,$$

then  $Q_n := P_n$  satisfies (2.3).

### Christoffel-Darboux identity

In the literature, the polynomials

$$K_n(x, y) = \sum_{k=0}^n p_k(x)p_k(y), \quad n \geq 0,$$

are usually called Kernel polynomials. The name comes from the fact that for any polynomial,  $q_n$ , of degree at most  $n$ , is given by

$$q_n(y) = \int_I q_n(x)K_n(x, y)d\mu(x).$$

The Kernel polynomial  $K_n$  can be represented in a simple way in terms of the polynomials  $p_n$  and  $p_{n+1}$  throughout the Christoffel-Darboux identity [45; 47; 53],

$$K_n(x, y) = a_{n+1} \frac{p_{n+1}(x)p_n(y) - p_n(x)p_{n+1}(y)}{x-y},$$

that can be deduced in a straightforward way from the three-term recurrence relation (2.5). When  $y$  tends to  $x$ , we obtain its confluent form

$$K_n(x, x) = a_{n+1} \left( p'_{n+1}(x)p_n(x) - p'_n(x)p_{n+1}(x) \right).$$

This last identity is used to prove two results that we show in this section, interlacing

---

of zeros and Gauss-Jacobi quadrature formula.

## Zeros

The fundamental theorem of algebra states that any polynomial of degree  $n$  has exactly  $n$  zeros (counting multiplicities). When dealing with orthogonal polynomials with respect to non-trivial probability measures supported on the real line, one can say much more about their localization. Two of the most relevant properties of zeros are the following:

- i) The zeros of  $p_n$  are all real, simple and are located in the interior of the convex hull<sup>1</sup> of  $I$ .
- ii) Suppose  $x_{n,1} < x_{n,2} < \dots < x_{n,n}$  are the zeros of  $p_n$ , then

$$x_{n,k} < x_{n-1,k} < x_{n,k+1}, \quad 1 \leq k \leq n-1.$$

The property ii) can also be proved using the Jacobi matrix  $\mathbf{J}_\mu$  from the inclusion principle for the eigenvalues of a hermitian matrix [92].

The following result is due to Wendroff [200]. Let  $P_{n+1}$  and  $P_n$  be two monic polynomials whose zeros are simple, real, and strictly interlacing. Then there is a positive Borel measure  $\mu$  for which they are the corresponding orthogonal polynomials of degrees  $n+1$  and  $n$ , respectively. All such measures have the same starting sequence  $P_{n+1}, P_n, P_{n-1}, \dots, P_0$ .

## Quadrature

A numerical quadrature consists of approximating the integral of a function  $f: I \subset \mathbb{R} \rightarrow \mathbb{R}$  by a finite sum which uses only  $n$  function evaluations. For a positive Borel measure  $\mu$  supported on  $I$ , an  $n$ -point quadrature rule is a set of points  $x_1, x_2, \dots, x_n$  and a set of associated numbers  $\lambda_1, \lambda_2, \dots, \lambda_n$ , such that

$$\int_I f(x) d\mu(x) \sim \sum_{k=1}^n \lambda_k f(x_k)$$

in some sense for a large class of functions as possible.

---

<sup>1</sup>By convex hull of a set  $E \subset \mathbb{C}$  we mean the smallest convex set containing  $E$ .  $G \subset \mathbb{C}$  is convex if for each pair of points  $x, y \in G$  the line connecting  $x$  and  $y$  is a subset of  $G$ .

## 2. ORTHOGONAL POLYNOMIALS

---

A Gauss-Jacobi quadrature [95; 182] is a quadrature rule constructed to yield an exact result for polynomials of degree at most  $2n - 1$ , by a suitable choice of the nodes and weights. If we choose the  $n$  nodes of a quadrature rule as the  $n$  zeros  $x_{n,1}, x_{n,2}, \dots, x_{n,n}$  of the orthogonal polynomial,  $P_n$ , with respect to  $\mu$  supported on  $I$  and if we denote the corresponding so-called Cotes or Christoffel numbers by  $\lambda_{n,1}, \lambda_{n,2}, \dots, \lambda_{n,n}$ , then for every polynomial  $Q_{2n-1}$  of degree at most  $2n - 1$ ,

$$\sum_{k=1}^n \lambda_{n,k} Q_{2n-1}(x_{n,k}) = \int_I Q_{2n-1}(x) d\mu(x).$$

The Christoffel numbers are positive and are given by

$$\lambda_{n,k} = \left( \sum_{k=0}^{n-1} p_k^2(x_{n,k}) \right)^{-1}.$$

### 2.1.1 Classical orthogonal polynomials

The most important polynomials on the real line are the classical orthogonal polynomials [188]. They are the Hermite polynomials, the Laguerre polynomials, the Jacobi polynomials (some special cases are the Gegenbauer polynomials, the Chebyshev polynomials, and the Legendre polynomials). These polynomials possess many properties that no other orthogonal polynomial system does. Among others, it is remarkable that the classical orthogonal polynomials satisfy a second order linear differential equation [17; 169]

$$\theta(x)y''(x) + \tau(x)y'(x) + \lambda_n y(x) = 0,$$

where  $\theta$  is a polynomial of degree at most 2 and  $\tau$  is a polynomial of degree 1, both independent of  $n$ . They also can be represented by a Rodrigues' distributional formula [51; 162]

$$P_n(x) = \frac{1}{c_n \omega(x)} \frac{d^n}{dx^n} (\omega(x)\theta^n(x)),$$

where  $\omega$  is the weight function and  $\theta$  is a polynomial independent of  $n$ . Moreover, for every classical orthogonal polynomial sequence, their derivatives constitute also an orthogonal polynomial sequence on the same interval of orthogonality [50; 110; 199].

---

## Jacobi polynomials

The Jacobi polynomials [46; 188], appear in the study of rotation groups <sup>1</sup>and in the solution to the equations of motion of the symmetric top [139]. They are orthogonal with respect to the absolutely continuous measure  $d\mu(\alpha, \beta; x) = (1-x)^\alpha(1+x)^\beta dx$ , supported on  $[-1, 1]$  where for integrability reasons we need to take  $\alpha, \beta > -1$ . These polynomials satisfy the orthogonality condition

$$\int_{-1}^1 P_n^{(\alpha, \beta)}(x) P_m^{(\alpha, \beta)}(x) d\mu(\alpha, \beta; x) = \frac{2^{\alpha+\beta+1}}{n!(2n+\alpha+\beta+1)} \frac{\Gamma(n+\alpha+1)\Gamma(n+\beta+1)}{\Gamma(n+\alpha+\beta+1)} \delta_{n,m},$$

where  $\Gamma$  is the Gamma function. From Rodrigues' formula we get

$$P_n^{(\alpha, \beta)}(x) = \frac{1}{(-2)^n n! (1-x)^\alpha (1+x)^\beta} \frac{d^n}{dx^n} \left( (1-x)^{\alpha+n} (1+x)^{\beta+n} \right),$$

or, equivalently, solving the differential equation by Frobenius' methods, the Jacobi polynomials are defined via the hypergeometric function as follows

$$\begin{aligned} P_n^{(\alpha, \beta)}(x) &= \frac{2^n (\alpha+1)_n^+}{(n+\alpha+\beta+1)_n^+} {}_2F_1 \left( -n, n+\alpha+\beta+1; \alpha+1; \frac{1-x}{2} \right) \\ &= \binom{n+\alpha}{n} \sum_{j=0}^n \frac{(-n)_k^+ (n+\alpha+\beta+1)_k^+}{(\alpha+1)_k^+ k!} \left( \frac{1-x}{2} \right)^j. \end{aligned}$$

The  $n$ -th Jacobi polynomial is the unique polynomial solution of the second order linear homogeneous differential equation

$$(x^2 - 1)y''(x) + ((2 + \alpha + \beta)x + \alpha - \beta)y'(x) - n(n + 1 + \alpha + \beta)y(x) = 0.$$

Particular cases are  $\alpha = \beta = -1/2$ , given the Chebyshev polynomials of first kind,

$$T_n(x) = 2^{2n} \frac{(n!)^2}{(2n)!} P_n^{(-1/2, -1/2)}(x).$$

The change of variable  $x = \cos \theta$  gives  $T_n(x) = \cos(n\theta)$ . The sequence  $\{T_n\}_{n \geq 0}$  is used as an approximation to a least squares fit, and it is a special case of the Gegenbauer polynomial with  $\alpha = 0$ .

---

<sup>1</sup>A rotation group is a group in which the elements are orthogonal matrices with determinant 1. In the case of three-dimensional space, the rotation group is known as the special orthogonal group.

## 2. ORTHOGONAL POLYNOMIALS

---

When  $\alpha = \beta = -1/2$ , we have the Chebyshev polynomial of second kind

$$U_n(x) = 2^{2n+1} \frac{((n+1)!)^2}{(2n+2)!} P_n^{(1/2, 1/2)}(x).$$

With  $x = \cos \theta$ , we get  $U_n(x) = \sin(n+1)\theta / \sin \theta$ . The sequence  $\{U_n\}_{n \geq 0}$  arises in the development of four-dimensional spherical harmonics in angular momentum theory.  $\{U_n\}_{n \geq 0}$  is also a special case of the Gegenbauer polynomial with  $\alpha = 1$ .

### Laguerre polynomials

The Laguerre polynomials [46; 188] arise in quantum mechanics, as the radial part of the solution of the Schrödinger's equation for the hydrogen atom. They are orthogonal on the positive half of the real line, satisfying

$$\int_0^\infty L_n^{(\alpha)}(x) L_m^{(\alpha)}(x) d\mu(\alpha; x) = \frac{\Gamma(n+\alpha+1)}{n!} \delta_{n,m},$$

where  $d\mu(\alpha; x) = x^\alpha e^{-x} dx$  and  $\alpha > -1$ . Rodrigues' formula for them is

$$L_n^{(\alpha)}(x) = \frac{e^x}{n! x^\alpha} \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha}).$$

The polynomial  $L_n^{(\alpha)}$  satisfies a second order linear differential equation that is a confluent hypergeometric equation

$$xy''(x) + (\alpha + 1 - x)y'(x) + ny(x) = 0,$$

and the Laguerre polynomials are a terminating confluent hypergeometric series

$$L_n^{(\alpha)}(x) = \frac{(-1)^n \Gamma(n+\alpha+1)}{\Gamma(\alpha+1)} {}_1F_1(-n, \alpha+1, x) = \sum_{j=0}^n \binom{n+\alpha}{n-j} \frac{(-x)^j}{j!}.$$

### Hermite polynomials

When  $d\mu(x) = e^{-x^2} dx$  on the whole real line, we have the Hermite polynomials [46; 188], satisfying the orthogonality relation

$$\int_{-\infty}^{\infty} H_n(x) H_m(x) d\mu(x) = \sqrt{\pi} 2^n n! \delta_{n,m}.$$

---

They arise in probability, such as the Edgeworth series, in numerical analysis as Gaussian quadrature, and in physics, where they give rise to the eigenstates of the quantum harmonic oscillator. From Rodrigues' formula,

$$H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2},$$

and we can deduce their explicit formula in terms of hypergeometric functions

$$H_{2n}(x) = (-1)^n (1/2)_n^+ {}_1F_1\left(-n, \frac{1}{2}; x^2\right),$$

$$H_{2n+1}(x) = (-1)^n (3/2)_n^+ x {}_1F_1\left(-n, \frac{3}{2}; x^2\right).$$

The choice  $\theta = 1$  and  $\tau(x) = -2x$  gives their characterization as the polynomial eigenfunctions of the second order linear differential operator

$$L[y(x)] = y''(x) - 2xy'(x).$$

## 2.1.2 $\mathcal{S}$ -functions and rational spectral transformations

### $\mathcal{S}$ -functions

The study of perturbations of the linear functional  $\mathcal{M}$  introduced in (2.1), and their effects on the corresponding  $\mathcal{S}$ -function

$$S(x) = \left\langle \mathcal{M}, \frac{1}{x-y} \right\rangle, \quad (2.11)$$

where the functional  $\mathcal{M}$  acts on the variable  $y$ , has a significant relevance in the theory of orthogonal polynomials on the real line.  $S$  admits, as a formal series expansion at infinity, the following equivalent representation

$$S(x) = \sum_{k=0}^{\infty} \frac{\mu_k}{x^{k+1}}, \quad (2.12)$$

i.e., it is a generating function of the sequence of moments for the linear functional  $\mathcal{M}$  (questions of convergence are not considered). If the moments associated with  $\mu$  are

## 2. ORTHOGONAL POLYNOMIALS

---

given by (2.7), the functions

$$S_n(x) = \int_I \frac{p_n(y)}{x-y} d\mu(y), \quad n \geq 0,$$

constitute a second (independent) solution of the difference equation

$$xy_n = a_{n+1}y_{n+1} + b_ny_n + a_ny_{n-1}, \quad n \geq 0.$$

They are called second kind functions associated with  $\mu$ . In this case, the  $\mathcal{S}$ -function [182] is given by

$$S_0(x) = S(x) = \int_I \frac{d\mu(y)}{x-y}.$$

One of the important properties of the  $\mathcal{S}$ -functions is its representation in terms of continued fractions, [182]

$$S(x) = \cfrac{1}{x-b_0} - \cfrac{a_1^2}{x-b_1} - \cfrac{a_2^2}{x-b_2} - \dots \quad (2.13)$$

In fact, (2.13) was a starting point for the general theory of orthogonal polynomials in pioneering works by Chebyshev [45] and Stieltjes [182].

### Spectral transformations

A rational spectral transformation [202] of the  $\mathcal{S}$ -function  $S$  is a new  $\mathcal{S}$ -function defined by

$$S_R(x) = \frac{a(x)S(x) + b(x)}{c(x)S(x) + d(x)}, \quad a(x)d(x) - b(x)c(x) \neq 0, \quad (2.14)$$

where  $a$ ,  $b$ ,  $c$ , and  $d$  are coprime polynomials. The spectral transformation

$$S_L(x) = \frac{a(x)S(x) + b(x)}{d(x)}, \quad (2.15)$$

where  $a$ ,  $b$ , and  $d$  are polynomials that provide a true asymptotic behavior to (2.14) is said to be linear. These polynomials should be chosen in such a way that the new  $\mathcal{S}$ -function,  $S_R$ , has the same asymptotic behavior as initial (2.12),

$$S_R(x) = \sum_{k=0}^{\infty} \frac{\tilde{\mu}_k}{x^{k+1}},$$



---

where  $\{\bar{\mu}_n\}_{n \geq 0}$  is the sequence of transformed moments. Hence, in general, the coefficients of the polynomials  $a$ ,  $b$ ,  $c$ , and  $d$  depend on the original moments  $\{\mu_n\}_{n \geq 0}$ . In particular, this means that the spectral transformations do not form a group. Indeed, for a given spectral transformation there exist many different reciprocal spectral transformations. Nevertheless, it is clear that one can always construct a composition of two spectral transformations, and moreover, for a given spectral transformation there is at least one reciprocal.

In terms of the moments, we can classify the spectral transformations of  $\mathcal{S}$ -functions as follows.

- i)* Local spectral transformations: spectral transformations under the modification of a finite number of moments.
- ii)* Global spectral transformations: spectral transformations under the modification of an infinite number of moments.

Notice that *i)* is a special case of general linear spectral transformations related with perturbations on the anti-diagonals of the Hankel matrix (2.2). A large subclass of *ii)* can be represented by the rational spectral transformation.

### Linear spectral transformations

Without loss of generality, we can assume that the measure  $\mu$  is normalized, i.e.,  $\mu_0 = 1$ . The Christoffel transformation [188] corresponds to a modification of the measure  $\mu$  defined by

$$d\mu_c(x) = \frac{x-\beta}{\mu_1-\beta} d\mu(x), \quad \beta \notin I. \quad (2.16)$$

The sequence of orthogonal polynomials  $\{P_n(\cdot; \mu_c)\}_{n \geq 0}$  associated with this transformation is given by

$$(x-\beta)P_n(x; \mu_c) = P_{n+1}(x) - \frac{P_{n+1}(\beta)}{P_n(\beta)} P_n(x), \quad n \geq 0.$$

Indeed, Christoffel transformation leads to the Kernel polynomial. Denoting the transformation (2.16) by  $\mathcal{R}_C(\beta)$ , the corresponding  $\mathcal{S}$ -function becomes

$$S_C(x) = \mathcal{R}_C(\beta)[S(x)] = \frac{(x-\beta)S(x) - 1}{\mu_1 - \beta}. \quad (2.17)$$

Conversely, if we start with a linear spectral transformation where  $a$  is a polynomial

## 2. ORTHOGONAL POLYNOMIALS

---

of first degree,  $b$  is constant,  $d \equiv 1$ , the only choice for such a spectral transformation is (2.17). In general, a linear spectral transformation with  $d \equiv 1$  is equivalent to a finite composition of Christoffel transformations [202].

The reciprocal of a Christoffel transformation is the so-called Geronimus transformation [74; 75], consisting of a perturbation of  $\mu$  such that

$$d\mu_g(x) = \frac{(\beta - x)^{-1} d\mu(x) + m\delta(x - \beta)}{m + S(\beta)}, \quad \beta \notin I, \quad m \in \mathbb{R}_+. \quad (2.18)$$

The sequence of orthogonal polynomials  $\{P_n(\cdot; \mu_g)\}_{n \geq 0}$  with respect to (2.18) can be written as

$$P_n(x; \mu_g) = xP_n(x) - \frac{Q_n(\beta; m)}{Q_{n-1}(\beta, m)} P_{n-1}(x), \quad n \geq 0$$

where  $P_0(\cdot; \mu_g) = 1$ , and  $Q_n(\beta, m)$  is a solution of the recurrence relation (2.3) with auxiliary parameter  $\beta$ ,

$$Q_n(\beta, m) = S_n(\beta) + mP_n(\beta), \quad n \geq 0.$$

This transformation is denoted by  $\mathcal{R}_G(\beta, m)$ . The corresponding  $\mathcal{S}$ -function is

$$S_G(x) = \mathcal{R}_G(\beta, m)[S(x)] = \frac{S(\beta) + m - S(x)}{(x - \beta)(m + S(\beta))}. \quad (2.19)$$

We can see that the transformation (2.19) is reciprocal to (2.17). However, in contrast to the Christoffel transformation, (2.19) contains two free parameters, where the second free parameter defines the value of additional discrete mass as is seen in (2.18). In general, one can prove that a linear spectral transformation with  $a \equiv 1$  is equivalent to a finite composition of Geronimus transformations [202].

It is easy to see that for different values of  $\beta$  we have

$$\mathcal{R}_C(\beta_1) \circ \mathcal{R}_G(\beta_2, m) = \mathcal{R}_G(\beta_2, m) \circ \mathcal{R}_C(\beta_1).$$

However, for the same parameter we have the following relations

$$\begin{aligned} \mathcal{R}_C(\beta) \circ \mathcal{R}_G(\beta, m) &= \mathcal{I} \quad (\text{Identity transformation}), \\ \mathcal{R}_G(\beta, m) \circ \mathcal{R}_C(\beta) &= \mathcal{R}_U(\beta, m) \quad (\text{Uvarov transformation}). \end{aligned}$$

The Uvarov transformation [74; 75] consists of the addition of a real positive mass

---

to the measure  $\mu$ ,

$$d\mu_u(x) = \frac{d\mu(x) + m\delta(x-\beta)}{1+m}, \quad \beta \notin I, \quad m \in \mathbb{R}_+.$$

The relation between the corresponding  $\mathcal{S}$ -functions is

$$S_U(x) = \mathcal{R}_U(\beta, m)[S(x)] = \frac{S(x) + m(x-\beta)^{-1}}{1+m}.$$

### Rational spectral transformations

In [202] it was proved that by means of  $\pm k$  associated transformations we can reduce (2.14) to the linear form. Combining Christoffel transformation (2.17), Geronimus transformations (2.19), and the  $\pm k$  associated transformations we get a wide class of rational spectral transformations (2.14).

From the sequence of monic orthogonal polynomials  $\{P_n\}_{n \geq 0}$  we can define the sequence of associated monic polynomials [76],  $\{P_n^{(k)}\}_{n \geq 0}$ ,  $k \geq 1$ , by means of the shifted recurrence relation

$$P_{n+1}^{(k)}(x) = (x - b_{n+k})P_n^{(k)}(x) - d_{n+k}P_{n-1}^{(k)}(x), \quad n \geq 0, \quad (2.20)$$

with  $P_{-1}^{(k)} = 0$  and  $P_0^{(k)} = 1$ . The recurrence relation (2.20) can also be written in the matrix form

$$x\mathbf{P}^{(k)}(x) = \mathbf{J}^{(k)}\mathbf{P}^{(k)}(x), \quad \mathbf{P}^{(k)} = [P_0^{(k)}, P_1^{(k)}, \dots]^T,$$

where  $\mathbf{J}^{(k)}$  is the tridiagonal matrix

$$\mathbf{J}^{(k)} = \begin{bmatrix} b_k & 1 & 0 & 0 & \cdots \\ d_{k+1} & b_{k+1} & 1 & 0 & \cdots \\ 0 & d_{k+2} & b_{k+2} & 1 & \ddots \\ 0 & 0 & d_{k+3} & b_{k+3} & \ddots \\ \vdots & \vdots & \ddots & \ddots & \ddots \end{bmatrix},$$

i.e., we have removed in the monic Jacobi matrix  $\mathbf{J}$  the first  $k$  rows and columns.

The  $\mathcal{S}$ -function corresponding to the associated polynomials of order  $k$ ,  $S^{(k)}$ , can



where  $\{\widetilde{d}_i\}_{i \geq 1} = \{d_{-i}\}_{i=k-1}^0 \cup \{d_i\}_{i \geq 1}$ . If  $k = 1$ , then the anti-associated polynomials of the first kind appear and its corresponding Stieltjes function  $S^{(-1)}$ , given by (2.21), is

$$S^{(-1)}(x) = \frac{1}{x - \widetilde{b}_0 - \widetilde{d}_1 S(x)}, \quad (2.22)$$

where  $S$  is the Stieltjes function associated with  $\mu$ , and  $\widetilde{b}_0$  and  $\widetilde{d}_1$  are free parameters. We denote  $\mathcal{R}_{(\widetilde{b}_0, \widetilde{d}_1)}^{(-1)}[S(x)] = S^{(-1)}(x)$ . Observe that  $\mathcal{R}^{(-1)}$  is not a unique inverse of  $\mathcal{R}^{(1)}$ , because of its dependence on the free parameters  $\widetilde{b}_0$  and  $\widetilde{d}_1$ .

## 2.2 Orthogonal polynomials on the unit circle

### Definition

Let  $\mathcal{L}$  be a linear functional in the linear space of Laurent polynomials with complex coefficients,  $\Lambda$ , satisfying

$$c_n = \langle \mathcal{L}, z^n \rangle = \overline{\langle \mathcal{L}, z^{-n} \rangle} = \bar{c}_{-n}, \quad n \in \mathbb{Z}. \quad (2.23)$$

$\mathcal{L}$  is said to be a hermitian linear functional. A bilinear functional associated with  $\mathcal{L}$  can be introduced in  $\mathbb{P}$  as follows

$$\langle f, g \rangle_{\mathcal{L}} = \langle \mathcal{L}, f(z)\bar{g}(z^{-1}) \rangle, \quad f, g \in \mathbb{P}.$$

The complex numbers  $\{c_n\}_{n \in \mathbb{Z}}$  are said to be the moments associated with  $\mathcal{L}$  and the infinite matrix

$$\mathbf{T} = \left[ \langle z^i, z^j \rangle_{\mathcal{L}} \right]_{i, j \geq 0} = \begin{bmatrix} c_0 & c_1 & \cdots & c_n & \cdots \\ c_{-1} & c_0 & \cdots & c_{n-1} & \cdots \\ \vdots & \vdots & \ddots & \vdots & \\ c_{-n} & c_{-n+1} & \cdots & c_0 & \cdots \\ \vdots & \vdots & & \vdots & \ddots \end{bmatrix}, \quad (2.24)$$

is the Gram matrix of the above bilinear functional in terms of the canonical basis  $\{z^n\}_{n \geq 0}$  of  $\mathbb{P}$ . It is known in the literature as a Toeplitz matrix, a matrix in which each descending diagonal from left to right is constant [92].

If  $\mathbf{T}_n$ , the  $(n+1) \times (n+1)$  principal leading submatrix of  $\mathbf{T}$ , is non-singular for every

## 2. ORTHOGONAL POLYNOMIALS

---

$n \geq 0$ ,  $\mathcal{L}$  is said to be quasi-definite, and there exists a sequence of monic polynomials  $\{\Phi_n\}_{n \geq 0}$ , orthogonal with respect to  $\mathcal{L}$ ,

$$\langle \Phi_n, \Phi_m \rangle_{\mathcal{L}} = \mathbf{k}_n \delta_{n,m}, \quad \mathbf{k}_n \neq 0, \quad n \geq 0.$$

### Szegő recurrence relations

We have seen that orthogonal polynomials on the real line satisfy a three-term recurrence relation. Such a recurrence relation does not hold for orthogonal polynomials on the unit circle, but there are also recurrence formulas. These polynomials satisfy the following forward and backward recurrence relations

$$\Phi_{n+1}(z) = z\Phi_n(z) + \Phi_{n+1}(0)\Phi_n^*(z), \quad n \geq 0, \quad (2.25)$$

$$\Phi_{n+1}(z) = \left(1 - |\Phi_{n+1}(0)|^2\right)z\Phi_n(z) + \Phi_{n+1}(0)\Phi_{n+1}^*(z), \quad n \geq 0, \quad (2.26)$$

where  $\Phi_n^*(z) = z^n \overline{\Phi_n(z^{-1})} = z^n (\Phi_n)_*(z)$  is the so-called reversed polynomial, and the complex numbers  $\{\Phi_n(0)\}_{n \geq 1}$ , with

$$|\Phi_n(0)| \neq 1, \quad n \geq 1,$$

are known as Verblunsky, Schur or reflection coefficients. The monic orthogonal polynomials are therefore completely determined by the sequence  $\{\Phi_n(0)\}_{n \geq 1}$ . To obtain the recurrence formula, we take into account the fact that the reversed polynomial  $\Phi_n^*(z)$  is the unique polynomial of degree at most  $n$  orthogonal to  $z^k$ ,  $1 \leq k \leq n$ . (2.25) and (2.26) are called either the Szegő recurrence or Szegő difference relations. Moreover, we have

$$\langle \Phi_n, \Phi_n \rangle_{\mathcal{L}} = \mathbf{k}_n = \frac{\det \mathbf{T}_n}{\det \mathbf{T}_{n-1}}, \quad n \geq 1, \quad \mathbf{k}_0 = c_0. \quad (2.27)$$

We can derive a recurrence formula which does not involve the reversed polynomials,

$$\Phi_n(0)\Phi_{n+1}(z) = (z\Phi_n(0) + \Phi_{n+1}(0))\Phi_n(z) - z\rho_n^2\Phi_{n+1}(0)\Phi_{n-1}(z), \quad n \geq 0, \quad (2.28)$$

if we assume  $\Phi_{-1} = 0$ . The polynomials  $\Phi_{n+1}$  can be found from  $\Phi_{n-1}$  and  $\Phi_n$ , if  $\Phi_n(0) \neq 0$ . This is an analogue of the three-term recurrence relation (2.3) for orthogonal polynomials on the real line, except for the factor  $z$  in the last term. In [52], the authors find bounds for complex zeros of polynomials generated by this kind of recurrence

relations.

### Integral representation

If  $c_0 = 1$  and  $\det \mathbf{T}_n > 0$ , for every  $n \geq 0$ ,  $\mathcal{L}$  is said to be positive definite and it has the following integral representation

$$\langle \mathcal{L}, f \rangle = \int_{\mathbb{T}} f(z) d\sigma(z), \quad f \in \mathbb{P}, \quad (2.29)$$

where  $\sigma$  is a non-trivial probability measure supported on the unit circle  $\mathbb{T}$ . In such a case, there exists a unique sequence of polynomials  $\{\phi_n\}_{n \geq 0}$  with positive leading coefficient, such that

$$\int_{\mathbb{T}} \phi_n(z) \overline{\phi_m(z)} d\sigma(z) = \delta_{m,n}.$$

$\{\phi_n\}_{n \geq 0}$  is said to be the sequence of orthonormal polynomials with respect to  $d\sigma$ . Denoting by  $\kappa_n$  the leading coefficient of  $\phi_n$ ,  $\Phi_n = \kappa_n^{-1} \phi_n$  is the corresponding monic orthogonal polynomial of degree  $n$ . Moreover,  $\langle \Phi_n, \Phi_n \rangle_{\mathcal{L}} = \|\Phi_n\|_{\sigma}^2 = \mathbf{k}_n > 0$ .

From the Pythagoras theorem, in (2.25) we get

$$\frac{\|\Phi_n\|_{\sigma}^2}{\|\Phi_{n-1}\|_{\sigma}^2} = 1 - |\Phi_n(0)|^2 > 0, \quad n \geq 1. \quad (2.30)$$

This shows that in the positive definite case the Verblunsky coefficients always satisfy

$$|\Phi_n(0)| < 1, \quad n \geq 1. \quad (2.31)$$

In this situation, we have an analogous of the Favard theorem [173; 174; 193], formulated as follows. Any sequence of complex numbers obeying (2.31) arises as the Verblunsky coefficients of a unique non-trivial probability measure supported on the unit circle.

We use the notation  $\rho_n = \sqrt{1 - |\Phi_n(0)|^2} = \|\Phi_n\|_{\sigma} / \|\Phi_{n-1}\|_{\sigma} = \kappa_{n-1} / \kappa_n$ . Hence, for the orthonormal polynomials  $\phi_n$ , the recurrence relations (2.25)-(2.28) become

$$\begin{aligned} \rho_{n+1} \phi_{n+1}(z) &= z \phi_n(z) - \Phi_{n+1}(0) \phi_n^*(z), \quad n \geq 0, \\ \phi_{n+1}(z) &= \rho_{n+1} z \phi_n(z) + \Phi_{n+1}(0) \phi_{n+1}^*(z), \quad n \geq 0, \\ \rho_{n+1} \Phi_n(0) \phi_{n+1}(z) &= (z \Phi_n(0) + \Phi_{n+1}(0)) \phi_n(z) - z \rho_n \Phi_{n+1}(0) \phi_{n-1}(z), \quad n \geq 0. \end{aligned} \quad (2.32)$$

## 2. ORTHOGONAL POLYNOMIALS

---

### Kernel polynomials

In the case of orthogonal polynomials on the unit circle we have a simple expression for the reproducing kernel [3; 71; 178], similar to the Christoffel-Darboux formula on the real line. The  $n$ -th polynomial kernel  $K_n(z, y)$  associated with  $\{\Phi_n\}_{n \geq 0}$  is defined by

$$\begin{aligned} K_n(z, y) &= \sum_{j=0}^n \frac{\overline{\Phi_j(y)}\Phi_j(z)}{\mathbf{k}_j} = \frac{\overline{\Phi_{n+1}^*(y)}\Phi_{n+1}^*(z) - \overline{\Phi_{n+1}(y)}\Phi_{n+1}(z)}{\mathbf{k}_{n+1}(1 - \bar{y}z)} \\ &= \frac{\overline{\phi_{n+1}^*(y)}\phi_{n+1}^*(z) - \overline{\phi_{n+1}(y)}\phi_{n+1}(z)}{1 - \bar{y}z}, \end{aligned} \quad (2.33)$$

and it satisfies the reproducing property,

$$\int_{\mathbb{T}} K_n(z, y) \overline{f(z)} d\sigma(z) = \overline{f(y)}, \quad (2.34)$$

for every polynomial  $f$  of degree at most  $n$ . Taking into account  $\phi_{n+1}^*(0) = \kappa_{n+1} \Phi_{n+1}^*(0) = \kappa_{n+1}$ , we find that

$$\Phi_n^*(z) = \frac{1}{\kappa_n^2} K_n(z, 0) = \mathbf{k}_n K_n(z, 0), \quad n \geq 0, \quad (2.35)$$

which is an expression for the reversed polynomials as a linear combination of the orthogonal polynomials up to degree  $n$ .

### GGT matrices

Using (2.35) and the forward recurrence formula (2.25), we are able to express  $z\phi_n(z)$  as a linear combination of  $\{\phi_k\}_{k=0}^{n+1}$ ,

$$z\phi_n(z) = \frac{\kappa_n}{\kappa_{n+1}} \phi_{n+1}(z) - \frac{1}{\kappa_n} \Phi_{n+1}(0) \sum_{k=0}^n \kappa_k \overline{\Phi_k(0)} \phi_k(z),$$

or, in the matrix form,

$$z\boldsymbol{\phi}(z) = \mathbf{H}_\sigma \boldsymbol{\phi}(z),$$



where  $\boldsymbol{\phi}(z) = [\phi_0(z), \phi_1(z), \dots]^T$ , and the matrix  $\mathbf{H}_\sigma$  is defined by

$$[\mathbf{H}_\sigma]_{i,j} = \langle z\phi_i, \phi_j \rangle_{\mathcal{L}} = \begin{cases} -\frac{\kappa_j}{\kappa_i} \Phi_{i+1}(0) \overline{\Phi_j(0)}, & j \leq i, \\ \frac{\kappa_i}{\kappa_{i+1}}, & j = i+1, \\ 0, & j > i+1. \end{cases}$$

This lower Hessenberg matrix [92], where the  $j$ -th row has at most its first  $j+1$  components non-zero, is called GGT representation of the multiplication by  $z$ , after [77; 86; 190].

In an analog way to the real line case, the zeros of the monic orthogonal polynomial  $\Phi_n$  are the eigenvalues of  $(\mathbf{H}_\sigma)_n$ , the  $n \times n$  principal leading sub-matrix of the GGT matrix  $\mathbf{H}_\sigma$ . Hence,  $\Phi_n$  is the characteristic polynomial of  $(\mathbf{H}_\sigma)_n$ ,

$$\Phi_n(z) = \det(z\mathbf{I}_n - (\mathbf{H}_\sigma)_n). \quad (2.36)$$

### Szegő extremum problem and $\mathcal{S}$ class

The measure of orthogonality  $d\sigma$  can be decomposed as the sum of a purely absolutely continuous measure with respect to the Lebesgue measure and a singular part. Thus, if we denote by  $\sigma'$ , the Radon-Nikodym derivative [170] of the measure  $\sigma$  supported in  $[-\pi, \pi]$ , then

$$d\sigma(\theta) = \sigma'(\theta) \frac{d\theta}{2\pi} + d\sigma_s, \quad (2.37)$$

where  $\sigma_s$  is the singular part of  $\sigma$ .

The Szegő extremum problem on the unit circle consists of finding

$$\lambda(z) = \lim_{n \rightarrow \infty} \lambda_n(z),$$

with

$$\lambda_n(z) = \inf_{f(z)=1} \left\{ \int_{-\pi}^{\pi} |f(e^{i\theta})|^2 d\sigma(\theta); f \in \mathbb{P}_n \right\}.$$

$\lambda(z)$  is said to be the Christoffel function. The solution of this problem for  $|z| < 1$  was given by Szegő in [185; 186].

In the literature, an important class of measures is the Szegő class  $\mathcal{S}$ . We summarize some relevant characterizations to the  $\mathcal{S}$  class. The following conditions are equivalent:

- i)  $\sigma \in \mathcal{S}$ .
- ii)  $\int_{-\pi}^{\pi} \log \sigma'(\theta) \frac{d\theta}{2\pi} > -\infty$ .

## 2. ORTHOGONAL POLYNOMIALS

---

$$iii) \sum_{n=0}^{\infty} |\Phi_n(0)|^2 < \infty. \quad iv) \lambda(0) = \prod_{n=0}^{\infty} (1 - |\Phi_{n+1}(0)|^2) < +\infty.$$

From this we deduce that if the measure  $\sigma$  does not belong to the  $\mathcal{S}$  class, the GGT matrix  $\mathbf{H}_\sigma$  is unitary. In general,  $\mathbf{H}_\sigma$  satisfies

$$i) \mathbf{H}_\sigma \mathbf{H}_\sigma^H = \mathbf{I}; \quad ii) \mathbf{H}_\sigma^H \mathbf{H}_\sigma = \mathbf{I} - \lambda(0) \boldsymbol{\phi}(0) \boldsymbol{\phi}(0)^H.$$

As a part of the analysis when  $\sigma \in \mathcal{S}$ , one can construct the Szegő function  $D$ , defined in  $\mathbb{D}$  as

$$D(z) = \exp\left(\frac{1}{4\pi} \int_{-\pi}^{\pi} \frac{e^{i\theta} + z}{e^{i\theta} - z} \log \sigma'(\theta) d\theta\right), \quad z \in \mathbb{D}.$$

Thus,  $|D|^2 = \sigma'$  almost everywhere on  $\mathbb{T}$ , and the solution of the Szegő extremum problem is given by

$$\lambda(z) = (1 - |z|^2) |D(z)|^2, \quad z \in \mathbb{D}.$$

### $\mathcal{N}$ class

We say that  $\sigma$  belongs to the Nevai class  $\mathcal{N}$ , if

$$\lim_{n \rightarrow \infty} \Phi_n(0) = \lim_{n \rightarrow \infty} \frac{\phi_n(0)}{\kappa_n} = 0.$$

The relation between the classes  $\mathcal{S}$  and  $\mathcal{N}$  can be viewed using the results in [136]. If  $\sigma \in \mathcal{S}$ , then it has a normal  $L^2$ -derivative behavior, i.e.,

$$\lim_{n \rightarrow \infty} \left( \int_{-\pi}^{\pi} \frac{|\phi'_n(e^{i\theta})|^2}{n^2} \sigma'(\theta) d\theta \right)^{\frac{1}{2}} = 1,$$

and thus  $\sigma \in \mathcal{N}$ . Furthermore, if  $\sigma \in \mathcal{N}$ ,

$$\left| \frac{\Phi_n(z)}{\Phi_{n-1}(z)} - z \right| \leq |\Phi_n(0)|, \quad z \in \mathbb{C} \setminus \mathbb{D}.$$

Thus,

$$\lim_{n \rightarrow \infty} \frac{\Phi_n(z)}{\Phi_{n-1}(z)} = z,$$

uniformly in compact subsets of  $\mathbb{C} \setminus \overline{\mathbb{D}}$ .

---

This result can be obtained under weaker conditions. A well known result of Rakhmanov [160] states that any probability measure  $\sigma$  with  $\sigma' > 0$  almost everywhere on  $\mathbb{T}$  belongs to the class  $\mathcal{N}$ .

### CMV matrices

The GGT matrix has several constraints. If  $\sigma \in \mathcal{S}$ ,  $\{\phi_n\}_{n \geq 0}$  is not basis on  $\Lambda$  and the matrix  $\mathbf{H}_\sigma$  is not unitary. Even more, all entries above the main diagonal and the first sub-diagonal are non-zero, and they depend on an unbounded number of Verblunsky coefficients. Consequently, the GGT matrix is somewhat difficult to manipulate. The more useful basis was discovered by Cantero, Moral, and Velazquez [29] (this result is one of the most interesting developments in the theory of orthogonal polynomials on the unit circle in recent years) as a matrix realization for the multiplication by  $z$ , with respect to the CMV orthonormal basis  $\{\chi_i\}_{i \geq 0}$ ,

$$\langle z\chi_i, \chi_j \rangle_{\mathcal{L}} = 0, \quad |i - j| > k, \quad k \geq 0.$$

In this case  $k = 2$  to be compared with  $k = 1$  for the Jacobi matrices which correspond to the real line case. The CMV basis  $\{\chi_n\}_{n \geq 0}$  is obtained by orthonormalizing  $\{1, z, z^{-1}, z^2, z^{-2}, \dots\}$  using the Gram-Schmidt process and the matrix, called the CMV matrix,

$$\mathbf{C} = \left[ \langle z\chi_i, \chi_j \rangle_{\mathcal{L}} \right]_{i, j \geq 0},$$

is five-diagonal. Its origins are outlined in [198] (see also [179]). Remarkably, the basis  $\{\chi_n\}_{n \geq 0}$  can be expressed in terms of the sequences  $\{\phi_n\}_{n \geq 0}$  and  $\{\phi_n^*\}_{n \geq 0}$ ,

$$\chi_{2n}(z) = z^{-n} \phi_{2n}^*(z), \quad \chi_{2n+1}(z) = z^{-n} \phi_{2n+1}(z), \quad n \geq 0.$$

There is an important relation between CMV and monic orthogonal polynomials as (2.36) for the GGT representation,

$$\Phi_n(z) = \det(z\mathbf{I}_n - \mathbf{C}_n),$$

where  $\mathbf{C}_n$  is the  $n \times n$  principal leading sub-matrix of the CMV matrix  $\mathbf{C}$ .

The CMV matrices play the same role in the study of orthogonal polynomials on the unit circle that Jacobi matrices in orthogonal polynomials on the real line.

## 2. ORTHOGONAL POLYNOMIALS

---

### Zeros

If  $\Phi_n$  is an orthogonal polynomial of degree  $n$ , all its zeros lie in the interior of the convex hull of the support of the measure of orthogonality [67], and we recover the properties of zeros for orthogonal polynomials on the real line. From the Christoffel-Darboux formula we have for  $z = y$ ,

$$K_n(z, z) = \sum_{k=0}^n |\phi_k(z)|^2 = \frac{|\phi_{n+1}^*(z)|^2 - |\phi_{n+1}(z)|^2}{1 - |z|^2}. \quad (2.38)$$

If  $z_{n,1}$  is a zero of  $\phi_n$  with  $|z_{n,1}| = 1$ , then using (2.38) for  $n - 1$ ,  $\phi_n^*(z_{n,1}) = 0$ , and from the recurrence relation (2.32) we get  $\phi_{n-1}(z_{n,1}) = 0$ . Repeating this argument, we have  $\phi_k(z_{n,1}) = 0$ ,  $k \leq n$ , but for  $k = 0$ ,  $\phi_0 = 1$ , which gives a contradiction. Hence we conclude that  $\phi_n$  has no zeros on the unit circle, and thus all the zeros of  $\phi_n$  are in  $\mathbb{D}$ .

In Section 2.1 we see that the interlacing property for the zeros of two polynomials  $P_{n-1}$  and  $P_n$ , means that they are the  $(n - 1)$ -st and  $n$ -th orthogonal polynomials associated with a measure  $d\mu$  supported on the real line. In the case of the unit circle, we have an analogous result, which is known in the literature as the Schur-Cohn-Jury criterion [12]. A monic polynomial  $f_n$  has its  $n$  zeros inside the unit circle if and only if the sequence of parameters  $\{h_k\}_{k=0}^n$ , defined by the following backward algorithm

$$\begin{aligned} g_n(z) &= f_n(z), \quad f_n(0) = h_n, \\ g_k(z) &= \frac{1}{z(1 - |h_{k+1}|^2)} \left( g_{k+1}(z) - h_{k+1} g_{k+1}^*(z) \right), \quad k = n - 1, n - 2, \dots, 0, \end{aligned}$$

satisfies  $|h_k| < 1$ ,  $k \geq 1$ .

### 2.2.1 C-functions and rational spectral transformations

#### C-functions

In the sequel, we consider that  $\sum_{k=0}^{\infty} c_{-k} z^k$  converges on  $|z| < r$ , for some  $r > 0$ , where  $\{c_n\}_{n \geq 0}$  is the sequence of moments (2.23). Let  $F : E \subset \mathbb{C} \rightarrow \mathbb{C}$  be a complex function associated with the linear functional  $\mathcal{L}$ , defined as follows

$$F(z) = \left\langle \mathcal{L}, \frac{y+z}{y-z} \right\rangle.$$

---

Here  $\mathcal{L}$  acts on  $y$ .  $F$  is said to be a  $C$ -function associated with the linear functional  $\mathcal{L}$ . Since  $\sum_{k=0}^{\infty} c_{-k} z^k$  converges on  $|z| \leq r$ ,  $F(z)$  is analytic in a neighborhood of  $z = 0$ , and we get the following representation of  $F(z)$  as a series expansion at  $z = 0$

$$F(z) = c_0 + 2 \sum_{k=1}^{\infty} c_{-k} z^k, \quad |z| \leq r, \quad (2.39)$$

where  $c_0 \in \mathbb{R}$  and  $c_{-k} \in \mathbb{C}$ .

If  $\mathcal{L}$  is positive definite and  $c_0 = 1$ , there is a probability measure  $\sigma$  supported on the unit circle such that  $F$  can be represented as a Riesz-Herglotz transformation [91; 165] of  $d\sigma$  as follows

$$F(z) = \int_{\mathbb{T}} \frac{y+z}{y-z} d\sigma(y). \quad (2.40)$$

A complex function  $F$  which has a representation of the form (2.40) is called a Carathéodory function [31]. It can be shown that  $F$  is a Carathéodory function if and only if  $F(z)$  is analytic in  $|z| < 1$  and  $\Re F(z) > 0$  for  $|z| < 1$ ; see Chapter 1. From this it follows immediately that  $F^{-1}$  is a Carathéodory function if and only if  $F$  is a Carathéodory function.

### Characterization of orthogonal polynomials

For a given polynomial  $\pi$  of degree  $n$  with leading coefficient  $\eta$ , the polynomial of the second kind of  $\pi$ ,  $\Pi$ , with respect to  $\mathcal{L}$ , is defined by

$$\Pi(z) = \begin{cases} \left\langle \mathcal{L}, \frac{z+y}{z-y} (\pi(y) - \pi(z)) \right\rangle, & \deg \pi > 0, \\ -\eta c_0, & \deg \pi = 0, \end{cases}$$

where  $\mathcal{L}$  acts on  $y$ .

Next, we give necessary and sufficient conditions for a polynomial to be orthogonal with respect to a linear, not necessarily positive definite functional  $\mathcal{L}$ , with the help of the associated  $C$ -function  $F$ . The following statement holds [155]:

$$\langle \Phi_n(z), z^k \rangle_{\mathcal{L}} = 0, \quad k = 0, \dots, n-1,$$

## 2. ORTHOGONAL POLYNOMIALS

---

if and only if

$$\Phi_n(z)F(z) - \Omega_n(z) = \mathcal{O}(z^n), \quad |z| < 1, \quad (2.41)$$

$$\Phi_n^*(z)F(z) + \Omega_n^*(z) = \mathcal{O}(z^{n+1}), \quad |z| < 1, \quad (2.42)$$

where  $\Omega_n$  is the polynomial of second kind associated with  $\Phi_n$  with respect to  $\mathcal{L}$ .

There is an interesting way to rephrase the previous result, namely, given a linear functional  $\mathcal{L}$  and its  $C$ -function  $F$ , and given  $Q_n$ , a monic polynomial of degree  $n$ , define  $\Omega_n$ , a monic polynomial, by (2.41). Then,  $Q_n = \Phi_n$  if and only if (2.42) holds.

### Spectral transformations

As in the real line case, by a spectral transformation of  $F$  we mean a new  $C$ -function associated with the hermitian linear functional  $\tilde{\mathcal{L}}$ , a modification of  $\mathcal{L}$ , such that

$$\tilde{F}(z) = \tilde{c}_0 + 2 \sum_{k=1}^{\infty} \tilde{c}_{-k} z^k, \quad |z| \leq r,$$

where  $\{\tilde{c}_n\}_{n \geq 0}$  is the sequence of transformed moments. Following the definition of Zhedanov [202] we refer to rational spectral transformation as a transformation of a  $C$ -function  $F$  given by

$$\tilde{F}(z) = \frac{A(z)F(z) + B(z)}{C(z)F(z) + D(z)}, \quad A(z)D(z) - B(z)C(z) \neq 0, \quad (2.43)$$

where  $A$ ,  $B$ ,  $C$ , and  $D$  are coprime polynomials or Laurent polynomials. If  $C = 0$ , and  $A$ ,  $B$ ,  $C$  are polynomials or Laurent polynomials that provide a true asymptotic behavior to (2.43), we have the linear spectral transformations. Following our classification in terms of the moments, the local spectral transformation is a special case of general linear spectral transformation related to perturbations on two symmetric sub-diagonals of the Toeplitz matrix (2.24).

### Linear spectral transformations

We use the  $C$ -function  $F$  as a main tool in the investigation of spectral transformations of orthogonal polynomials. As an analog of the Christoffel transformation (2.16)

---

for  $\mathcal{S}$ -functions, we consider a perturbation  $\mathcal{L}_R$  of  $\mathcal{L}$  defined by

$$\langle \mathcal{L}_R, f \rangle = \left\langle \mathcal{L}, \frac{1}{2} \left( z - \alpha + z^{-1} - \bar{\alpha} \right) f(z) \right\rangle, \quad f \in \Lambda, \quad \alpha \in \mathbb{C}. \quad (2.44)$$

It constitutes an example of linear spectral transformation and it was introduced in [28; 183]. It is natural to analyze the existence of the inverse transformation, i.e., if there exists a linear functional  $\mathcal{L}_{R(-)}$  such that

$$\left\langle \mathcal{L}_{R(-)}, \left( z - \alpha + z^{-1} - \bar{\alpha} \right) f(z) \right\rangle = \langle \mathcal{L}, f \rangle, \quad f \in \Lambda, \quad (2.45)$$

as well as to analyze if the quasi-definite character of the linear functional is preserved by such a transformation. Notice that this transformation does not define a unique linear functional  $\mathcal{L}_{R(-)}$ . The uniqueness depends on a free parameter. Recently, the spectral transformations (2.44) and (2.45) have been studied with a new approach in the framework of inverse problems for sequences of monic orthogonal polynomials [30].

For all values of  $\alpha$ , such that  $|\Re(\alpha)| > 1$ , the Laurent polynomial  $z - \alpha + z^{-1} - \bar{\alpha}$  can be represented as a polynomial of the form  $-\frac{1}{\beta}|z - \beta|^2$ , where  $\beta \in \mathbb{R} \setminus \{0\}$ . The particular cases (2.44) and (2.45) with  $|\Re(\alpha)| > 1$  have been extensively covered in [72] and the references therein.

## Rational spectral transformations

Two remarkable examples of rational spectral transformations are due to Peherstorfer [152]. We denote by  $\{\Phi_n^{(k)}\}_{n \geq 0}$  the  $k$ -th associated sequence of polynomials of order  $k \geq 1$  for the monic orthogonal sequence  $\{\Phi_n\}_{n \geq 0}$ , that constitutes the analog of the associated polynomials, satisfying (2.20). In this case they are generated by the recurrence relation

$$\Phi_{n+1}^{(k)}(z) = z\Phi_n^{(k)}(z) + \Phi_{n+k+1}(0)\left(\Phi_n^{(k)}(z)\right)^*, \quad n \geq 0. \quad (2.46)$$

Notice that  $\{\Phi_n^{(k)}\}_{n \geq 0}$  is again a sequence of orthogonal polynomials with respect to a new hermitian linear functional such that  $F^{(k)}$  is the corresponding  $\mathcal{C}$ -function.

Denoting by  $\mathcal{F}^{(k)}[F(z)] = F^{(k)}(z)$  the forward transformation of  $F$ , the correspond-

## 2. ORTHOGONAL POLYNOMIALS

---

ing  $C$ -function is a rational spectral transformation given by

$$F^{(k)}(z) = \frac{(\Phi_k(z) + \Phi_k^*(z))F(z) - \Omega_k(z) + \Omega_k^*(z)}{(\Phi_k(z) - \Phi_k^*(z))F(z) - \Omega_k(z) - \Omega_k^*(z)},$$

where  $\Omega_n$  is the polynomial of second kind of  $\Phi_n$  with respect to  $\mathcal{L}$ . In other words, we remove the first  $k$  Verblunsky coefficients from the original sequence.

On the other hand, if we add complex numbers  $z_1, z_2, \dots, z_k$  with  $|z_i| \neq 1$ ,  $1 \leq i \leq k$ , to the original sequence of Verblunsky coefficients, we have the backward associated sequence of polynomials  $\{\Phi_n^{(-k)}\}_{n \geq 0}$  as a sequence of monic orthogonal polynomials generated by  $\{z_i\}_{i=1}^k \cup \{\phi_n(0)\}_{n \geq 1}$ .

Denoting by  $\mathcal{F}^{(-k)}[F(z)] = F^{(-k)}(z)$  the backward transformation of  $F$ , the corresponding  $C$ -function is a rational spectral transformation given by

$$F^{(-k)}(z) = \frac{-(\widetilde{\Omega}_k(z) + \widetilde{\Omega}_k^*(z))F(z) - \widetilde{\Omega}_k^*(z) + \widetilde{\Omega}_k(z)}{(\widetilde{\Phi}_k^*(z) - \widetilde{\Phi}_k(z))F(z) + \widetilde{\Phi}_k(z) + \widetilde{\Phi}_k^*(z)},$$

where  $\widetilde{\Phi}_k$  (respectively  $\widetilde{\Omega}_k$ ) is the  $k$ -th degree polynomial generated using the complex numbers  $z_1, z_2, \dots, z_k$  (respectively  $-z_1, -z_2, \dots, -z_k$ ) through the recurrence relation (2.46), i.e.,  $\widetilde{\Omega}_k$  is the polynomial of second kind associated with  $\widetilde{\Phi}_k$ . It is easily verified that  $\mathcal{F}^{(k)} \circ \mathcal{F}^{(-k)} = \mathcal{I}$ . Generally, the inverse is not always true since it depends on the choice of free parameters.

### 2.2.2 Connection with orthogonal polynomials on $[-1, 1]$

Given a non-trivial probability measure  $\mu$ , supported on the interval  $[-1, 1]$ , then there exists a sequence of orthonormal polynomials  $\{p_n\}_{n \geq 0}$  such that

$$\int_{-1}^1 p_n(x)p_m(x)d\mu(x) = \delta_{n,m}.$$

We can define a measure  $\sigma$  supported on  $[-\pi, \pi]$  such that

$$d\sigma(\theta) = \frac{1}{2}|d\mu(\cos \theta)|.$$



---

In particular, if  $\mu$  is an absolutely continuous measure, i.e.,  $d\mu(x) = \omega(x)dx$ , then

$$d\sigma(\theta) = \frac{1}{2}\omega(\cos\theta)|\sin\theta|d\theta.$$

This is the so-called Szegő transformation of probability measures supported on  $[-1, 1]$  to probability measures supported on  $\mathbb{T}$ .

If  $\mu$  is a non-trivial probability measure on  $[-1, 1]$  (this is the reason why we introduced the factor  $1/2$ ),  $\sigma$  is also a symmetric probability measure on the unit circle and, as a consequence, there exists a sequence of orthonormal polynomials  $\{\phi_n\}_{n \geq 0}$  such that

$$\int_{-\pi}^{\pi} \phi_n(e^{i\theta}) \overline{\phi_m(e^{i\theta})} d\sigma(\theta) = \delta_{n,m},$$

as well as the corresponding sequence of monic orthogonal polynomials. In this case,

$$\Phi_n(0) \in (-1, 1), \quad n \geq 1.$$

There is a relation between the sequence of orthogonal polynomials associated with a measure  $\mu$  supported on  $[-1, 1]$  and the sequence of orthogonal polynomials associated with the measure  $\sigma$  supported on the unit circle. The sequence of orthogonal polynomials  $\{\Phi_n\}_{n \geq 0}$  on the unit circle associated with the measure  $\sigma$  has real coefficients. In addition, if  $2x = z + z^{-1}$ , so that  $x = \cos\theta$  corresponds to  $z = e^{i\theta}$ , then

$$p_n(x) = \frac{\kappa_{2n}}{\sqrt{2(1 + \Phi_{2n}(0))}} (z^{-n} \Phi_{2n}(z) + z^n \Phi_{2n}(1/z)). \quad (2.47)$$

From (2.47) one can obtain a relation between the coefficients of the recurrence relations (2.5) and (2.25)-(2.26),

$$\begin{aligned} 2a_n &= \sqrt{(1 - \Phi_{2n}(0))(1 - \Phi_{2n-1}^2(0))(1 + \Phi_{2n-2}(0))}, \quad n \geq 1, \\ 2b_n &= \Phi_{2n-1}(0)(1 - \Phi_{2n}(0)) - \Phi_{2n+1}(0)(1 + \Phi_{2n}(0)), \quad n \geq 0. \end{aligned}$$

Conversely, if  $R_n = P_{n+1}/P_n$ , then

$$\Phi_{2n}(0) = R_n(1) - R_n(-1) - 1, \quad \Phi_{2n+1}(0) = \frac{R_n(1) + R_n(-1)}{R_n(1) - R_n(-1)}.$$

There is also a relation between the  $\mathcal{S}$ -function and  $\mathcal{C}$ -function associated with  $\mu$

## 2. ORTHOGONAL POLYNOMIALS

---

and  $\sigma$ , respectively, as follows

$$F(z) = \frac{1-z^2}{2z} \int_{-1}^1 \frac{d\mu(y)}{x-y} = \frac{1-z^2}{2z} S(x),$$

or, equivalently,

$$S(x) = \frac{F(z)}{\sqrt{x^2-1}},$$

with  $2x = z + z^{-1}$  and  $z = x - \sqrt{x^2-1}$ .

## Chapter 3

# On special classes of Szegő polynomials

The notion of Schur parameters <sup>1</sup> is fundamental not only in the theory of orthogonal polynomials . . .

— A. Mukaihira and Y. Nakamura. [145]

From Chapter 2 we know that the Verblunsky coefficients  $\{\Phi_n(0)\}_{n \geq 1}$  associated with a measure supported on the unit circle satisfy (2.31). Moreover, by the Favard theorem, any sequence obeying (2.31) arises as the Verblunsky coefficients of a unique non-trivial probability measure supported on the unit circle. An interesting question arises: What happens when  $|\Phi_n(0)| > 1$  at most for some  $n$ ? Clearly, these polynomials can not be orthogonal on the unit circle. In 1999, Vinet and Zhedanov [194] constructed special classes of Szegő polynomials when  $|\Phi_n(0)| > 1$ ,  $n \geq 1$ . This situation is more interesting, because there are sequences  $\{\Phi_n(0)\}_{n \geq 1}$  for which the moment problem is indeterminate. They consider two possible choices of the Verblunsky coefficients when the support of the associated measure lies on the real line. They note that if  $\Phi_n(0) > 1$ ,  $n \geq 1$ , the corresponding orthogonality measure is supported on the negative half side of the real line. On the other hand, if  $(-1)^n \Phi_n(0) > 1$ ,  $n \geq 1$ , the corresponding orthogonality measure is supported on the positive half side side of the real line. Their main tool is a mapping from symmetric polynomials on the real line to the Szegő

---

<sup>1</sup>Verblunsky coefficients.

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

polynomials [58; 60]; see also Chapter 1.

The aim of this chapter is to study the properties of the sequence of monic orthogonal polynomials  $\{\Phi_n\}_{n \geq 0}$ , which satisfy the same recurrence relation as orthogonal polynomials on the unit circle (Szegő orthogonality) with

$$\Phi_n(0) \in \mathbb{C}, \quad |\Phi_n(0)| > 1, \quad n \geq 1.$$

In our development a central role is played by continued fractions. We emphasize the natural parallelism existing between the theories of orthogonal polynomials on the real line and those on the unit circle. An analysis of the Perron-Carathéodory continued fractions [97] shows that these polynomials satisfy the Szegő orthogonality, where the linear functional  $\mathcal{L}$  defined in (2.23) satisfies

$$\bar{c}_{-n} = c_n, \quad (-1)^{n(n+1)/2} \det \mathbf{T}_n > 0, \quad n \geq 0.$$

The relations between the polynomials  $\Phi_n(\omega; z) = \Phi_n(z) - \omega z \Phi_{n-1}(z)$ ,  $\omega \neq 1$ , and the para-orthogonal polynomials [97],  $\Phi_n(z) + \tau \Phi_n^*(z)$ ,  $|\tau| = 1$  are also analysed. In the two particular cases considered in [194], the zeros of the Szegő polynomials, those of para-orthogonal polynomials, and the associated quadrature rules have been studied. As a consequence of this study, we solve the moment problem (1.11) associated with the Chebyshev polynomials of the first kind. This chapter could be considered in many aspects as a continuation of the introductory theory of orthogonal polynomials, using a different approach to the subject based on continued fractions and their modified approximates.

#### 3.1 Special class of Szegő polynomials

The treatment of this section is similar to that is given in [99], where the authors assume that  $|\Phi_n(0)| \neq 1$ ,  $n \geq 1$ . The results given here, in addition to making the chapter self-contained, help the reader to see the specific properties satisfied by the associated Toeplitz and Hankel determinants when  $|\Phi_n(0)| > 1$ ,  $n \geq 1$ .

### 3.1.1 Szegő polynomials from continued fractions

We start with the so-called Perron-Carathéodory continued fraction

$$\beta_0 - \frac{2\beta_0}{1} - \frac{1}{\bar{\alpha}_1 z} - \frac{(|\alpha_1|^2 - 1)z}{\alpha_1} - \frac{1}{\bar{\alpha}_2 z} - \dots, \quad (3.1)$$

where we assume that  $\beta_0 > 0$  and  $|\alpha_n| > 1$ ,  $n \geq 1$ . This continued fraction was introduced in [97]; see also [98; 99]. If we compare (3.1) with (2.8), then we have  $q_0 = \beta_0$ ,  $r_1 = -2\beta_0$ ,  $q_1 = 1$ ,  $r_{2n} = -1$ ,  $r_{2n+1} = -(|\alpha_n|^2 - 1)z$ ,  $q_{2n} = \bar{\alpha}_n z$ , and  $q_{2n+1} = \alpha_n$ ,  $n \geq 1$ .

Let  $\{A_n\}_{n \geq 0}$  and  $\{B_n\}_{n \geq 0}$  be the sequence of numerator and denominator polynomials of (3.1), respectively. Then these polynomials satisfy the recurrence relations

$$\begin{bmatrix} A_{2n}(z) \\ B_{2n}(z) \end{bmatrix} = \bar{\alpha}_n z \begin{bmatrix} A_{2n-1}(z) \\ B_{2n-1}(z) \end{bmatrix} - \begin{bmatrix} A_{2n-2}(z) \\ B_{2n-2}(z) \end{bmatrix}, \quad n \geq 1, \quad (3.2)$$

$$\begin{bmatrix} A_{2n+1}(z) \\ B_{2n+1}(z) \end{bmatrix} = \alpha_n \begin{bmatrix} A_{2n}(z) \\ B_{2n}(z) \end{bmatrix} - (|\alpha_n|^2 - 1)z \begin{bmatrix} A_{2n-1}(z) \\ B_{2n-1}(z) \end{bmatrix}, \quad n \geq 1, \quad (3.3)$$

where  $B_0 = 1$ ,  $B_1 = 1$ ,  $A_0 = \beta_0$ , and  $A_1 = -\beta_0$ . From these recurrence relations, since  $\alpha_n \neq 0$ ,  $n \geq 1$ , we get

$$\begin{bmatrix} A_{2n+2}(z) \\ B_{2n+2}(z) \end{bmatrix} = \left( \frac{\bar{\alpha}_{n+1}}{\bar{\alpha}_n} z - 1 \right) \begin{bmatrix} A_{2n}(z) \\ B_{2n}(z) \end{bmatrix} - \frac{\bar{\alpha}_{n+1}}{\bar{\alpha}_n} (|\alpha_n|^2 - 1)z \begin{bmatrix} A_{2n-2}(z) \\ B_{2n-2}(z) \end{bmatrix}, \quad n \geq 1,$$

with  $B_0 = 1$ ,  $B_2(z) = \bar{\alpha}_1 z - 1$ ,  $A_0 = \beta_0$ ,  $A_2(z) = -\beta_0(\bar{\alpha}_1 z + 1)$ , and

$$\begin{bmatrix} A_{2n+3}(z) \\ B_{2n+3}(z) \end{bmatrix} = \left( z - \frac{\alpha_{n+1}}{\alpha_n} \right) \begin{bmatrix} A_{2n+1}(z) \\ B_{2n+1}(z) \end{bmatrix} - \frac{\alpha_{n+1}}{\alpha_n} (|\alpha_n|^2 - 1)z \begin{bmatrix} A_{2n-1}(z) \\ B_{2n-1}(z) \end{bmatrix}, \quad n \geq 1.$$

Here  $B_1 = 1$ ,  $B_3(z) = z - \alpha_1$ ,  $A_1 = -\beta_0$ , and  $A_3(z) = -\beta_0(z + \alpha_1)$ . From these recurrence relations one can easily observe that both  $B_{2n}$  and  $B_{2n+1}$  are polynomials of exact degree  $n$ . Likewise, both  $A_{2n}$  and  $A_{2n+1}$  are polynomials of exact degree  $n$ . More precisely,

$$\begin{aligned} B_{2n}(z) &= \bar{\alpha}_n z^n + \dots + (-1)^n, & B_{2n+1}(z) &= z^n + \dots + (-1)^n \alpha_n, \\ A_{2n}(z) &= -\beta_0(\bar{\alpha}_n z^n + \dots + (-1)^{n-1}), & A_{2n+1}(z) &= -\beta_0(z^n + \dots + (-1)^{n-1} \alpha_n), \quad n \geq 1. \end{aligned}$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

Using the above recurrence relations one can also easily conclude that

$$A_{2n+1}^*(z) = z^n \overline{A_{2n+1}(1/\bar{z})} = z^n \bar{A}_{2n+1}(1/z) = (-1)^{n+1} A_{2n}(z), \quad n \geq 0, \quad (3.4)$$

$$B_{2n+1}^*(z) = z^n \overline{B_{2n+1}(1/\bar{z})} = z^n \bar{B}_{2n+1}(1/z) = (-1)^n B_{2n}(z), \quad n \geq 0. \quad (3.5)$$

Moreover, from these recurrence relations we obtain

$$\frac{A_{2n+2}(z)}{B_{2n+2}(z)} - \frac{A_{2n}(z)}{B_{2n}(z)} = \begin{cases} \left( 2\beta_0 \bar{\alpha}_{n+1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^{n+1} + \mathcal{O}(z^{n+2}), \\ - \left( 2\beta_0 \bar{\alpha}_n^{-1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^{-n} + \mathcal{O}(z^{-(n+1)}), \end{cases} \quad n \geq 1,$$

$$\frac{A_{2n+3}(z)}{B_{2n+3}(z)} - \frac{A_{2n+1}(z)}{B_{2n+1}(z)} = \begin{cases} \left( 2\beta_0 \alpha_n^{-1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^n + \mathcal{O}(z^{n+1}), \\ - \left( 2\beta_0 \alpha_{n+1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^{-(n+1)} + \mathcal{O}(z^{-(n+2)}), \end{cases} \quad n \geq 1,$$

$$\frac{A_{2n+1}(z)}{B_{2n+1}(z)} - \frac{A_{2n}(z)}{B_{2n}(z)} = \begin{cases} - \left( 2\beta_0 \alpha_n^{-1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^n + \mathcal{O}(z^{n+1}), \\ - \left( 2\beta_0 \bar{\alpha}_n^{-1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^{-n} + \mathcal{O}(z^{-(n+1)}), \end{cases} \quad n \geq 1.$$

Thus, there exists a pair of formal power series

$$F(z) = c_0 + 2 \sum_{n=1}^{\infty} c_{-n} z^n, \quad |z| < 1, \quad F_{\infty}(z) = -c_0 - 2 \sum_{n=1}^{\infty} c_n z^{-n}, \quad |z| > 1, \quad (3.6)$$

with  $c_0 = \beta_0$  and  $\bar{c}_{-1} = c_1 = \beta_0 \alpha_1$ , such that

$$F(z) - \frac{A_{2n}(z)}{B_{2n}(z)} = \left( 2\beta_0 \bar{\alpha}_{n+1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^{n+1} + \mathcal{O}(z^{n+2}), \quad n \geq 1,$$

$$F_{\infty}(z) - \frac{A_{2n}(z)}{B_{2n}(z)} = - \left( 2\beta_0 \bar{\alpha}_n^{-1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^{-n} + \mathcal{O}(z^{-(n+1)}), \quad n \geq 1,$$

$$F(z) - \frac{A_{2n+1}(z)}{B_{2n+1}(z)} = \left( 2\beta_0 \alpha_n^{-1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^n + \mathcal{O}(z^{n+1}), \quad n \geq 1, \quad (3.7)$$

$$F_{\infty}(z) - \frac{A_{2n+1}(z)}{B_{2n+1}(z)} = - \left( 2\beta_0 \alpha_{n+1} \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) z^{-(n+1)} + \mathcal{O}(z^{-(n+2)}), \quad n \geq 1.$$

That is, (3.1) corresponds to the formal power series expansions  $F$  and  $F_\infty$ . Using the reciprocal properties (3.4)-(3.5), we also conclude that these formal power series expansions are such that

$$\bar{c}_{-n} = c_n, \quad n \geq 1. \quad (3.8)$$

If we write  $B_{2n+1}(z) = \sum_{r=0}^n b_{n,r} z^r$  and  $A_{2n+1}(z) = \sum_{r=0}^n a_{n,r} z^r$ , then from (3.7),

$$F(z) \sum_{r=0}^n b_{n,r} z^r - \sum_{r=0}^n a_{n,r} z^r = \gamma_n z^n + O(z^{n+1}), \quad n \geq 1,$$

$$F_\infty(z) \sum_{r=0}^n b_{n,r} z^{r-n} - \sum_{r=0}^n a_{n,r} z^{r-n} = O(z^{-n-2}), \quad n \geq 1,$$

where  $\gamma_n = (-1)^n 2\beta_0 \prod_{r=1}^n (|\alpha_r|^2 - 1)$ . This leads to two systems of  $n+1$  linear equations and  $n+1$  unknowns as follows

$$\mathbf{A}_1 \mathbf{b} = \mathbf{a} + \mathbf{c}, \quad \mathbf{A}_2 \mathbf{b} = -\mathbf{a},$$

where  $\mathbf{a} = [a_{n,0}, a_{n,1}, \dots, a_{n,n-1}, a_{n,n}]^T$ ,  $\mathbf{b} = [b_{n,0}, b_{n,1}, \dots, b_{n,n-1}, b_{n,n}]^T$ ,  $\mathbf{c} = [0, 0, \dots, 0, \gamma_n]^T$ ,

$$\mathbf{A}_1 = \begin{bmatrix} c_0 & 0 & \cdots & 0 & 0 \\ 2c_{-1} & c_0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 2c_{-n+1} & 2c_{-n+2} & \cdots & c_0 & 0 \\ 2c_{-n} & 2c_{-n+1} & \cdots & 2c_{-1} & c_0 \end{bmatrix}, \quad \mathbf{A}_2 = \begin{bmatrix} c_0 & 2c_1 & \cdots & 2c_{n-1} & 2c_n \\ 0 & c_0 & \cdots & 2c_{n-2} & 2c_{n-1} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & c_0 & 2c_1 \\ 0 & 0 & \cdots & 0 & c_0 \end{bmatrix},$$

or, equivalently, the linear system

$$\begin{bmatrix} c_0 & c_1 & \cdots & c_{n-1} & c_n \\ c_{-1} & c_0 & \cdots & c_{n-2} & c_{n-1} \\ \vdots & \vdots & & \vdots & \vdots \\ c_{-n+1} & c_{-n+2} & \cdots & c_0 & c_1 \\ c_{-n} & c_{-n+1} & \cdots & c_{-1} & c_0 \end{bmatrix} \begin{bmatrix} b_{n,0} \\ b_{n,1} \\ \vdots \\ b_{n,n-1} \\ b_{n,n} \end{bmatrix} = \frac{1}{2} \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 0 \\ \gamma_n \end{bmatrix}. \quad (3.9)$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

Applying Cramer's rule for the coefficient  $b_{n,n} = 1$  we get

$$\det \mathbf{T}_0 = c_0 = \beta_0, \quad \det \mathbf{T}_n = \left[ (-1)^n \beta_0 \prod_{r=1}^n (|\alpha_r|^2 - 1) \right] \det \mathbf{T}_{n-1}, \quad n \geq 1.$$

Therefore,

$$\begin{aligned} \det \mathbf{T}_n &= \left( (-1)^n \beta_0 \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) \left( (-1)^{n-1} \beta_0 \prod_{r=1}^{n-1} (|\alpha_r|^2 - 1) \right) \det \mathbf{T}_{n-1} = \dots, \\ &= \left( (-1)^n \beta_0 \prod_{r=1}^n (|\alpha_r|^2 - 1) \right) \left( (-1)^{n-1} \beta_0 \prod_{r=1}^{n-1} (|\alpha_r|^2 - 1) \right) \dots \left( (-1) \beta_0 \prod_{r=1}^1 (|\alpha_r|^2 - 1) \beta_0 \right), \\ &= (-1)^{n(n+1)/2} \beta_0^{n+1} \prod_{m=1}^n \left( \prod_{r=1}^m (|\alpha_r|^2 - 1) \right). \end{aligned}$$

Hence, if  $\beta_0 > 0$ , then

$$\det \mathbf{T}_0 = \beta_0 > 0, \quad (-1)^{n(n+1)/2} \det \mathbf{T}_n = \beta_0^{n+1} \prod_{m=1}^n \left[ \prod_{r=1}^m (|\alpha_r|^2 - 1) \right] > 0, \quad n \geq 1.$$

Since  $(-1)^{n(n+1)/2} = (-1)^{\lfloor (n+1)/2 \rfloor}$ , if we look at the Hankel determinants

$$\det \mathbf{H}_1^{(0)} = c_0, \quad \det \mathbf{H}_{n+1}^{(-n)} = \begin{vmatrix} c_{-n} & c_{-n+1} & \cdots & c_0 \\ c_{-n+1} & c_{-n+2} & \cdots & c_1 \\ \vdots & \vdots & & \vdots \\ c_0 & c_1 & \cdots & c_n \end{vmatrix},$$

which can be obtained by a rearrangement of rows of the Toeplitz determinants  $\det \mathbf{T}_n$ , then we have

$$\det \mathbf{H}_{n+1}^{(-n)} > 0, \quad n \geq 0. \quad (3.10)$$

Applying Cramer's rule in (3.9) for the coefficient  $b_{n,0} = (-1)^n \alpha_n$ , we obtain

$$\alpha_n = \frac{1}{\det \mathbf{T}_{n-1}} \begin{vmatrix} c_1 & \cdots & c_n \\ c_0 & \cdots & c_{n-1} \\ \vdots & & \vdots \\ c_{-n+2} & \cdots & c_1 \end{vmatrix} = \frac{1}{\det \mathbf{T}_{n-1}} \begin{vmatrix} c_{-1} & \cdots & c_{n-2} \\ c_{-2} & \cdots & c_{n-3} \\ \vdots & & \vdots \\ c_{-n} & \cdots & c_{-1} \end{vmatrix}.$$



---

The above determinant expression follows by considering the transpose and then using (3.8). Hence, through an interchanging of rows in the determinants,

$$\det \mathbf{H}_n^{(-n+2)} = \overline{\det \mathbf{H}_n^{(-n)}} = \alpha_n \det \mathbf{H}_n^{(-n+1)}, \quad n \geq 1. \quad (3.11)$$

Since  $|\alpha_n| > 1$ , this means

$$\left| \det \mathbf{H}_n^{(-n+2)} \right| = \left| \det \mathbf{H}_n^{(-n)} \right| > \det \mathbf{H}_n^{(-n+1)} > 0, \quad n \geq 1.$$

Now, with respect to the formal power series expansions  $F$  and  $F_\infty$  in (3.6), we define the linear functional  $\mathcal{L}$  on  $\Lambda$ . If  $l_{r,s}(z) = \sum_{m=r}^s \lambda_m z^m$ ,  $\lambda_m \in \mathbb{C}$  and  $-\infty < r \leq s < \infty$ , then

$$\langle \mathcal{L}, l_{r,s} \rangle = \sum_{m=r}^s \lambda_m c_m.$$

Therefore, from (3.9), we have for the monic polynomials  $\Phi_n = B_{2n+1}$ ,

$$\langle \Phi_n(z), z^m \rangle_{\mathcal{L}} = \frac{1}{2} \gamma_n \delta_{n,m}, \quad 0 \leq m \leq n, \quad n \geq 1.$$

Also, from (3.2) and (3.4)-(3.5),

$$\Phi_n(z) = z\Phi_{n-1}(z) + (-1)^n \alpha_n \Phi_{n-1}^*(z), \quad n \geq 1,$$

with  $\Phi_0 = 1$ ,  $\Phi_1(z) = z - \alpha_1$ . Based on these facts, we refer to  $\{B_{2n+1}\}_{n \geq 0}$  as the sequence of Szegő polynomials generated by the Perron-Carathéodory continued fraction (3.1).

### 3.1.2 Szegő polynomials from series expansions

Under the conditions (3.8) and (3.10),  $\mathcal{L}$  is referred to as an sq-definite moment functional, meaning 'special quasi-definite moment functional'. When the moments are also real, the name rsq-definite moment functional is used.

The sequence of polynomials  $\{\Phi_n\}_{n \geq 0}$  is defined by  $\Phi_0 = 1$ , and  $\Phi_n$  a monic polynomial of degree  $n$  orthogonal with respect to the sq-definite linear functional  $\mathcal{L}$ . Then, with the condition (3.10), these polynomials always exist. In fact, we can easily obtain

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

an analog result to (2.6),

$$\Phi_n(z) = \frac{1}{\det \mathbf{T}_{n-1}} \begin{vmatrix} c_0 & c_1 & \cdots & c_{n-1} & c_n \\ c_{-1} & c_0 & \cdots & c_{n-2} & c_{n-1} \\ \vdots & \vdots & & \vdots & \\ c_{-n+1} & c_{-n+2} & \cdots & c_0 & c_1 \\ 1 & z & \cdots & z^{n-1} & z^n \end{vmatrix}, \quad n \geq 1. \quad (3.12)$$

Moreover,

$$\langle \Phi_n(z), z^m \rangle_{\mathcal{L}} = \langle \Phi_n^*(z), z^{n-m} \rangle_{\mathcal{L}} = \frac{\det \mathbf{T}_n}{\det \mathbf{T}_{n-1}} \delta_{n,m}, \quad m \leq n, \quad n \geq 1. \quad (3.13)$$

The information on  $\langle \Phi_n^*(z), z^{n-m} \rangle_{\mathcal{L}}$  follows from (3.8), and with this symmetric property we have

$$\overline{\langle l_{r,s}(z), z^m \rangle_{\mathcal{L}}} = \langle z^m, l_{r,s}(z) \rangle_{\mathcal{L}}, \quad l_{r,s} \in \Lambda. \quad (3.14)$$

We call  $\{\Phi_n\}_{n \geq 0}$  the sequence of Szegő polynomials associated with the sq-definite moment functional  $\mathcal{L}$ .

**Theorem 3.1.1.** *Let  $\alpha_n = (-1)^n \Phi_n(0)$ ,  $n \geq 1$ , where  $\{\Phi_n\}_{n \geq 0}$  is the sequence of Szegő polynomials associated with the sq-definite moment functional  $\mathcal{L}$ . Then the following statements hold:*

i) *The polynomials  $\{\Phi_n\}_{n \geq 0}$  satisfy the recurrence relations*

$$\begin{aligned} \Phi_n^*(z) &= (-1)^n \bar{\alpha}_n z \Phi_{n-1}(z) + \Phi_{n-1}^*(z), \quad n \geq 1, \\ \Phi_n(z) &= (-1)^n \alpha_n \Phi_n^*(z) - (|\alpha_n|^2 - 1) z \Phi_{n-1}(z), \quad n \geq 1. \end{aligned}$$

ii)  $|\alpha_n| > 1$ ,  $n \geq 1$ .

iii) *If we set  $B_{2n}(z) = (-1)^n \Phi_n^*(z)$ ,  $B_{2n+1}(z) = \Phi_n(z)$ ,  $n \geq 0$ , then  $\{B_n\}_{n \geq 0}$  is the sequence of the denominator polynomials of (3.1).*

*Proof.* Set

$$A(z) = \Phi_n^*(z) - \alpha_n z \Phi_{n-1}(z) - \Phi_{n-1}^*(z), \quad n \geq 1,$$

where  $\alpha_n = -\langle \Phi_{n-1}^*(z), z^n \rangle_{\mathcal{L}} / \langle \Phi_{n-1}(z), z^{n-1} \rangle_{\mathcal{L}}$ . We consider  $A$  as a polynomial of de-

gree  $n$ , and we will show that it is identically zero. Clearly,

$$\langle A, z^m \rangle_{\mathcal{L}} = 0, \quad 1 \leq m \leq n. \quad (3.15)$$

Since  $A(0) = 0$ , we can write  $A^*(z) = \sum_{k=0}^{n-1} a_{n,k} \Phi_k(z)$ , and hence

$$A(z) = \sum_{k=0}^{n-1} \bar{a}_{n,k} z^{n-k} \Phi_k^*(z).$$

Using here the results of (3.15) for  $m = n, m = n - 1$  until  $m = 1$  and noting at each stage that  $\langle \mathcal{L}, \Phi_{n-m}^* \rangle \neq 0$ , we successively obtain that  $a_0 = 0, a_1 = 0$  until  $a_{n-1} = 0$ . Hence,  $A = 0$  and thus

$$\Phi_n^*(z) = \alpha_n z \Phi_{n-1}(z) + \Phi_{n-1}^*(z).$$

Comparing the coefficients of  $z^n$ , we obtain  $\bar{\alpha}_n = \Phi_n(0)$ , thus the first of the recurrence relations follows.

Now, to prove the second recurrence relation, we set

$$A(z) = \Phi_n(z) - (-1)^n \alpha_n \Phi_n^*(z) + \alpha_n z \Phi_{n-1}(z), \quad n \geq 1$$

with  $\alpha_n = -\langle \Phi_n(z), z^n \rangle_{\mathcal{L}} / \langle \Phi_{n-1}(z), z^{n-1} \rangle_{\mathcal{L}}$ . Hence,

$$\langle A, z^m \rangle_{\mathcal{L}} = 0, \quad 1 \leq m \leq n.$$

Since  $A(0) = 0$ , we can follow the same procedure as above and we get  $A = 0$ . Thus,

$$\Phi_n(z) = (-1)^n \alpha_n \Phi_n^*(z) - \alpha_n z \Phi_{n-1}(z).$$

Comparing the coefficients of  $z^n$ , we obtain  $\bar{\alpha}_n = |\alpha_n|^2 - 1$ , thus the second of the recurrence relations is deduced.

Since (2.27), we obtain from the above recurrence relation,

$$\begin{aligned} \frac{\det \mathbf{T}_n}{\det \mathbf{T}_{n-1}} &= \langle \Phi_n(z), z^n \rangle_{\mathcal{L}} = (-1)^n \alpha_n \langle \Phi_n^*(z), z^n \rangle_{\mathcal{L}} - (|\alpha_n|^2 - 1) \langle \Phi_{n-1}(z), z^{n-1} \rangle_{\mathcal{L}} \\ &= -(|\alpha_n|^2 - 1) \frac{\det \mathbf{T}_{n-1}}{\det \mathbf{T}_{n-2}}, \quad n \geq 1. \end{aligned}$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

Thus,

$$|\alpha_n|^2 - 1 = -\frac{\det \mathbf{T}_n \det \mathbf{T}_{n-2}}{\det \mathbf{T}_{n-1}^2}, \quad n \geq 1.$$

By convention,  $\det \mathbf{T}_{-1} = 1$ . We can easily verify that to get (3.10) is sufficient to establish that  $|\alpha_n|^2 - 1 > 0$ ,  $n \geq 1$ , and with this, *ii*).

To verify *iii*), let notice that by setting  $B_{2n}(z) = (-1)^n \Phi_n^*(z)$ ,  $B_{2n+1}(z) = \Phi_n(z)$ ,  $n \geq 0$ , we obtain from *i*),

$$\begin{aligned} B_{2n}(z) &= \bar{\alpha}_n z B_{2n-1}(z) - B_{2n-2}(z), \quad n \geq 1 \\ B_{2n+1}(z) &= \alpha_n B_{2n}(z) - (|\alpha_n|^2 - 1) z B_{2n-1}(z), \quad n \geq 1, \end{aligned}$$

with  $B_0 = B_1 = 1$ . Hence, from the theory of continued fractions the result follows.  $\square$

Since  $\alpha_n = (-1)^n \Phi_n(0)$ , from (2.27) we obtain, such as in (3.11),

$$\alpha_n = \frac{\det \mathbf{H}_n^{(-n+2)}}{\det \mathbf{H}_n^{(-n+1)}} = \frac{\overline{\det \mathbf{H}_n^{(-n)}}}{\det \mathbf{H}_n^{(-n+1)}}, \quad n \geq 1. \quad (3.16)$$

Since  $\alpha_n \neq 0$ , from *i*) of the above theorem, the following recurrence relation always holds,

$$\Phi_{n+1}(z) = (z - \beta_{n+1}) \Phi_n(z) - \varsigma_{n+1} z \Phi_{n-1}(z), \quad n \geq 1, \quad (3.17)$$

where  $\Phi_0 = 1$ ,  $\Phi_1(z) = z - \beta_1$ ,  $\beta_1 = \alpha_1$ ,  $\varsigma_{n+1} = \beta_{n+1} (|\alpha_n|^2 - 1)$  and  $\beta_{n+1} = \frac{\alpha_{n+1}}{\alpha_n}$ ,  $n \geq 1$ .

#### 3.1.3 Polynomials of second kind

Let  $\{\Omega_n\}_{n \geq 0}$  be the polynomials of the second kind of  $\{\Phi_n\}_{n \geq 0}$ . It is easily verified that  $\Omega_n$  is a polynomial of exact degree  $n$ . More precisely,

$$\Omega_n(z) = -c_0 (z^n + \dots + (-1)^{n-1} \alpha_n).$$

**Theorem 3.1.2.** *Let  $\{\Phi_n\}_{n \geq 0}$  be the Szegő polynomials associated with the sq-definite moment functional  $\mathcal{L}$ . Then, the polynomials of second kind  $\Omega_n$  satisfy:*

$$i) \quad \Omega_n(z) = \left\langle \mathcal{L}, \frac{z+y}{z-y} \left( \frac{z^k}{y^k} \Phi_n(y) - \Phi_n(z) \right) \right\rangle, \quad 0 \leq k \leq n-1, \quad n \geq 1.$$

$$ii) \quad -\Omega_n^*(z) = \left\langle \mathcal{L}, \frac{z+y}{z-y} \left( \frac{z^k}{y^k} \Phi_n^*(y) - \Phi_n^*(z) \right) \right\rangle, \quad 1 \leq k \leq n, \quad n \geq 1.$$

---


$$\begin{aligned} \text{iii) } -\Omega_n^*(z) &= (-1)^n \bar{\alpha}_n z \Omega_{n-1}(z) - \Omega_{n-1}^*(z), \\ \Omega_n(z) &= (-1)^{n+1} \alpha_n \Omega_n^*(z) - (|\alpha_n|^2 - 1) z \Omega_{n-1}(z), \quad n \geq 1. \end{aligned}$$

iv) For  $n \geq 0$ , if we set  $A_{2n}(z) = (-1)^{n+1} \Omega_n^*(z)$ ,  $A_{2n+1}(z) = \Omega_n(z)$ , then  $\{A_n\}_{n \geq 0}$  is the sequence of numerator polynomials of (3.1), with  $\beta_0 = c_0$  and  $\alpha_n = (-1)^n \Phi_n(0)$ ,  $n \geq 1$ .

*Proof.* The proof of the results in this theorem is the same as in [99].  $\square$

From the above theorem, it also follows that the polynomials  $\{\Omega_n\}_{n \geq 0}$  satisfy the three term recurrence relation

$$\Omega_{n+1}(z) = (z - \beta_{n+1}) \Omega_n(z) - \zeta_{n+1} z \Omega_{n-1}(z), \quad n \geq 1, \quad (3.18)$$

with  $R_0 = -c_0$  and  $\Omega_1(z) = -c_0(z + \alpha_1)$ . The values of  $\zeta_n$  and  $\beta_n$  are as in (3.17).

### 3.1.4 Para-orthogonal polynomials

For  $\omega \in \mathbb{C}$ , we consider the sequence of polynomials  $\{\Phi_n(\omega; \cdot)\}_{n \geq 0}$  and  $\{\Omega_n(\omega; \cdot)\}_{n \geq 0}$  given by  $\Phi_0(\omega; \cdot) = 1$ ,  $\Omega_0(\omega; \cdot) = -c_0$ , and

$$\Phi_n(\omega; z) = \Phi_n(z) - \omega z \Phi_{n-1}(z), \quad \Omega_n(\omega; z) = \Omega_n(z) - \omega z \Omega_{n-1}(z), \quad n \geq 1.$$

Obviously,  $\Phi_n(0; z) = \Phi_n(z)$  and for any  $\omega$ ,

$$\Phi_n(\omega; 0) = \Phi_n(0) = (-1)^n \alpha_n \neq 0, \quad n \geq 1.$$

**Theorem 3.1.3.** Let  $\alpha_n = (-1)^n \Phi_n(0)$ ,  $n \geq 1$ , where  $\{\Phi_n\}_{n \geq 0}$  is the sequence of Szegő polynomials associated with the sq-definite moment functional  $\mathcal{L}$ . Then, the following hold:

- i)  $\langle \Phi_n(\omega; z), z^m \rangle_{\mathcal{L}} = 0, \quad 1 \leq m \leq n-1.$
- ii)  $\Omega_n(\omega; z) = \left\langle \mathcal{L}, \frac{z+y}{z-y} \left( \frac{z^k}{y^k} \Phi_n(\omega; y) - \Phi_n(\omega; z) \right) \right\rangle, \quad 1 \leq k \leq n-1, \quad n \geq 2.$
- iii)  $-\Omega_n^*(\omega; z) = \left\langle \mathcal{L}, \frac{z+y}{z-y} \left( \frac{z^k}{y^k} \Phi_n^*(\omega; y) - \Phi_n^*(\omega; z) \right) \right\rangle, \quad 1 \leq k \leq n-1, \quad n \geq 2.$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

*Proof.* The proof of *i*) is immediate. To prove *ii*), we can write using *i*) of Theorem 3.1.2,

$$\begin{aligned}\Omega_n(\omega; z) &= \Omega_n(z) - \omega z \Omega_{n-1}(z), \quad n \geq 2, \\ &= \left\langle \mathcal{L}, \frac{z+y}{z-y} \left( \frac{z^k}{y^k} \Phi_n(y) - \Phi_n(z) \right) \right\rangle \\ &\quad - \omega z \left\langle \mathcal{L}, \frac{z+y}{z-y} \left( \frac{z^{k-1}}{y^{k-1}} \Phi_{n-1}(y) - \Phi_{n-1}(z) \right) \right\rangle, \quad 1 \leq k \leq n-1.\end{aligned}$$

This leads us to the required result. Similarly, *iii*) follows from *ii*) of Theorem 3.1.2.  $\square$

For convenience, we write  $\widehat{\Phi}_n(\omega; z) = \overline{\Phi}_n(z) - \overline{\omega z} \overline{\Phi}_{n-1}(z)$  and  $\widehat{R}_n(\omega; z) = \overline{R}_n(z) - \overline{\omega z} \overline{R}_{n-1}(z)$ ,  $n \geq 1$ . Hence, observe that

$$\Phi_n^*(\omega; z) = z^n \overline{\widehat{\Phi}_n(\omega; 1/\bar{z})} = z^n \widehat{S}_n(\omega; 1/z) = \Phi_n^*(z) - \overline{\omega} \Phi_{n-1}^*(z), \quad n \geq 1.$$

From the recurrence relations for  $\Phi_n$  in Theorem 3.1.1,

$$\Phi_n(1; z) = (-1)^n \alpha_n \Phi_{n-1}^*(0; z), \quad \Phi_n(-|\alpha_n|^2 + 1; z) = (-1)^n \alpha_n \Phi_n^*(0; z), \quad n \geq 1. \quad (3.19)$$

**Theorem 3.1.4.** *Let  $\alpha_n = (-1)^n \Phi_n(0)$ ,  $n \geq 1$ , where  $\{\Phi_n\}_{n \geq 0}$  are the Szegő polynomials associated with the  $sq$ -definite moment functional  $\mathcal{L}$ . If  $\bar{v} = \frac{1 - \omega - |\alpha_n|^2}{1 - \omega}$ , then*

$$\begin{aligned}i) \quad \Phi_n^*(v; z) &= z^n \widehat{\Phi}_n(v; 1/z) = \frac{1 - \bar{v}}{(-1)^n \alpha_n} \Phi_n(\omega; z), \quad n \geq 1; \\ ii) \quad \Omega_n^*(v; z) &= z^n \widehat{R}_n(v; 1/z) = \frac{1 - \bar{v}}{(-1)^{n-1} \alpha_n} \Omega_n(\omega; z), \quad n \geq 1; \\ iii) \quad \frac{d\Phi_n^*(v; z)}{dz} &= n z^{n-1} \widehat{\Phi}_n(v; 1/z) - z^{n-2} \widehat{\Phi}'_n(v; 1/z) = \frac{1 - \bar{v}}{(-1)^n \alpha_n} \Phi'_n(\omega; z), \quad n \geq 1.\end{aligned}$$

*In particular, if  $\omega_n = 1 - \xi \bar{\alpha}_n$ , with  $|\xi| = 1$ , then*

$$\begin{aligned}iv) \quad \Phi_n^*(\omega_n; z) &= z^n \widehat{\Phi}_n(\omega_n; 1/z) = (-1)^n \bar{\xi} \Phi_n(\omega_n; z), \quad n \geq 1; \\ v) \quad \Omega_n^*(\omega_n; z) &= z^n \widehat{R}_n(\omega_n; 1/z) = (-1)^{n-1} \bar{\xi} \Omega_n(\omega_n; z), \quad n \geq 1; \\ vi) \quad n z^{n-1} \widehat{\Phi}_n(\omega_n; 1/z) &- z^{n-2} \widehat{\Phi}'_n(\omega_n; 1/z) = (-1)^n \bar{\xi} \Phi'_n(\omega_n; z), \quad n \geq 1.\end{aligned}$$

Here,  $\widehat{\Phi}'_n(\omega; z) = \frac{d\widehat{\Phi}_n(\omega; z)}{dz}$  and  $\Phi'_n(\omega; z) = \frac{d\Phi_n(\omega; z)}{dz}$ .

---

*Proof.* Since  $\omega$  and  $\nu$  are different from 1, the proof of *i*) follows from the relations in (3.19). *ii*), especially when  $n \geq 2$ , follows from Theorem 3.1.3. *iii*) is obtained from *i*) by taking derivatives. To obtain the remaining parts, we let  $\omega = \nu = \omega_n$  and this gives  $|\omega_n - 1|^2 = |\alpha_n|^2$ .  $\square$

Notice that we can also write

$$\Phi_n(\omega_n; z) = \Phi_n(1 - \xi \bar{\alpha}_n; z) = \frac{|\alpha_n|^2 - \xi \bar{\alpha}_n}{|\alpha_n|^2 - 1} (\Phi_n(z) + \tau \Phi_n^*(z)), \quad (3.20)$$

where  $\tau = (-1)^n \frac{\alpha_n}{\bar{\alpha}_n} \frac{1 - \xi \bar{\alpha}_n}{\xi - \alpha_n}$ . Again, following the notation used in the case of the classical Szegő polynomials [99], we refer to the polynomials  $\Phi_n(z) + \tau \Phi_n^*(z)$ , when  $|\tau| = 1$ , as the para-orthogonal polynomial of degree  $n$  associated with  $\{\Phi_n\}_{n \geq 0}$ . Since  $|\tau| = 1$ , whenever  $|\xi| = 1$ , the polynomials  $\Phi_n(\omega_n; \cdot)$  are hence para-orthogonal polynomials multiplied by constant factors.

From *i*) of Theorem 3.1.3, we have for  $n \geq 1$ ,

$$F(z) - \frac{\Omega_n(\omega; z)}{\Phi_n(\omega; z)} = O(z^n), \quad F_\infty(z) - \frac{\Omega_n(\omega; z)}{\Phi_n(\omega; z)} = \begin{cases} O\left(\frac{1}{z^{n+1}}\right), & \omega = 0, \\ O\left(\frac{1}{z^{n-1}}\right), & \omega = 1, \\ O\left(\frac{1}{z^n}\right), & \omega \neq \{0, 1\}. \end{cases} \quad (3.21)$$

Let consider the sequence of polynomials  $\{\chi_n^{(1)}(\omega; \cdot)\}_{n \geq 1}$  and  $\{\chi_n^{(2)}(\omega; \cdot)\}_{n \geq 1}$ ,

$$\begin{aligned} \chi_n^{(1)}(\omega; z) &= \Omega_n(\omega; z) \Phi_{n-1}(z) - \Omega_{n-1}(z) \Phi_n(\omega; z), \quad n \geq 1, \\ \chi_n^{(2)}(\omega; z) &= \Phi'_n(\omega; z) \Phi_{n-1}(z) - \Phi'_{n-1}(z) \Phi_n(\omega; z), \quad n \geq 1. \end{aligned}$$

From the recurrence relations (3.17) and (3.18),

$$\begin{aligned} \chi_n^{(1)}(\omega; z) &= \Omega_n(z) \Phi_{n-1}(z) - \Omega_{n-1}(z) \Phi_n(z), \\ &= -\varsigma_n \varsigma_{n-1} z^2 (\Omega_{n-2}(z) \Phi_{n-3}(z) + \Omega_{n-3}(z) \Phi_{n-2}(z)) = \dots \\ &= \varsigma_n \varsigma_{n-1} \dots \varsigma_3 \varsigma_2 z^{n-1} (-\Omega_0(z) \Phi_1(z) + \Omega_1(z) \Phi_0(z)) \\ &= -2c_0 \alpha_1 \varsigma_2 \dots \varsigma_n z^{n-1}, \quad n \geq 2, \end{aligned} \quad (3.22)$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

$$\begin{aligned}
 \chi_n^{(2)}(\omega; z) &= \alpha_n z \left( \Phi_{n-2}(z) \Phi'_{n-1}(z) - \Phi'_{n-2}(z) \Phi_{n-1}(z) \right) - \varsigma_n \Phi_{n-2}(z) \Phi_{n-1}(z) + (1-\omega) \Phi_{n-1}^2(z) \\
 &= \varsigma_n \varsigma_{n-1} z^2 \left( \Phi'_{n-2}(z) \Phi_{n-3}(z) - \Phi'_{n-3}(z) \Phi_{n-2}(z) \right) - \varsigma_n \Phi_{n-2}(z) \Phi_{n-1}(z) + \\
 &+ (1-\omega) \Phi_{n-1}^2(z) + \varsigma_n z \Phi_{n-2}^2(z) - \varsigma_n \varsigma_{n-1} z \Phi_{n-2}(z) \Phi_{n-3}(z) \\
 &= \varsigma_n \varsigma_{n-1} z^2 \chi_{n-2}^{(2)}(0; z) + (1-\omega) \Phi_{n-1}^2(z) + \alpha_n \beta_{n-1} \Phi_{n-2}^2(z), \quad n \geq 2, \quad (3.23)
 \end{aligned}$$

with  $\chi_1^{(1)}(\omega; z) = -2c_0\alpha_1$ ,  $\alpha_1\chi_0^{(2)}(\omega; z) = 0$ , and  $\chi_1^{(2)}(\omega; z) = 1 - \omega$ .

The proof of the following result follows to the same method of the proof given in [52].

**Theorem 3.1.5.** *Let  $\alpha_n = (-1)^n \Phi_n(0)$ ,  $n \geq 1$ , where  $\{\Phi_n\}_{n \geq 0}$  are the Szegő polynomials associated with the sq-definite moment functional  $\mathcal{L}$ . Then, for  $\omega \neq 1$ , the zeros of  $\Phi_n(\omega; \cdot)$  are the eigenvalues of the following lower Hessenberg matrix,*

$$\begin{bmatrix}
 \eta_1 & \varsigma_2 & 0 & \cdots & 0 & 0 \\
 \eta_1 & \eta_2 & \varsigma_3 & \cdots & 0 & 0 \\
 \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\
 \eta_1 & \eta_2 & \eta_3 & \cdots & \varsigma_{n-1} & 0 \\
 \eta_1 & \eta_2 & \eta_3 & \cdots & \eta_{n-1} & \frac{\varsigma_n}{(1-\omega)} \\
 \eta_1 & \eta_2 & \eta_3 & \cdots & \eta_{n-1} & \frac{\eta_n}{(1-\omega)}
 \end{bmatrix},$$

where  $\eta_1 = \alpha_1$ ,  $\eta_r = \alpha_r \bar{\alpha}_{r-1}$ , and  $\varsigma_r = \alpha_r \bar{\alpha}_{r-1} - \alpha_r / \alpha_{r-1}$ ,  $r = 2, \dots, n$ .

*Proof.* From (3.17) and the definition of  $\{\Phi_n(\omega, \cdot)\}_{n \geq 0}$ , we have

$$\Phi_n(\omega; z) = \begin{vmatrix}
 z - \beta_1 & -\varsigma_2 & 0 & \cdots & 0 & 0 \\
 -z & z - \beta_2 & -\varsigma_3 & \cdots & 0 & 0 \\
 0 & -z & z - \beta_3 & & 0 & 0 \\
 \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\
 0 & 0 & 0 & \cdots & z - \beta_{n-1} & -\varsigma_n \\
 0 & 0 & 0 & \cdots & -z & (1-\omega)z - \beta_n
 \end{vmatrix},$$

or, equivalently,  $\Phi_n(\omega; z) = \det(z\mathbf{A}_n - \mathbf{B}_n)$ , where  $\mathbf{A}_n$  and  $\mathbf{B}_n$  are matrices of order  $n$



given by

$$\mathbf{A}_n = \begin{bmatrix} 1 & 0 & 0 & \cdots & 0 \\ -1 & 1 & 0 & \cdots & 0 \\ 0 & -1 & 1 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & 1-\omega \end{bmatrix}, \quad \mathbf{B}_n = \begin{bmatrix} \beta_1 & \varsigma_2 & 0 & \cdots & 0 \\ 0 & \beta_2 & \varsigma_3 & \cdots & 0 \\ 0 & 0 & \beta_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \beta_n \end{bmatrix}.$$

Since  $\mathbf{A}_n$  is non-singular,  $\Phi_n(\omega; z) = \det \mathbf{A}_n \det (z\mathbf{I}_n - \mathbf{A}_n^{-1}\mathbf{B}_n) = (1-\omega) \det \mathbf{G}_n$ , where

$$\mathbf{G}_n = \begin{bmatrix} z-\beta_1 & \varsigma_2 & 0 & \cdots & 0 & 0 \\ \beta_1 & z-(\varsigma_2+\beta_2) & \alpha_3 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ \beta_1 & \varsigma_2+\beta_2 & \varsigma_3+\beta_3 & \cdots & z-(\varsigma_{n-1}+\beta_{n-1}) & \varsigma_n \\ \frac{\beta_1}{1-\omega} & \frac{\varsigma_2+\beta_2}{1-\omega} & \frac{\varsigma_3+\beta_3}{1-\omega} & \cdots & \frac{\varsigma_{n-1}+\beta_{n-1}}{1-\omega} & \frac{z(1-\omega)-(\varsigma_n+\beta_n)}{1-\omega} \end{bmatrix}.$$

Thus, the result follows.  $\square$

## 3.2 Polynomials with real zeros

Let  $\mathcal{L}$  be an rsq-definite moment functional. That is, all the moments  $\{c_n\}_{n \in \mathbb{Z}}$  are real and the properties (3.8) and (3.10) hold. Clearly, the associated Szegő polynomials  $\{\Phi_n\}_{n \geq 0}$  are all real and, in particular, we can state the following theorem.

**Theorem 3.2.1.** *Let  $\alpha_n = (-1)^n \Phi_n(0)$ ,  $n \geq 1$ , where  $\{\Phi_n\}_{n \geq 0}$  are the Szegő polynomials associated with the rsq-definite moment functional  $\mathcal{L}$ . Then, the following statements are equivalent.*

i)  $\det \mathbf{H}_n^{(-n)} > 0$ ,  $n \geq 1$ .

ii)  $\alpha_n > 1$ ,  $n \geq 1$ .

iii) *The zeros of the polynomials  $\Phi_n(\omega; \cdot)$ ,  $n \geq 1$ , when  $\omega < 1$ , are simple and lie on the positive half side of the real line.*

*Proof.* If  $\mathcal{L}$  is an rsq-definite moment functional then their moments are real and  $\det \mathbf{H}_n^{(-n+1)} > 0$ ,  $n \geq 1$ . Hence, from (3.16),  $\det \mathbf{H}_n^{(-n)} > 0$ ,  $n \geq 1$ , and  $\alpha_n > 1$ ,  $n \geq 1$ .

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

On the other hand, if  $\alpha_n > 1$ ,  $n \geq 1$ , then  $\det \mathbf{H}_n^{(-n)} > 0$ ,  $n \geq 1$ , follows from (3.11). Therefore *i*) and *ii*) are equivalent.

We now prove the equivalence of *ii*) and *iii*). Suppose that all zeros of  $\Phi_n(\omega; \cdot)$ ,  $n \geq 1$ , are positive. Since the leading coefficient  $1 - \omega$  of  $\Phi_n(\omega; \cdot)$  is positive,  $\alpha_n = (-1)^n \Phi_n(\omega; 0) > 0$ ,  $n \geq 1$ . Combining this with the fact that  $\mathcal{L}$  is an rsq-definite moment functional, *ii*) follows. On the other hand, if  $\alpha_n > 1$ ,  $n \geq 1$ , then, from (3.23) and by induction, we can prove that  $\chi_n^{(2)}(\omega; z) > 0$ ,  $n \geq 1$ , for  $\omega < 1$  and for any real values of  $z$ . Using this with the additional conditions  $(-1)^n \Phi_n(0) > 0$ ,  $n \geq 1$ , we easily establish that the zeros of  $\Phi_n(\omega; \cdot)$  are positive and simple (and, even more, interlace with the zeros of  $\Phi_{n-1}$ ). This proves the equivalence of *ii*) and *iii*). Consequently, our theorem is proved.  $\square$

The recurrence relation (3.17), which corresponds to the conditions of the above theorem, has been studied, for instance, in [161].

**Remark 3.2.1.** *One can also establish in an analog way (or using the polynomials  $\Psi_n(z) = (-1)^n \Phi_n(-z)$ ,  $n \geq 1$ ) the equivalence of the statements.*

$$i) \quad (-1)^n \det \mathbf{H}_n^{(-n)} > 0 \quad n \geq 1.$$

$$ii) \quad (-1)^n \alpha_n > 1 \quad n \geq 1.$$

*iii) The zeros of the polynomials  $\Phi_n(\omega; \cdot)$ ,  $n \geq 1$ , when  $\omega < 1$ , are simple and lie on the negative half side of the real line.*

In fact, when  $(-1)^n \alpha_n > 1$ ,  $n \geq 1$ , the polynomials  $\{\Phi_n\}_{n \geq 0}$  are the denominator polynomials of a positive continued fraction,

$$\sqrt{\frac{\lambda_1 z}{1 + \eta_1 z}} + \sqrt{\frac{\lambda_2 z}{1 + \eta_2 z}} + \sqrt{\frac{\lambda_3 z}{1 + \eta_3 z}} + \cdots, \quad \lambda_n > 0, \quad \eta_n > 0, \quad n \geq 0.$$

This kind of continued fractions are known as  $\mathcal{T}$ -fractions. We can thus also use some known results on the denominator polynomials of a positive  $\mathcal{T}$ -fraction given in [101]. From now on we restrict our analysis to the case when  $\alpha_n > 1$ ,  $n \geq 1$ , and we refer to the associated moment functional as an rsq-definite moment functional on  $(0, \infty)$ .

The results given from here can be easily extended to the case when  $(-1)^n \alpha_n > 1$ ,  $n \geq 1$ , that is, when the moment functional is an rsq-definite moment functional on  $(-\infty, 0)$ .

---

**Corollary 3.2.1.** *Let  $\{\Phi_n\}_{n \geq 0}$  be the sequence of Szegő polynomials associated with an rsq-definite moment functional  $\mathcal{L}$  on  $(0, \infty)$ . Then, for the zeros  $z_{n,r}(\omega)$  of  $\Phi_n(\omega; \cdot)$ , we have*

$$i) \quad 0 < z_{n,1}(\omega) < z_{n,2}(\omega) < \dots < z_{n,n-1}(\omega) < z_{n,n}(\omega), \quad \omega < 1.$$

$$ii) \quad z_{n,1}(\omega) < 0 < z_{n,2}(\omega) < \dots < z_{n,n-1}(\omega) < z_{n,n}(\omega), \quad \omega > 1.$$

*Proof.* All we need to verify is that when  $\omega > 1$ , the zeros of  $\Phi_n(\omega; \cdot)$  are still real and simple, and that one of the zeros is of the opposite sign. We verify this using the monic polynomial  $(1 - \omega)^{-1} \Phi_n(\omega; \cdot)$ . Substituting  $z = 0$  and  $z = z_{n,r}(0)$ ,  $r = 1, 2, \dots, n$ , in the definition of  $\Phi_n(\omega; \cdot)$ ,

$$\prod_{r=1}^n z_{n,r}(\omega) = (1 - \omega)^{-1} \prod_{r=1}^n z_{n,r}(0)$$

and

$$(1 - \omega)^{-1} \Phi_n(\omega; z_{n,r}(0)) = \frac{\omega}{\omega - 1} z_{n,r}(0) \Phi_{n-1}(z_{n,r}(0)), \quad r = 1, 2, \dots, n.$$

Since the zeros of  $\Phi_n(0; \cdot) = \Phi_n$  are positive, simple and interlace with the zeros of  $\Phi_{n-1}$ , we can conclude that the zeros of  $\Phi_n(\omega; \cdot)$  interlace with the zeros of  $\Phi_n$ , proving that they are real and simple. Moreover, at the most one zero of  $\Phi_n(\omega; \cdot)$  can have negative sign, which certainly happens if  $(1 - \omega)$  is negative.  $\square$

We can now state the following results on the quadrature rules associated with the polynomials  $\{\Phi_n(\omega; \cdot)\}_{n \geq 0}$ .

**Theorem 3.2.2.** *Let  $\omega < 1$  and let  $\{\Phi_n\}_{n \geq 0}$  be the Szegő polynomials associated with an rsq-definite moment functional  $\mathcal{L}$  on  $(0, \infty)$ . Then, the following statements hold.*

$$i) \quad \frac{\Omega_n(\omega; z)}{\Phi_n(\omega; z)} = \sum_{r=1}^n \frac{z_{n,r}(\omega) + z}{z_{n,r}(\omega) - z} \lambda_{n,r}(\omega), \quad \lambda_{n,r}(\omega) = \frac{\Omega_n(\omega; z_{n,r}(\omega))}{-2z_{n,r}(\omega) \Phi_n'(\omega; z_{n,r}(\omega))}, \quad n \geq 1.$$

$$ii) \quad \lambda_{n,r}(\omega) > 0, \quad \sum_{r=1}^n \lambda_{n,r}(\omega) = c_0, \quad 1 \leq r \leq n.$$

iii) *For any  $n \geq 1$ , if  $l_{k,s}(z) = \sum_{j=k}^s a_j z^j$  is a Laurent polynomial such that  $-n + 1 \leq k \leq s \leq n - 1$ , then the quadrature rule holds*

$$\langle \mathcal{L}, l_{k,s} \rangle = \sum_{r=1}^n \lambda_{n,r}(\omega) l_{k,s}(z_{n,r}(\omega)).$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

*Proof.* To prove *i*), first observe that the zeros of  $\Phi_n(\omega; \cdot)$  are simple and lie within  $(0, \infty)$ . Hence, a partial fraction decomposition of the given type verifies *i*).

Notice that  $\lambda_{n,r}(\omega)$  can be written as

$$\begin{aligned} \lambda_{n,r}(\omega) &= \frac{-\Omega_n(\omega; z_{n,r}(\omega))\Phi_{n-1}(z)/z_{n,r}(\omega) + \Omega_{n-1}(z_{n,r}(\omega))\Phi_n(\omega; z_{n,r}(\omega))}{2\Phi'_n(\omega; z_{n,r}(\omega))\Phi_{n-1}(z) - \Phi'_{n-1}(z)\Phi_n(\omega; z_{n,r}(\omega))} \\ &= \frac{-\chi_n^{(1)}(\omega; z_{n,r}(\omega))/z_{n,r}(\omega)}{2\chi_n^{(2)}(\omega; z_{n,r}(\omega))}, \\ &= \frac{c_0(|\alpha_1|^2 - 1)(|\alpha_2|^2 - 1) \dots (|\alpha_{n-1}|^2 - 1)\alpha_n z_{n-r}^{n-2}(\omega)}{\chi_n^{(2)}(\omega; z_{n,r}(\omega))}, \end{aligned}$$

where  $\chi_n^{(1)}(\omega; \cdot)$  and  $\chi_n^{(2)}(\omega; \cdot)$  are as in (3.22) and (3.23). Since  $\alpha_n > 1$  for  $n \geq 1$ , one can easily verify that the numerator and denominator above are positive and, hence *ii*) is established.

When  $n = 1$ , *iii*) clearly follows from *ii*). To obtain *iii*) when  $n \geq 2$ , we note that  $\tilde{l}_{k,s}(y) = y^{n-1}l_{k,s}(y) \in \mathbb{P}_{2n-2}$ . Hence, the interpolation on the zeros of  $\Phi_n(\omega; \cdot)$  gives

$$\begin{aligned} \tilde{l}_{k,s}(y) &= \sum_{r=1}^n \frac{\Phi_n(\omega; y)}{(y - z_{n,r}(\omega))\Phi'_n(\omega; z_{n,r}(\omega))} \tilde{l}_{k,s}(z_{n,r}(\omega)) \\ &\quad + \tilde{l}_{k,s}(z_{n,1}(\omega), \dots, z_{n,n}(\omega); y)\Phi_n(\omega; y). \end{aligned}$$

Here, we use the divided difference  $\tilde{l}_{k,s}(z_{n,1}(\omega), \dots, z_{n,n}(\omega); y) \in \mathbb{P}_{n-2}$ , and therefore,

$$\left\langle \tilde{l}_{k,s}(z_{n,1}(\omega), \dots, z_{n,n}(\omega); y)\Phi_n(\omega; y), y^{n-1} \right\rangle_{\mathcal{L}} = 0.$$

This means,

$$\langle \mathcal{L}, l_{k,s} \rangle = \left\langle \tilde{l}_{k,s}, y^{-n+1} \right\rangle_{\mathcal{L}} = \sum_{r=1}^n \tilde{\lambda}_{n,r}(\omega)(z_{n,r}(\omega))^{n-1} l_{k,s}(z_{n,r}(\omega)),$$

where

$$\tilde{\lambda}_{n,r}(\omega) = \left\langle \frac{\Phi_n(\omega; y)}{(y - z_{n,r}(\omega))\Phi'_n(\omega; z_{n,r}(\omega))}, y^{n-1} \right\rangle_{\mathcal{L}}.$$

Taking into account

$$\Omega_n(\omega; z) + 2z^n \left\langle \frac{\Phi_n(\omega; y)}{y - z}, y^{-n+1} \right\rangle_{\mathcal{L}} = \Phi_n(\omega; z) \left\langle \mathcal{L}, \frac{y+z}{y-z} \right\rangle,$$

we get the statement of our theorem.  $\square$

Moreover, using the results given in Theorem 3.1.4 and Theorem 3.2.2 the following results can also be easily verified. If  $\omega < 1$  and  $\nu = \frac{1 - \omega - |\alpha_n|^2}{1 - \omega}$ , then for  $n \geq 1$ ,

$$z_{n,r}(\omega) = 1/z_{n,n-r+1}(\nu), \quad \lambda_{n,r}(\omega) = \lambda_{n,n-r+1}(\nu), \quad r = 1, 2, \dots, n. \quad (3.24)$$

The results in Theorem 3.2.2 allow one to define the step function  $\mu_n(\omega; \cdot)$  by

$$\mu_n(\omega; y) = \begin{cases} 0, & 0 < y \leq z_{n,1}(\omega), \\ \sum_{r=1}^s \lambda_{n,r}(\omega), & z_{n,s}(\omega) < y \leq z_{n,s+1}(\omega), \\ c_0, & z_{n,n}(\omega) < y < \infty. \end{cases}$$

Then, from *i*) of Theorem 3.2.2 we get

$$\frac{\Omega_n(\omega; z)}{\Phi_n(\omega; z)} = \int_0^\infty \frac{y+z}{y-z} d\mu_n(\omega; y), \quad n \geq 1$$

and hence, from (3.21), for  $n \geq 1$ ,

$$c_s = \int_0^\infty y^s d\mu_n(\omega; y), \quad -n+1 \leq s \leq n-1.$$

Helly's selection principle [46] states that a sequence of functions which is locally of bounded total variation has a convergent subsequence. Hence, using the Helly selection theorem, there exists a subsequence  $\{n_k\}$  such that  $\{\mu_{n_k}(\omega)\}$  converges to a bounded non-decreasing function  $\mu$  defined on the positive half side of the real line. The function  $\mu$  is such that it has infinitely many points of increase in  $(0, \infty)$  and

$$\lim_{k \rightarrow \infty} \frac{\Omega_{n_k}(\omega; z)}{\Phi_{n_k}(\omega; z)} = \lim_{k \rightarrow \infty} \int_0^\infty \frac{y+z}{y-z} d\mu_{n_k}(\omega; t) = \int_0^\infty \frac{y+z}{y-z} d\mu(t), \quad 0 < \arg(z) < 2\pi.$$

The convergence is also uniform for compact subsets within  $0 < \arg(z) < 2\pi$ . The points of increase (support) of  $\mu$  lie entirely on the positive half side of the real line. Moreover,

$$c_n = \int_0^\infty y^n d\mu(y), \quad n \in \mathbb{Z}, \quad (3.25)$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

and

$$\int_0^\infty y^{-m} \Phi_n(y) d\mu(y) = \int_0^\infty y^{-n+m} \Phi_n^*(y) d\mu(y) = \frac{\det \mathbf{T}_n}{\det \mathbf{T}_{n-1}} \delta_{n,m}, \quad 0 \leq m \leq n, \quad n \geq 1. \quad (3.26)$$

Assuming  $\mathcal{L}$  to be an rsq-definite moment functional on  $(0, \infty)$ , we now consider the sequence of polynomials  $\{\Phi_n(\widehat{\omega}_n; \cdot)\}_{n \geq 0}$ , where  $\widehat{\omega}_n = 1 - \alpha_n$ . The zeros of  $\Phi_n(\widehat{\omega}_n; \cdot)$  are on the positive half side of the real line. Using the results given in Theorem 3.1.4, we can say more about these zeros and also the quadrature weights  $\lambda_{n,r}(\widehat{\omega}_n)$  which follow from Theorem 3.2.2.

$$z_{n,r}(\widehat{\omega}_n) = 1/z_{n,n-r+1}(\widehat{\omega}_n), \quad \lambda_{n,r}(\widehat{\omega}_n) = \lambda_{n,n-r+1}(\widehat{\omega}_n), \quad r = 1, 2, \dots, n, \quad n \geq 1.$$

This means that the distribution given by the step function  $\mu_n(\widehat{\omega}_n; \cdot)$  satisfies

$$d\mu_n(\widehat{\omega}_n; y) = -d\mu_n(\widehat{\omega}_n; 1/y), \quad y \in (0, \infty).$$

Hence, applying the Helly selection theorem we can state the following result.

**Theorem 3.2.3.** *Let  $\alpha_n = (-1)^n \Phi_n(0)$ , where  $\{\Phi_n\}_{n \geq 0}$  are the Szegő polynomials associated with  $\mathcal{L}$ , which is an rsq-definite moment functional on  $(0, \infty)$ . Then there exists a bounded non-decreasing function  $\mu$ , with all its points of increase on  $(0, \infty)$ , such that  $d\mu(y) = -d\mu(1/y)$ . Moreover, (3.25) and (3.26) hold.*

Let  $0 \leq a < b \leq \infty$ . We say that the strong positive measure  $\nu$ , defined on  $(a, b)$ , belongs to the symmetric class  $\mathcal{S}^3[\xi, \beta, b]$  if

$$\frac{d\nu(y)}{y^\xi} = -\frac{d\nu(\beta^2/y)}{(\beta^2/y)^\xi}, \quad y \in [a, b],$$

where  $0 < \beta < b$ ,  $a = \beta^2/b$  and  $2\xi \in \mathbb{Z}$ . The classification of the symmetry is according to the value of  $\xi$  [19; 48; 161]. Notice that our orthogonality measure  $\mu$  is classified as the class  $\mathcal{S}^3(0, 1, b)$ ,  $1 < b \leq \infty$ .

To be able to talk about Szegő polynomials on a positive interval  $[a, b]$ , it is important that the measure belongs to the class  $\mathcal{S}^3(0, 1, b)$ . Otherwise, the monic polynomials  $\{\Phi_n\}_{n \geq 0}$  defined on  $[a, b]$  by (3.26) do not satisfy *i*) of Theorem 3.1.1. For example, if the measure belongs to the class  $\mathcal{S}^3(1/2, 1, b)$ , then for the monic polynomials  $\{\Phi_n\}_{n \geq 0}$

defined by (3.26) we would have, instead of  $i$  in Theorem 3.1.1,

$$\Phi_n(z) = \tilde{\delta}_n \Phi_n(z) = z\Phi_{n-1}(z) + \tilde{\delta}_n \Phi_{n-1}(z) - \tilde{\alpha}_n z \Phi_{n-2}(z), \quad n \geq 2,$$

where  $\tilde{\delta}_n = \Phi_n(0)$  and  $\tilde{\alpha}_n > 0$ .

We now look at some properties of the sequences of polynomials  $\{\Phi_n(\widehat{\omega}_n; \cdot)\}_{n \geq 0}$  and  $\{\Phi_n(\check{\omega}_n; \cdot)\}_{n \geq 0}$ , where  $\widehat{\omega}_n = 1 - \alpha_n$  as above, and  $\check{\omega}_n = 1 + \alpha_n$ , when  $\mathcal{L}$  is an rsq-definite moment functional on  $(0, \infty)$ . From Theorem 3.1.4 and Corollary 3.2.1, the only negative zero of  $\Phi_n(\check{\omega}_n; \cdot)$  is  $-1$ .

In (3.20), letting  $\xi = \alpha_n/|\alpha_n|$  and  $\xi = -\alpha_n/|\alpha_n|$ , we have for the monic polynomials  $\Phi_n(\widehat{\omega}_n; z)/|\alpha_n|$  and  $-\Phi_n(\check{\omega}_n; z)/|\alpha_n|$ ,

$$\frac{\Phi_n(\widehat{\omega}_n; z)}{|\alpha_n|} = \frac{\Phi_n(z) + \tau_n \Phi_n^*(z)}{1 + |\alpha_n|}, \quad \frac{\Phi_n(\check{\omega}_n; z)}{-|\alpha_n|} = \frac{\Phi_n(z) - \tau_n \Phi_n^*(z)}{1 - |\alpha_n|}, \quad n \geq 1,$$

where  $\tau_n = (-1)^n \alpha_n/|\alpha_n|$ . Hence, from  $i$ ) of Theorem 3.1.1, we obtain

$$\begin{aligned} \frac{\Phi_{n+1}(\widehat{\omega}_{n+1}; z)}{|\alpha_{n+1}|} &= \left( z - \frac{|\alpha_n|}{\alpha_n} \frac{\alpha_{n+1}}{|\alpha_{n+1}|} \right) \frac{\Phi_n(\widehat{\omega}_n; z)}{|\alpha_n|} \\ &\quad - \frac{|\alpha_n|}{\alpha_n} \frac{\alpha_{n+1}}{|\alpha_{n+1}|} (|\alpha_n| - 1)(|\alpha_{n-1}| + 1) z \frac{\Phi_{n-1}(z) + \tilde{\tau}_{n-1} \Phi_{n-1}^*(z)}{1 + |\alpha_{n-1}|}, \\ \frac{\Phi_{n+1}(\check{\omega}_{n+1}; z)}{-|\alpha_{n+1}|} &= \left( z - \frac{|\alpha_n|}{\alpha_n} \frac{\alpha_{n+1}}{|\alpha_{n+1}|} \right) \frac{\Phi_n(\check{\omega}_n; z)}{-|\alpha_n|} \\ &\quad - \frac{|\alpha_n|}{\alpha_n} \frac{\alpha_{n+1}}{|\alpha_{n+1}|} (|\alpha_n| + 1)(|\alpha_{n-1}| - 1) z \frac{\Phi_{n-1}(z) - \tilde{\tau}_{n-1} \Phi_{n-1}^*(z)}{1 - |\alpha_{n-1}|}, \end{aligned}$$

where  $\tilde{\tau}_{n-1} = (-1)^{n-1} \frac{\alpha_n^2}{|\alpha_n|^2} \frac{|\alpha_{n+1}|}{\alpha_{n+1}}$ . Consequently, these lead to recurrence relations for  $\frac{\Phi_n(\widehat{\omega}_n; z)}{|\alpha_n|}$  and  $\frac{-\Phi_n(\check{\omega}_n; z)}{|\alpha_n|}$ , since  $\tilde{\tau}_{n-1} = \tau_{n-1}$  when  $\alpha_n > 1$ ,  $n \geq 1$ .

**Theorem 3.2.4.** *Let  $\alpha_n = (-1)^n \Phi_n(0)$ , where  $\{\Phi_n\}_{n \geq 0}$  are the Szegő polynomials associated with  $\mathcal{L}$ , which is an rsq-definite moment functional on  $(0, \infty)$ . Then,*

$$\begin{aligned} \frac{\Phi_{n+1}(\widehat{\omega}_{n+1}; z)}{\alpha_{n+1}} &= (z - 1) \frac{\Phi_n(\widehat{\omega}_n; z)}{\alpha_n} - (\alpha_n - 1)(\alpha_{n-1} + 1) z \frac{\Phi_{n-1}(\widehat{\omega}_{n-1}; z)}{\alpha_{n-1}}, \quad n \geq 1, \\ \frac{\Phi_{n+1}(\check{\omega}_{n+1}; z)}{-\alpha_{n+1}} &= (z - 1) \frac{\Phi_n(\check{\omega}_n; z)}{-\alpha_n} - (\alpha_n + 1)(\alpha_{n-1} - 1) z \frac{\Phi_{n-1}(\check{\omega}_{n-1}; z)}{-\alpha_{n-1}}, \quad n \geq 1, \end{aligned}$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

with  $\frac{\Phi_1(\widehat{\omega}_1; z)}{\alpha_1} = z - 1$ ,  $\frac{\Phi_0(\widehat{\omega}_0; z)}{\alpha_0} = 1$ ,  $\frac{\Phi_1(\check{\omega}_1; z)}{-\alpha_1} = z + 1$ , and  $\frac{\Phi_0(\check{\omega}_0; z)}{-\alpha_0} = 0$ .

Let  $\mathcal{L}$  be an rsq-definite moment functional on  $(0, \infty)$  defined as in (3.8)-(3.10) and satisfying the conditions of Theorem 3.2.1. If  $\Phi_n$  denotes the  $n$ -th Szegő polynomial associated with this moment functional, then the zeros  $z_{n,r}$  of  $\Phi_n$  are positive. We arrange them in an increasing order

$$0 < z_{n,1} < z_{n,2} < \cdots < z_{n,n-1} < z_{n,n}.$$

In what follows we denote by  $z_{n,r}^*$  and  $k_{n,r}$  the zeros of  $\Phi_n^*$  and  $K_n(\alpha, \cdot)$ , respectively. Interesting inequalities between zeros of these polynomials follow from Theorem 3.1.1 and from (2.33). Our results read as follows.

**Theorem 3.2.5.** *The zeros of  $\Phi_n$  and  $\Phi_n^*$  interlace. More precisely,*

$$0 < z_{n,1}^* < z_{n,1} < z_{n-1,1} < \cdots < z_{n,n-1}^* < z_{n,n-1} < z_{n-1,n-1} < z_{n,n}^* < z_{n,n}.$$

*Proof.* By Theorem 3.1.1, we have

$$\Phi_n(z) = (-1)^n \alpha_n \Phi_n^*(z) - (|\alpha_n|^2 - 1)z \Phi_{n-1}(z), \quad n \geq 1.$$

Since the zeros of  $\Phi_n$  and  $\Phi_{n-1}$  interlace, we obtain from the above relation

$$\operatorname{sgn}((-1)^n \Phi_n^*(z_{n,r})) = \operatorname{sgn}(\Phi_{n-1}(z_{n,r}))$$

and

$$\operatorname{sgn}(\Phi_n(z_{n-1,r})) = \operatorname{sgn}((-1)^n \Phi_n^*(z_{n-1,r})).$$

Hence there exist  $z_{n,r}^*$ ,  $r = 1, \dots, n$ , zeros of  $\Phi_n^*$ , such that

$$0 < z_{n,1}^* < z_{n,1} < z_{n-1,1} < \cdots < z_{n,n-1}^* < z_{n,n-1} < z_{n-1,n-1} < z_{n,n}^* < z_{n,n},$$

thus proving the theorem. □

**Theorem 3.2.6.** *If  $\alpha < 0$ , then the zeros of  $\Phi_n$  and  $K_n(\alpha, \cdot)$  satisfy the interlacing property*

$$0 < z_{n,1}^* < z_{n,1} < k_{n-1,1} < \cdots < z_{n,n-1}^* < z_{n,n-1} < k_{n-1,n-1} < z_{n,n}^* < z_{n,n}.$$



---

*Proof.* By (2.33)

$$\mathbf{k}_n(1 - \alpha z)K_{n-1}(z, \alpha) = \Phi_n^*(\alpha)\Phi_n^*(z) - \Phi_n(\alpha)\Phi_n(z).$$

Since  $(-1)^{n(n+1)/2} \det \mathbf{T}_n > 0$  and (2.27), we have  $\text{sgn}(\mathbf{k}_n) = (-1)^n$ . We also have  $\text{sgn}(\Phi_n(\alpha)) = (-1)^n$  and  $\Phi_n^*(\alpha) > 0$ . On the other hand, by Theorem 3.2.5,  $\Phi_n$  and  $\Phi_n^*$  have interlacing zeros. Therefore, from the above relation

$$(-1)^n \text{sgn}(K_{n-1}(z_{n,r}, \alpha)) = \text{sgn}(\Phi_n^*(z_{n,r})), \quad 1 \leq r \leq n,$$

and

$$\text{sgn}(K_{n-1}(z_{n,r}^*, \alpha)) = -\text{sgn}(\Phi_n(z_{n,r}^*)), \quad 1 \leq r \leq n.$$

Hence, there exist zeros  $k_{n-1,r}$ ,  $r = 1, \dots, n-1$ , of  $K_{n-1}(z, \alpha)$ , satisfying  $0 < z_{n,1}^* < z_{n,1} < k_{n-1,1} < \dots < z_{n,n-1}^* < z_{n,n-1} < k_{n-1,n-1} < z_{n,n}^* < z_{n,n}$ .  $\square$

### 3.2.1 Associated moment problem

The necessary and sufficient conditions for the existence of a solution of the strong Stieltjes moment problem introduced in Chapter 1 are the following

$$\det \mathbf{H}_n^{(-n)} > 0, \quad \det \mathbf{H}_n^{(-n+1)} > 0, \quad n \geq 0.$$

From the Hamburger moment problem, these conditions become

$$\det \mathbf{H}_{2n}^{(-2n)} > 0, \quad \det \mathbf{H}_{2n+1}^{(-2n+1)} > 0, \quad n \geq 0.$$

The objective of this section is to solve the moment problem (1.11) formulated in the introduction. We can easily verify that the modified moments associated with the Chebyshev polynomials can be rewritten as follows

$$c_n = \int_0^\infty T_n(\cosh \theta) d\bar{\mu}(\cosh \theta) = -\frac{1}{2} \int_{-\infty}^0 e^{n\theta} d\bar{\mu}(\cosh \theta) + \frac{1}{2} \int_0^\infty e^{n\theta} d\bar{\mu}(\cosh \theta).$$

Substituting  $x = e^\theta$ , we obtain

$$c_n = -\frac{1}{2} \int_0^1 x^n d\bar{\mu}\left(\frac{x+x^{-1}}{2}\right) + \frac{1}{2} \int_1^\infty x^n d\bar{\mu}\left(\frac{x+x^{-1}}{2}\right) = \int_0^\infty x^n d\mu(x),$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

where  $\mu$  is a non-decreasing distribution function supported on  $(0, \infty)$

$$d\mu(x) = \begin{cases} -\frac{1}{2} d\bar{\mu}\left(\frac{x+x^{-1}}{2}\right), & 0 < x \leq 1; \\ \frac{1}{2} d\bar{\mu}\left(\frac{x+x^{-1}}{2}\right), & 1 \leq x < \infty. \end{cases}$$

Hence, our moment problem can be stated as follows. Given a sequence of real numbers  $\{c_n\}_{n \geq 0}$  find necessary and sufficient conditions for the existence of a measure  $\mu$  supported on  $(0, \infty)$ , with the symmetry  $d\mu(x) = -d\mu(1/x)$ , such that,

$$c_n = \int_0^\infty x^n d\mu(x), \quad n \geq 0. \quad (3.27)$$

**Remark 3.2.2.** Consider the following moment problem: Given a sequence  $\{c_n\}_{n \geq 0}$  of real numbers find necessary and sufficient conditions for the existence of a measure  $\bar{\mu}$  supported on the whole real line, with the symmetry  $d\bar{\mu}(\theta) = -d\bar{\mu}(-\theta)$ , such that

$$\mu_n = \int_{-\infty}^\infty e^{n\theta} d\bar{\mu}(\theta), \quad n \geq 0.$$

Notice that with the substitution  $t = e^\theta$  and  $x = \cosh(\theta)$  this moment problem is equivalent to the moment problem (3.27).

**Theorem 3.2.7.** The moment problem associated with the Chebyshev polynomials of the first kind has at least one solution if and only if  $c_n = c_{-n}$ , and

$$\det \mathbf{H}_{n+1}^{(-n)} > 0, \quad \det \mathbf{H}_{n+1}^{(-n-1)} > 0, \quad n \geq 0.$$

*Proof.* The existence of the measure  $\mu$  follows immediately from Theorem 3.2.3. Conversely, let  $\mu$  be a measure supported on  $(0, \infty)$ . Given the following quadratic form

$$J(n, l) = \sum_{i=-n}^n \sum_{j=-n}^n \mu_{i+j+l} v_i v_j, \quad l \geq 0, \quad n \geq 1,$$

it is easily verified that

$$J(n, l) = \sum_{i=-n}^n \sum_{j=-n}^n \left( \int_0^\infty t^{i+j+l} d\mu(x) \right) v_i v_j = \int_0^\infty t^l \left( \sum_{i=-n}^n x^i v_i \right)^2 d\mu(x) > 0.$$

Since  $J(n, l)$  is positive definite, we can establish that

$$\det \mathbf{H}_{k+1}^{(-2n+l)} = \begin{vmatrix} c_{-2n+l} & c_{-2n+1+l} & \cdots & c_{-2n+k+l} \\ c_{-2n+1+l} & c_{-2n+2+l} & \cdots & c_{-2n+k+1+l} \\ \vdots & \vdots & \ddots & \vdots \\ c_{-2n+k+l} & c_{-2n+k+1+l} & \cdots & c_{-2n+2k+l} \end{vmatrix} > 0, \quad k = 0, 1, 2, \dots, 2n.$$

Thus, for  $l = 0$  ( $l = 1$ ) with  $k = 2n$  and  $k = 2n - 1$  ( $k = 2n - 1$  and  $k = 2n - 2$ ), the result follows.  $\square$

**Theorem 3.2.8.** *Let  $\alpha_n = (-1)^n \Phi_n(0)$ , where  $\{\Phi_n\}_{n \geq 0}$  are the Szegő polynomials orthogonal with respect to the symmetric measure  $\mu$ . The moment problem associated with the Chebyshev polynomials of the first kind is determinate if and only if*

$$\sum_{n=1}^{\infty} e_n = \infty, \quad \text{or} \quad \sum_{n=1}^{\infty} d_n = \infty, \quad (3.28)$$

where  $e_1 = \frac{1}{c_0 \alpha_1}$ ,  $d_1 = \frac{1}{\mu_0}$ ,

$$\begin{aligned} d_{2n} &= \frac{1}{\alpha_{2n}} \prod_{r=2}^n \left( \left( \frac{\alpha_{2r-1}^2}{\alpha_{2r-1}^2 - 1} \right) \left( \frac{\alpha_{2r-2}^2 - 1}{\alpha_{2r-2}^2} \right) \right) \frac{\alpha_1^2}{\alpha_1^2 - 1}, \\ e_{2n} &= \frac{1}{\alpha_{2n-1}} \prod_{r=2}^n \left( \left( \frac{\alpha_{2r-1}^2}{\alpha_{2r-1}^2 - 1} \right) \left( \frac{\alpha_{2r-2}^2 - 1}{\alpha_{2r-2}^2} \right) \right) \frac{\alpha_1^2}{\alpha_1^2 - 1}, \\ d_{2n+1} &= \frac{1}{\alpha_{2n+1}} \prod_{r=1}^n \left( \left( \frac{\alpha_{2r-1}^2 - 1}{\alpha_{2r-1}^2} \right) \left( \frac{\alpha_{2r}^2}{\alpha_{2r}^2 - 1} \right) \right), \\ e_{2n+1} &= \frac{1}{\alpha_{2n}} \prod_{r=1}^n \left( \left( \frac{\alpha_{2r-1}^2 - 1}{\alpha_{2r-1}^2} \right) \left( \frac{\alpha_{2r}^2}{\alpha_{2r}^2 - 1} \right) \right). \end{aligned}$$

*Proof.* From *ii)* of Theorem 3.1.1, and *iii)* of Theorem 3.1.2, we have the following Perron-Carathéodory continued fraction,

$$\frac{\Omega_n(z)}{\Phi_n(z)} = c_0 - \frac{2c_0}{1} - \frac{1}{\alpha_1 z} - \frac{(\alpha_1^2 - 1)z}{\alpha_1} - \dots - \frac{1}{\alpha_{n-1} z} - \frac{(\alpha_n^2 - 1)z}{\alpha_n}. \quad (3.29)$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

The results in Theorem 3.2.2 allow us to define the step function  $\mu_n$  by

$$\mu_n(x) = \begin{cases} 0, & 0 < x \leq z_{n,1}, \\ \sum_{r=1}^s \lambda_{n,r}, & z_{n,s} < x \leq z_{n,s+1}, \\ \mu_0, & z_{n,n} < x < \infty, \end{cases}$$

where  $z_{n,r}$  are the zeros of the polynomial  $\Phi_n$ . Then, from *i*) of Theorem 3.2.2,

$$\frac{\Omega_n(z)}{\Phi_n(z)} = \int_0^\infty \frac{x+z}{x-z} d\mu_n(x), \quad n \geq 1,$$

and

$$c_s = \int_0^\infty x^s d\mu_n(x), \quad -n+1 \leq s \leq n-1, \quad n \geq 1,$$

from the previous consideration, we have

$$\frac{w}{2} \left( \frac{c_0 \Phi_n(-w) + \Omega_n(-w)}{\Phi_n(-w)} \right) = \int_0^\infty \frac{w}{x+w} d\widehat{\mu}_n(x),$$

where  $d\widehat{\mu}(x) = x d\mu(x)$ . Expanding  $\int_0^\infty \frac{w}{x+w} d\widehat{\mu}_n(x)$ , we can easily verify that

$$\frac{\widehat{\Omega}_n(w)}{\widehat{\Phi}_n(w)} = \begin{cases} -a_{-1}w - a_{-2}w^2 - \dots - a_{-n}w^n + O(w^{(n+1)}), \\ a_0 + a_1w^{-1} + \dots + a_{n-1}w^{-(n-1)} + O(w^{-n}), \end{cases}$$

where

$$\widehat{\Omega}_n(w) = (-1)^n \frac{w}{2} (c_0 \Phi_n(-w) + \Omega_n(-w)), \quad \widehat{\Phi}_n(w) = (-1)^n \Phi_n(-w).$$

From Theorem 3.2.7,  $\det \mathbf{H}_{k+1}^{(-2n+l)} > 0$ ,  $k = 0, 1, \dots, 2n$ ,  $l = 0, \pm 1, \pm 2, \dots$ , we can prove that the Hankel determinants

$$\det \widehat{\mathbf{H}}_n^{(r)} = \begin{vmatrix} a_r & a_{r+1} & \dots & a_{r+n-1} \\ a_{r+1} & a_{r+2} & \dots & a_{r+n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{r+n-2} & a_{r+n-1} & \dots & a_{r+2n-3} \\ a_{r+n-1} & a_{r+n} & \dots & a_{r+2n-2} \end{vmatrix}, \quad n \geq 1, \quad (3.30)$$

satisfy the following conditions,

$$\det \widehat{\mathbf{H}}_{n+1}^{(-n)} > 0, \quad n \geq 0, \\ \det \widehat{\mathbf{H}}_{2n}^{(-2n)} > 0, \quad \det \widehat{\mathbf{H}}_{2n-1}^{(-(2n-1))} < 0, \quad n \geq 1. \quad (3.31)$$

Therefore, the Chebyshev moment problem can be identified as a strong Stieltjes moment problem. Then there exists a solution if (3.31) holds [101]. Obviously,  $d\widehat{\mu}(x) = x d\mu(x)$  is a solution, taking into account that

$$a_n = \int_0^\infty (-x)^n d\widehat{\mu}(x), \quad n \in \mathbb{Z}.$$

Furthermore, we have

$$\int_0^\infty \frac{w}{x+w} d\widehat{\mu}_{2n}(x) = \frac{\widehat{\Omega}_{2n}(w)}{\widehat{\Phi}_{2n}(w)} < \int_0^\infty \frac{w}{x+w} d\widehat{\mu}(x) < \frac{\widehat{\Omega}_{2n+1}(w)}{\widehat{\Phi}_{2n+1}(w)} = \int_0^\infty \frac{w}{x+w} d\widehat{\mu}_{2n+1}(x),$$

where  $w > 0$ . Moreover, there also exist solutions  $d\widehat{\mu}^{(0)}(x)$  and  $d\widehat{\mu}^{(1)}(x)$ , limits of  $d\widehat{\mu}_{2n}(x)$  and  $d\widehat{\mu}_{2n+1}(x)$ , respectively, such that

$$\int_0^\infty \frac{w}{x+w} d\widehat{\mu}^{(0)}(x) \leq \int_0^\infty \frac{w}{x+w} d\widehat{\mu}(t) \leq \int_0^\infty \frac{w}{x+w} d\widehat{\mu}^{(1)}(x).$$

If  $\widehat{\Omega}_n/\widehat{\Phi}_n$  converges, then the uniqueness of the solution is proved. Hence, we need to verify the convergence of the following Perron-Carathéodory continued fraction

$$\frac{\Omega_n(z)}{\Phi_n(z)} = \frac{\alpha_1 w}{w + \beta_1} + \frac{\alpha_2 w}{w + \beta_2} + \frac{\alpha_3 w}{w + \beta_3} + \dots, \quad (3.32)$$

where  $\alpha_1 = c_1$ ,  $\beta_n = \alpha_n/\alpha_{n-1}$ , and  $\alpha_n = \beta_n(\alpha_{n-1}^2 - 1)$ . Notice that (3.32) can be expressed as follows

$$\frac{\Omega_n(z)}{\Phi_n(z)} = \frac{w}{e_1 + d_1 \omega} + \frac{w}{e_2 + d_2 \omega} + \frac{w}{e_3 + d_3 \omega} + \dots, \quad (3.33)$$

where

$$e_1 = \frac{1}{\mu_0 \alpha_1}, \quad d_1 = \frac{1}{\mu_0}, \quad d_{2n} = \prod_{r=1}^n \left( \frac{\alpha_{2r-1}}{\alpha_{2r}} \right), \quad e_{2n} = \beta_{2n} d_{2n}, \\ d_{2n+1} = \frac{1}{\alpha_{2n+1} d_{2n}}, \quad e_{2n+1} = \beta_{2n+1} d_{2n+1}, \quad n \geq 1.$$

### 3. ON SPECIAL CLASSES OF SZEGŐ POLYNOMIALS

---

By [101], the Perron-Carathéodory continued fraction (3.33) converges uniformly on compact subsets of  $\{w; |\arg(w)| < \pi\}$  if and only if (3.28) holds, as required.  $\square$

## Chapter 4

# Spectral transformations of moment matrices

Our hero is the intrepid, yet sensitive matrix  $\mathbf{A}$ .

Our villain is  $\mathbf{E}$ , who keeps perturbing  $\mathbf{A}$ .

When  $\mathbf{A}$  is perturbed he puts on a crumpled hat:  $\tilde{\mathbf{A}} = \mathbf{A} + \mathbf{E}$ .

— G. W. Stewart and J. -G. Sun. *Matrix perturbation theory*. Academic Press, New York, 1990

It is very well known that the Gram matrices of the bilinear functionals associated with (2.1) and (2.23) in the canonical basis  $\{z^n\}_{n \geq 0}$  of  $\mathbb{P}$  are Hankel and Toeplitz matrices, respectively. The main objective of this chapter is to study the perturbation of a fixed moment of the corresponding moment matrix. We refer to [7; 8; 10; 14; 25; 26; 83; 111; 125; 147] for other contributions involving related perturbations of moment functionals.

This chapter is divided into two parts. In the first one, given a strongly regular Hankel matrix, and its associated sequence of moments – which defines a quasi-definite moment linear functional  $\mathcal{M}$  – we study the perturbation on one anti-diagonal of the corresponding Hankel matrix. We define a linear functional  $\mathcal{M}_j$ , whose action results in such a perturbation and we establish necessary and sufficient conditions in order to preserve the quasi-definite character. A relation between the corresponding sequences of orthogonal polynomials is obtained, as well as the asymptotic behavior of their zeros. In the second one, we analyze a general linear spectral transformation of  $\mathcal{L}$ ,  $\mathcal{L}_j$ , such

## 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

---

that the corresponding Toeplitz matrix is the result of the addition of a constant in two symmetric sub-diagonals of the initial Toeplitz matrix. We focus our attention on the analysis of the quasi-definite character of the perturbed linear functional, as well as on the explicit expressions of the new monic orthogonal polynomial sequence in terms of the first one. Some illustrative examples are pointed out.

### 4.1 Hankel matrices

Before introducing the problem to be analyzed in this section, let us briefly discuss one rather straightforward but interesting example where the moments are modified in a natural way. Instead of the canonical basis of  $\mathbb{P}$ , let us consider the basis  $\{1, (x - a), (x - a)^2, \dots\}$ , where  $a \in \mathbb{R}$ . Then, the new sequence of moments  $\{v_n\}_{n \geq 0}$  is given by

$$v_n = \langle \mathcal{M}, (x - a)^n \rangle = \left\langle \mathcal{M}, \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} a^{n-j} x^j \right\rangle = \sum_{j=0}^n \binom{n}{j} (-1)^{n-j} a^{n-j} \mu_j, \quad n \geq 0. \quad (4.1)$$

As a consequence,  $\widehat{\mathbf{H}}$  the Hankel matrix associated with the new basis is

$$\widehat{\mathbf{H}} = \left[ \langle \mathcal{M}, (x - a)^{i+j} \rangle \right]_{i,j \geq 0} = \begin{bmatrix} \mu_0 & \mu_1 + m_1 & \cdots & \mu_n + m_n & \cdots \\ \mu_1 + m_1 & \mu_2 + m_2 & \cdots & \mu_{n+1} + m_{n+1} & \cdots \\ \vdots & \vdots & \ddots & \vdots & \cdots \\ \mu_n + m_n & \mu_{n+1} + m_{n+1} & \cdots & \mu_{2n+1} + m_{2n+1} & \cdots \\ \vdots & \vdots & \vdots & \vdots & \ddots \end{bmatrix},$$

where  $m_0 = 0$  and  $m_i = \sum_{j=1}^{i-1} \binom{i}{j} (-1)^{i-j} a^{i-j} \mu_j$ . Thus, if  $\mathcal{M}$  is a quasi-definite moment linear functional, then the polynomials

$$\widehat{P}_n(x) = \frac{1}{\det \widehat{\mathbf{H}}_{n-1}} \begin{vmatrix} v_0 & v_1 & v_2 & \cdots & v_n \\ v_1 & v_2 & v_3 & \cdots & v_{n+1} \\ \vdots & \vdots & \cdots & \vdots & \vdots \\ v_{n-1} & v_n & v_{n+1} & \cdots & v_{2n-1} \\ 1 & (x - a) & (x - a)^2 & \cdots & (x - a)^n \end{vmatrix}, \quad n \geq 0, \quad (4.2)$$



---

constitute a sequence of monic polynomials orthogonal with respect to  $\mathcal{M}$ , using the new basis, where  $\widehat{P}_n(x) = P_n(x)$ . Notice that this simple change of the basis results in a perturbation on the anti-diagonals of the original moment matrix (2.2). Namely, the  $j$ -th anti-diagonal is perturbed by the addition of the constant  $m_j$ . In the remaining of the manuscript, we will use the basis  $\{1, (x-a), (x-a)^2, \dots\}$ , since most of the required calculations can be performed in a most simple way. Now a natural question arises: Is there a linear functional  $\mathcal{M}_j$  such that its action results on a perturbation of (only) the moment  $u_j$  or, equivalently, the  $(j+1)$ -th anti-diagonal of the corresponding Hankel matrix  $\widetilde{\mathbf{H}}$ ? In other words, we are interested in the properties of the functional  $\mathcal{M}_j$  whose moments are given by

$$\widetilde{v}_n = \langle \mathcal{M}_j, (x-a)^n \rangle = \begin{cases} u_n, & n \neq j, \\ u_n + m, & n = j, \end{cases} \quad (4.3)$$

for some  $m \in \mathbb{R}$ .

#### 4.1.1 Perturbation on the anti-diagonals of a Hankel matrix

In order to state our main result, we need some definitions. Given a moment linear functional  $\mathcal{M}$ , the usual distributional derivative  $D\mathcal{M}$  [135] is given by

$$\langle D\mathcal{M}, p \rangle = -\langle \mathcal{M}, p' \rangle, \quad p \in \mathbb{P}.$$

In particular, if  $j$  is a non-negative integer, then

$$\langle D^{(j)}\delta(x-a), p(x) \rangle = (-1)^j p^{(j)}(a).$$

In the last years, modifications by means of real Dirac's deltas have been considered by several authors from different points of view (hypergeometric character, holonomic equations that these polynomials satisfy, electrostatic interpretation of zeros, asymptotic properties, among others). In a pioneering result by Krall [111] were obtained three new classes of polynomials orthogonal with respect to measures which are not absolutely continuous with respect to the Lebesgue measure; the resulting polynomials satisfy a fourth-order linear differential equation. Nevai [147] considered the addition of finitely many mass points to a positive measure supported on a bounded subset of the real line, and studied the asymptotic behavior of the corresponding orthogonal

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

---

polynomials.

We begin with a simple but important remark. The moment linear functional  $\mathcal{M}_j$  discussed in the introduction of this chapter is given by

$$\langle \mathcal{M}_j, p \rangle = \langle \mathcal{M}, p \rangle + (-1)^j \frac{m_j}{j!} \langle D^{(j)} \delta(x-a), p(x) \rangle = \langle \mathcal{M}, p \rangle + \frac{m_j}{j!} p^{(j)}(a), \quad a \in \mathbb{R}. \quad (4.4)$$

It is easy to see that all the moments associated with  $\mathcal{M}_j$  are equal to the moments  $\{v_n\}_{n \geq 0}$  of  $\mathcal{M}$  in the basis  $\{1, (x-a), (x-a)^2, \dots\}$ , except for the  $j$ -th one, which is equal to  $v_j + m_j$ . Notice that this perturbation is the simplest one that preserves the Hankel structure of the moment matrix. We can now state the result, which establishes necessary and sufficient conditions in order to the linear functional  $\mathcal{M}_j$  preserves the quasi-definite character, and we provide the relation between the corresponding sequences of monic orthogonal polynomials.

**Theorem 4.1.1.** *Let  $\mathcal{M}$  be a quasi-definite moment linear functional and  $\{P_n\}_{n \geq 0}$  its corresponding sequence of monic orthogonal polynomials. Then, the following statements are equivalent:*

i) *The moment linear functional  $\mathcal{M}_j$ , defined in (4.4), is quasi-definite.*

ii) *The matrix  $\mathbf{I}_{j+1} + \mathbf{K}_{j+1} \mathbf{D}_{j+1}$ , where*

$$\mathbf{D}_{j+1} = \frac{m}{j!} \begin{bmatrix} \binom{j}{j} & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \binom{j}{0} \end{bmatrix}, \quad \mathbf{K}_{j+1} = \begin{bmatrix} K_{n-1}^{(j,0)}(a, a) & K_{n-1}^{(j-1,0)}(a, a) & \cdots & K_{n-1}^{(0,0)}(a, a) \\ K_{n-1}^{(j,1)}(a, a) & K_{n-1}^{(j-1,1)}(a, a) & \cdots & K_{n-1}^{(0,1)}(a, a) \\ \vdots & \vdots & \ddots & \vdots \\ K_{n-1}^{(j,j)}(a, a) & K_{n-1}^{(j-1,j)}(a, a) & \cdots & K_{n-1}^{(0,j)}(a, a) \end{bmatrix},$$

*is non-singular, and*

$$\langle \mathcal{M}_j, P_n(j; \cdot) P_n \rangle = \langle \mathcal{M}, P_n^2 \rangle + \begin{bmatrix} P_n^{(j)}(a) \\ P_n^{(j-1)}(a) \\ \vdots \\ P_n(a) \end{bmatrix}^T \mathbf{D}_{j+1} (\mathbf{I}_{j+1} + \mathbf{K}_{j+1} \mathbf{D}_{j+1})^{-1} \begin{bmatrix} P_n(a) \\ P_n^{(1)}(a) \\ \vdots \\ P_n^{(j)}(a) \end{bmatrix} \neq 0.$$

*Moreover, if both  $\mathcal{M}$  and  $\mathcal{M}_j$  are quasi-definite, and  $\{P_n(j; \cdot)\}_{n \geq 0}$  is the sequence of*

---

monic orthogonal polynomials with respect to  $\mathcal{M}_j$ , then

$$P_n(j; x) = P_n(x) - \begin{bmatrix} K_{n-1}^{(j,0)}(a, x) \\ K_{n-1}^{(j-1,0)}(a, x) \\ \vdots \\ K_{n-1}^{(0,0)}(a, x) \end{bmatrix}^T \mathbf{D}_{j+1}(\mathbf{I}_{j+1} + \mathbf{K}_{j+1}\mathbf{D}_{j+1})^{-1} \begin{bmatrix} P_n(a) \\ P_n^{(1)}(a) \\ \vdots \\ P_n^{(j)}(a) \end{bmatrix}. \quad (4.5)$$

*Proof.* Suppose that  $\mathcal{M}_j$  is a quasi-definite moment linear functional. Since  $\{P_n\}_{n \geq 0}$  is the sequence of monic orthogonal polynomials with respect to  $\mathcal{M}$ , there exist constants  $\lambda_{n,0}, \dots, \lambda_{n,n-1}$  such that

$$P_n(j; x) = P_n(x) + \sum_{k=0}^{n-1} \lambda_{n,k} P_k(x). \quad (4.6)$$

Thus, by the orthogonality property,

$$\lambda_{n,k} = -\frac{m_j}{j!} \frac{\sum_{l=0}^r \binom{j}{l} P_n^{(j-l)}(j; a) P_k^{(l)}(a)}{\langle \mathcal{M}, P_k^2 \rangle}, \quad r = \min\{k, j\}, \quad 0 \leq k \leq n-1.$$

Substituting the above expression in (4.6), and taking the  $i$ -th derivative of  $P_n(j; \cdot)$ , we obtain

$$P_n^{(i)}(j; x) = P_n^{(i)}(x) - \frac{m_j}{j!} \sum_{l=0}^j \binom{j}{l} P_n^{(j-l)}(j; a) K_{n-1}^{(l,i)}(a, x), \quad i = 0, 1, \dots, j. \quad (4.7)$$

Evaluating (4.7) at  $x = a$ , we have the following linear system

$$(\mathbf{I}_{j+1} + \mathbf{K}_{j+1}\mathbf{D}_{j+1}) \begin{bmatrix} P_n(j; a) \\ P_n^{(1)}(j; a) \\ \vdots \\ P_n^{(j)}(j; a) \end{bmatrix} = \begin{bmatrix} P_n(a) \\ P_n^{(1)}(a) \\ \vdots \\ P_n^{(j)}(a) \end{bmatrix}.$$

Since  $\mathcal{M}_j$  is quasi-definite, there exists a unique sequence of monic orthogonal polynomials with respect to  $\mathcal{M}_j$ . Thus, the linear system has a unique solution and therefore the  $(j+1) \times (j+1)$  matrix  $\mathbf{I}_{j+1} + \mathbf{K}_{j+1}\mathbf{D}_{j+1}$  is non-singular. Furthermore, (4.7) with  $i = 0$  reduces to (4.5).

## 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

---

On the other hand, notice that for  $m = 0, 1, \dots, n-1$ , we have

$$\begin{aligned} \langle \mathcal{M}_j, P_n(j; \cdot) P_m \rangle &= \langle \mathcal{M}, P_n P_m \rangle \\ &= -\frac{m_j}{j!} \sum_{l=0}^j \binom{j}{l} P_n^{(j-l)}(j; a) P_m^{(l)}(a) + \frac{m_j}{j!} (P_n(j; x) P_m(x))^{(j)} \Big|_{x=a} = 0, \end{aligned}$$

and

$$0 \neq \langle \mathcal{M}_j, P_n(j; \cdot) P_n \rangle = \langle \mathcal{M}, P_n^2 \rangle + \begin{bmatrix} P_n^{(j)}(a) \\ P_n^{(j-1)}(a) \\ \vdots \\ P_n(a) \end{bmatrix}^T \mathbf{D}_{j+1} (\mathbf{I}_{j+1} + \mathbf{K}_{j+1} \mathbf{D}_{j+1})^{-1} \begin{bmatrix} P_n(a) \\ P_n^{(1)}(a) \\ \vdots \\ P_n^{(j)}(a) \end{bmatrix}. \quad (4.8)$$

For the converse, let us assume that *ii*) holds, and define  $\{P_n(j; \cdot)\}_{n \geq 0}$  as in (4.5). Then it is straightforward to show that  $\{P_n(j; \cdot)\}_{n \geq 0}$  is the sequence of monic orthogonal polynomials with respect to  $\mathcal{M}_j$ , and its quasi-definite character is proved.  $\square$

### 4.1.2 Zeros

In this subsection we assume that the linear functional  $\mathcal{M}$  is positive definite, i.e., it has an associated positive measure supported on some interval  $I \subseteq \mathbb{R}$ , and that  $\mathcal{M}_j$  is quasi-definite. We show some properties regarding the zeros of its corresponding sequence of monic orthogonal polynomials.

Let  $x_1, \dots, x_r$  be the zeros of  $P_n(j; \cdot)$  on  $I$  with odd multiplicity, and define  $Q_r(x) = (x - x_1) \dots (x - x_r)$ . Then,  $P_n(j; x) Q_r(x) (x - a)^{2k}$ , where  $k$  is the smallest integer such that  $k \geq (j+1)/2$ , is a polynomial that does not change sign on  $I$  and, furthermore, we have

$$\langle \mathcal{M}_j, P_n(j; x) Q_r(x) (x - a)^{2k} \rangle = \langle \mathcal{M}, P_n(j; x) Q_r(x) (x - a)^{2k} \rangle \neq 0.$$

From the orthogonality of  $P_n(j; \cdot)$  with respect to  $\mathcal{M}_j$  we have

**Theorem 4.1.2.** *For  $n > 2k$ , the polynomial  $P_n(j; \cdot)$  has at least  $n - 2k$  different zeros with odd multiplicity on  $I$ .*

We now analyze the asymptotic behavior of the zeros of  $\{P_n(j; \cdot)\}_{n \geq 0}$  when the mass

$m_j$  tends to infinity. Notice that,

$$P_n(j; x) = \frac{1}{\det \widetilde{\mathbf{H}}_{n-1}} \begin{bmatrix} u_0 & \cdots & u_j + m_j & \cdots & u_n \\ \vdots & & \vdots & & \vdots \\ u_j + m_j & & u_{2j} & & u_{n+j} \\ \vdots & & \vdots & \ddots & \vdots \\ u_{n-1} & & u_{n+j-1} & & u_{2n-1} \\ 1 & \cdots & (x-a)^j & \cdots & (x-a)^n \end{bmatrix}. \quad (4.9)$$

On the other hand, let  $\{R_n^k(a; \cdot)\}_{n \geq 0}$  be the sequence of monic orthogonal polynomials with respect to the linear functional  $\widehat{\mathcal{M}}$  defined by

$$\langle \widehat{\mathcal{M}}, p \rangle = \langle \mathcal{M}, (x-a)^k p(x) \rangle, \quad k \geq 0.$$

Here, we assume that  $a$  is not a zero of the polynomials  $\{P_n\}_{n \geq 0}$  in order  $\widehat{\mathcal{M}}$  to be quasi-definite. The Hankel matrix associated with  $\widehat{\mathcal{M}}$  is

$$\mathbf{H}^{(k)} = \begin{bmatrix} \mu_k & \mu_{1+k} & \cdots & \mu_{n+k} & \cdots \\ \mu_{1+k} & \mu_{2+k} & \cdots & \mu_{n+1+k} & \cdots \\ \vdots & \vdots & \ddots & \vdots & \\ \mu_{n+k} & \mu_{n+1+k} & \cdots & \mu_{2n+k} & \cdots \\ \vdots & \vdots & & \vdots & \ddots \end{bmatrix}.$$

From (4.2),

$$R_{n-j-1}^{2(j+1)}(a; x) = \frac{1}{\det \mathbf{H}_{n-j-2}^{(2j+2)}} \begin{bmatrix} u_{2j+2} & u_{2j+3} & \cdots & u_{n+j} & u_{n+j+1} \\ u_{2j+3} & u_{2j+4} & \cdots & u_{n+j+1} & u_{n+j+2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ u_{n+j} & u_{n+j+1} & \cdots & u_{2n-2} & u_{2n-1} \\ 1 & (x-a) & \cdots & (x-a)^{n-j-2} & (x-a)^{n-j-1} \end{bmatrix}, \quad n > j+1,$$

If the matrix in (4.9) is block partitioned into  $\begin{bmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{bmatrix}$ , where  $\mathbf{A}$  is a  $(j+1) \times (j+1)$  matrix, then

$$\begin{vmatrix} \mathbf{A} & \mathbf{B} \\ \mathbf{C} & \mathbf{D} \end{vmatrix} = \det \mathbf{D} \det(\mathbf{A} - \mathbf{B} \mathbf{D}^{-1} \mathbf{C}).$$

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

---

It is clear that  $\det \mathbf{D} = \det \mathbf{H}_{n-j-2}^{(2j+2)}(x-a)^{j+1} R_{n-j-1}^{2(j+1)}(a; x)$ . Moreover,  $\mathbf{B}\mathbf{D}^{-1}\mathbf{C}$  is a  $(j+1) \times (j+1)$  matrix that does not depend on  $m$ , and thus

$$P_n(j; x) = \frac{\det \mathbf{H}_{n-j-2}^{(2j+2)}(x-a)^{j+1} R_{n-j-1}^{2(j+1)}(a; x) Q(m_j)}{\det \mathbf{H}_{n-j-2}^{(2j+2)} R(m_j)}, \quad (4.10)$$

where  $Q(m_j)$  and  $R(m_j)$  are monic polynomials in  $m_j$  of degree  $j+1$ . Therefore,

$$\lim_{m_j \rightarrow \infty} P_n(j; x) = (x-a)^{j+1} R_{n-(j+1)}^{2(j+1)}(a; x), \quad (4.11)$$

and we conclude

**Theorem 4.1.3.** *The zeros  $x_{n,k}(j; m_j)$ ,  $k = 1, \dots, n$ , of the polynomial  $P_n(j; \cdot)$  converge to the zeros of the polynomial  $(x-a)^{j+1} R_{n-(j+1)}^{2(j+1)}(a; \cdot)$  when  $m_j$  tends to infinity.*

Observe that when  $m_j$  tends to infinity the mass point  $a$  attracts  $j+1$  zeros of  $P_n(j; \cdot)$ .

A rather natural question is if  $x_{n,k}(j; m_j)$ , considered as functions of  $m_j$ , tend to the zeros of  $(x-a)^{j+1} R_{n-(j+1)}^{2(j+1)}(a; x)$  in a monotonic way. For the particular case when  $j=0$ , it was proved [61; 62] that the zeros of the so-called Laguerre and Jacobi type orthogonal polynomials, which are particular cases of the Uvarov spectral transformation, behave monotonically with respect to  $m_j$ . Unfortunately, this phenomenon does not occur for every positive integer  $j$ . We have performed some numerical experiments with specific classical measures. For example, if the initial measure is the one associated with the Laguerre polynomials  $L_n^{(\alpha)}$ ,  $j=1$ , and  $a=0$ , then the zeros  $x_{n,k}(1; m)$  of the corresponding polynomials  $L_n(1; \cdot)$  converge to those of  $x^2 L_{n-2}^{\alpha+4}(x)$ , although they are not monotonic functions of  $m_j$  when it varies in  $(0, \infty)$ .

We present some tables that show the behavior of the zeros of  $T_n(j; \cdot)$  as a function of  $m_j$ , when the initial measure is  $d\mu(-1/2, -1/2; x)$  (Chebyshev polynomials of the first kind; see Chapter 1) for  $j=1$  and  $n=2, 3$ .

Notice that there exist complex zeros depending on the values of the parameter  $m_j$ . It is also observed that two zeros of the polynomial approach the point  $x=a$  as  $m_j$  increases, as established in Theorem 4.1.3.

Table 4.1: Zeros of  $T_2(j; \cdot)$  for  $a = 3$ ,  $j = 1$  for some values of  $m$

$m$	$x_{2,1}(j; m)$	$x_{2,2}(j; m)$
0	-0.707107	0.707107
0.1	-0.4034	1.61713
0.5	-0.317089	3.16098
1	-0.355225	3.74581
5	-0.95446	4.54373
10	-2.39228	4.881
$10^2$	$3.18582 + 1.08651i$	$3.18582 - 1.08651i$
$10^3$	$3.01522 + 0.317155i$	$3.01522 - 0.317155i$
$10^4$	$3.0015 + 0.0995593i$	$3.0015 - 0.0995593i$
$10^5$	$3.00015 + 0.0314605i$	$3.00015 - 0.0314605i$

Table 4.2: Zeros of  $T_3(j; \cdot)$  for  $a = 3$ ,  $j = 1$  for some values of  $m$

$m$	$x_{3,1}(j; m)$	$x_{3,2}(j; m)$	$x_{3,3}(j; m)$
0	-0.866025	0	0.866025
0.1	-0.801321	0.51227	3.61105
0.5	-1.18576	0.0479705	3.7638
1	-3.70458	-0.305553	3.86415
5	-0.510437	$3.38703 + 0.805069i$	$3.38703 - 0.805069i$
10	-0.525903	$3.15823 + 0.549961i$	$3.15823 - 0.549961i$
$10^2$	-0.538469	$3.01358 + 0.167363i$	$3.01358 - 0.167363i$
$10^3$	-0.539661	$3.00134 + 0.052716i$	$3.00134 - 0.052716i$
$10^4$	-0.539779	$3.00013 + 0.0166637i$	$3.00013 - 0.0166637i$
$10^5$	-0.539791	$3.00001 + 0.0052693i$	$3.00001 - 0.0052693i$

## 4.2 Toeplitz matrices

From the point of view of perturbations of positive definite hermitian Toeplitz matrices or, equivalently, probability measures supported on the unit circle, there is a wide literature [21; 54; 78; 81; 82; 117; 127; 128], emphasizing the analytic properties of orthogonal polynomials with respect to the perturbed measures. In [25] a spectral transformation associated with a modification of a measure on the unit circle by the addition of the normalized Lebesgue measure was introduced. The translation, in terms of the entries of the new Toeplitz matrix, means that we only perturb the main diagonal of the original Toeplitz matrix. In the introduction of their work the authors emphasized that this problem is strongly related with the method introduced by Pisarenko [158] for retrieving harmonics from a covariance function. Four years later in [26], the same authors generalized their previous result and studied the sequence of orthogonal poly-

## 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

---

nomials associated with the sum of a measure supported on the unit circle in the class  $\mathcal{S}$  and a Bernstein-Szegő measure [178]. Indeed, they deduced that the corresponding measure belongs to the  $\mathcal{S}$  class and obtained several properties about the norms of the associated sequence of orthogonal polynomials.

### 4.2.1 Diagonal perturbation of a Toeplitz matrix

Let  $\mathcal{L}$  be the moment linear functional introduced in (2.23). We define a new linear functional  $\mathcal{L}_0$  such that its associated bilinear form satisfies

$$\langle f, g \rangle_{\mathcal{L}_0} = \langle f, g \rangle_{\mathcal{L}} + m \int_{\mathbb{T}} f(z) \overline{g(z)} \frac{dz}{2\pi iz}, \quad f, g \in \mathbb{P}, \quad m \in \mathbb{R}. \quad (4.12)$$

Assume that  $\mathcal{L}$  is a positive definite linear functional and let  $\sigma$  be its corresponding measure as defined in (2.29). Then, this transformation can be expressed in terms of the corresponding measure  $\sigma$  as

$$d\tilde{\sigma}(z) = d\sigma(z) + m \frac{dz}{2\pi iz}, \quad (4.13)$$

i.e., the addition of a Lebesgue measure to  $\sigma$ . We assume  $m \in \mathbb{R}_+$  in order  $\tilde{\sigma}$  to be a positive Borel measure supported in  $\mathbb{T}$ . The moments  $\{\tilde{c}_k\}_{k \in \mathbb{Z}}$  associated with  $\mathcal{L}_0$  are given by

$$\tilde{c}_0 = c_0 + m, \quad \tilde{c}_k = c_k, \quad k \in \mathbb{Z}. \quad (4.14)$$

As a consequence, the  $C$ -function of the linear functional  $\mathcal{L}_0$  is

$$F_0(z) = F(z) + m. \quad (4.15)$$

Notice that  $\tilde{\mathbf{T}}$ , the Toeplitz matrix associated with  $\mathcal{L}_0$ , is

$$\tilde{\mathbf{T}} = \mathbf{T} + m\mathbf{I}, \quad (4.16)$$

i.e., a constant is added to the main diagonal of  $\mathbf{T}$ . We now proceed to obtain the sequence of monic orthogonal polynomials with respect to  $\mathcal{L}_0$ .

**Theorem 4.2.1.** *Let  $\mathcal{L}$  be a positive definite linear functional, and denote by  $\{\Phi_n\}_{n \geq 0}$  its associated sequence of monic orthogonal polynomials. Then,  $\{\Psi_n\}_{n \geq 0}$ , the sequence*



of monic polynomials orthogonal with respect to  $\mathcal{L}_0$  defined by (4.12), is given by

$$\Psi_n(z) = \Phi_n(z) - \mathbf{K}_{n-1}^T(z, 0)(m^{-1}\mathbf{D}_n^{-2} + \mathbf{P}_{n-1}\mathbf{P}_{n-1}^T)^{-1}\Phi_n(0), \quad (4.17)$$

with  $\mathbf{K}_{n-1}(z, 0) = [K_{n-1}(z, 0), K_{n-1}^{(0,1)}(z, 0), \dots, K_{n-1}^{(0,n-1)}(z, 0)]^T$ ,  $\mathbf{D}_n = \text{diag}\{\frac{1}{0!}, \dots, \frac{1}{(n-1)!}\}$ ,  $\Phi_n(0) = [\Phi_n(0), \Phi_n'(0), \dots, \Phi_n^{(n-1)}(0)]^T$ , and

$$\mathbf{P}_{n-1} = \begin{bmatrix} \phi_0(0) & \phi_1(0) & \cdots & \phi_{n-1}(0) \\ 0 & \phi_1'(0) & \cdots & \phi_{n-1}'(0) \\ \vdots & 0 & \ddots & \vdots \\ 0 & \cdots & 0 & \phi_{n-1}^{(n-1)}(0) \end{bmatrix}.$$

*Proof.* Set

$$\Psi_n(z) = \Phi_n(z) + \sum_{k=0}^{n-1} \lambda_{n,k} \Phi_k(z),$$

where, for  $0 \leq k \leq n-1$ ,

$$\lambda_{n,k} = \frac{1}{\mathbf{k}_k} \langle \Psi_n, \Phi_k \rangle_{\mathcal{L}_0} - m \int_{\mathbb{T}} \Psi_n(y) \overline{\Phi_k(y)} \frac{dy}{2\pi iy} = -\frac{m}{\mathbf{k}_k} \int_{\mathbb{T}} \Psi_n(y) \overline{\Phi_k(y)} \frac{dy}{2\pi iy}.$$

Thus,

$$\Psi_n(z) = \Phi_n(z) - m \int_{\mathbb{T}} \Psi_n(y) K_{n-1}(z, y) \frac{dy}{2\pi iy} = \Phi_n(z) - m \sum_{j=0}^{n-1} \frac{\Psi_n^{(j)}(0)}{j!} \frac{K_{n-1}^{(0,j)}(z, 0)}{j!}. \quad (4.18)$$

In particular, for  $0 \leq i \leq n-1$ , we get

$$\Psi_n^{(i)}(0) = \Phi_n^{(i)}(0) - m \sum_{j=0}^{n-1} \frac{\Psi_n^{(j)}(0)}{j!} \frac{K_{n-1}^{(i,j)}(0, 0)}{j!}. \quad (4.19)$$

So, we have the following system of  $n$  linear equations and  $n$  unknowns

$$\Psi_n^{(i)}(0) = \Phi_n^{(i)}(0) - m \sum_{j=0}^{n-1} \frac{\Psi_n^{(j)}(0)}{j!} \frac{K_{n-1}^{(i,j)}(0, 0)}{j!}, \quad i = 0, 1, \dots, n-1,$$

which reads as

$$(\mathbf{I}_n + m\mathbf{R}_{n-1}\mathbf{D}_n^2)\Psi_n(0) = \Phi_n(0), \quad (4.20)$$

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

where  $\Psi_n(0) = [\Psi_n(0), \dots, \Psi_n^{(n-1)}(0)]^T$ ,

$$\mathbf{R}_{n-1} = \begin{bmatrix} K_{n-1}^{(0,0)}(0,0) & K_{n-1}^{(0,1)}(0,0) & \cdots & K_{n-1}^{(0,n-1)}(0,0) \\ K_{n-1}^{(1,0)}(0,0) & K_{n-1}^{(1,1)}(0,0) & \cdots & K_{n-1}^{(1,n-1)}(0,0) \\ \vdots & \vdots & \ddots & \vdots \\ K_{n-1}^{(n-1,0)}(0,0) & K_{n-1}^{(n-1,1)}(0,0) & \cdots & K_{n-1}^{(n-1,n-1)}(0,0) \end{bmatrix}, \quad (4.21)$$

As a consequence, (4.20) becomes  $\Psi_n(0) = m^{-1}(m^{-1}\mathbf{I}_n + \mathbf{R}_{n-1}\mathbf{D}_n^2)^{-1}\Phi_n(0)$ . Thus, (4.18) can be written

$$\Psi_n(z) = \Phi_n(z) - m\mathbf{K}_{n-1}^T(z,0)\mathbf{D}_n^2\Psi_n(0) = \Phi_n(z) - \mathbf{K}_{n-1}^T(z,0)(m^{-1}\mathbf{D}_n^{-2} + \mathbf{R}_{n-1})^{-1}\Phi_n(0),$$

which is (4.17), since  $\mathbf{R}_{n-1} = \mathbf{P}_{n-1}\mathbf{P}_{n-1}^T$ .  $\square$

On the other hand, considering the derivatives of order  $j$  with respect to the variable  $z$  in the Christoffel-Darboux formula (2.33) we get

$$K_{n-1}^{(j,0)}(z,y) = \frac{\overline{\Phi_n^*(y)}}{\mathbf{k}_n} \left( \frac{\Phi_n^*(z)}{1-\bar{y}z} \right)^{(j)} - \frac{\overline{\Phi_n(y)}}{\mathbf{k}_n} \left( \frac{\Phi_n(z)}{1-\bar{y}z} \right)^{(j)}.$$

Thus,

$$K_{n-1}^{(j,0)}(0,y) = \frac{\overline{\Phi_n^*(y)}}{\mathbf{k}_n} \left( \frac{\Phi_n^*(z)}{1-\bar{y}z} \right)^{(j)}(0) - \frac{\overline{\Phi_n(y)}}{\mathbf{k}_n} \left( \frac{\Phi_n(z)}{1-\bar{y}z} \right)^{(j)}(0).$$

Now, by Leibniz rule

$$\left( \frac{\Phi_n(z)}{1-\bar{y}z} \right)^{(j)} = \sum_{k=0}^j \binom{j}{k} \Phi_n^{(j-k)}(z) \frac{k! \bar{y}^k}{(1-\bar{y}z)^{k+1}},$$

the evaluation at  $z = 0$  yields

$$j! \sum_{k=0}^j \frac{\Phi_n^{(j-k)}(0)}{(j-k)!} \bar{y}^k = j! \sum_{k=0}^j \frac{\overline{\Phi_n^{(j-k)}(0)}}{(j-k)!} \bar{y}^k = j! \overline{T_j^*(\Phi_n(y); 0)},$$

where  $T_j(p(y); 0)$  denotes the  $j$ -th Taylor polynomial of  $p(y)$  around  $y = 0$ . In an analog way,

$$j! \sum_{k=0}^j \frac{\Phi_n^{*(j-k)}(0)}{(j-k)!} \bar{y}^k = j! \overline{T_j^*(\Phi_n^*(y); 0)}.$$

Therefore, we have proved

**Theorem 4.2.2.**

$$K_{n-1}^{(0,j)}(z, 0) = j! \left( \frac{\Phi_n^*(z)}{\mathbf{k}_n} T_j^*(\Phi_n^*(z); 0) - \frac{\Phi_n(z)}{\mathbf{k}_n} T_j^*(\Phi_n(z); 0) \right).$$

From the previous theorem, if we denote

$$\mathbf{T}(\Phi_n(z); 0) = \left[ T_0^*(\Phi_n(z); 0), T_1^*(\Phi_n(z); 0), \dots, T_{n-1}^*(\Phi_n(z); 0) \right]^T,$$

then, (4.17) becomes

$$\begin{aligned} \Psi_n(z) &= \Phi_n(z) - \frac{\Phi_n(z)}{\mathbf{k}_n} \mathbf{T}^T(\Phi_n(z); 0) \mathbf{D}_n^{-1} (m^{-1} \mathbf{D}_n^{-2} + \mathbf{P}_{n-1} \mathbf{P}_{n-1}^T)^{-1} \Phi_n(0) \\ &\quad - \frac{\Phi_n^*(z)}{\mathbf{k}_n} \mathbf{T}^T(\Phi_n^*(z); 0) \mathbf{D}_n^{-1} (m^{-1} \mathbf{D}_n^{-2} + \mathbf{P}_{n-1} \mathbf{P}_{n-1}^T)^{-1} \Phi_n(0) \end{aligned} \quad (4.22)$$

$$= a(z; n) \Phi_n(z) + b(z; n) \Phi_n^*(z), \quad (4.23)$$

where

$$a(z; n) = 1 - \frac{1}{\mathbf{k}_n} \mathbf{T}^T(\Phi_n(z); 0) \mathbf{D}_n^{-1} (m^{-1} \mathbf{D}_n^{-2} + \mathbf{P}_{n-1} \mathbf{P}_{n-1}^T)^{-1} \Phi_n(0),$$

$$b(z; n) = -\frac{1}{\mathbf{k}_n} \mathbf{T}^T(\Phi_n^*(z); 0) \mathbf{D}_n^{-1} (m^{-1} \mathbf{D}_n^{-2} + \mathbf{P}_{n-1} \mathbf{P}_{n-1}^T)^{-1} \Phi_n(0).$$

**Example: Bernstein-Szegő polynomials**

Let us consider the measure  $\sigma$  such that

$$d\sigma(\theta) = \frac{1 - |\beta|^2}{|e^{i\theta} - \beta|^2} \frac{d\theta}{2\pi}, \quad |\beta| < 1, \quad (4.24)$$

and apply the transformation studied in this section, i.e., let us define a new measure  $\tilde{\sigma}$

$$d\tilde{\sigma}(\theta) = \frac{1 - |\beta|^2}{|e^{i\theta} - \beta|^2} \frac{d\theta}{2\pi} + m \frac{d\theta}{2\pi}, \quad m \in \mathbb{R}_+. \quad (4.25)$$

Our aim is to find the sequence of monic polynomials orthogonal with respect to (4.25), that will be denoted by  $\{\Psi_n\}_{n \geq 0}$ . (4.24) is known in the literature as Bernstein-Szegő measure, and its corresponding sequence of monic orthogonal polynomials is [178],  $\Phi_n(z) = z^{n-1}(z - \beta)$ ,  $n \geq 1$ . Furthermore, we have  $\Phi_n^*(z) = 1 - \bar{\beta}z$ ,  $n \geq 1$ . Notice

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

that for  $n = 1$ , from (4.23) we get

$$\Psi_1(z) = \left(1 - \beta^2(1 - |\beta|^2)\left(m + (1 - |\beta|^2)^{-1}\right)\right)(z - \beta) - \left(\beta(1 - |\beta|^2)\left(m + (1 - |\beta|^2)^{-1}\right)\right)(1 - \bar{\beta}z).$$

For  $n \geq 2$ ,  $T_j(\Phi_n(0); 0) = 0$ ,  $0 \leq j \leq n-2$ , and  $T_{n-1}(\Phi_n(z); 0) = -\beta z^{n-1}$ . Thus,  $\mathbf{T}(\Phi_n(z); 0) = [0, 0, \dots, -\bar{\beta}]^T$ . On the other hand,  $T_0(\Phi_n^*(0); 0) = 1$ , and  $T_j(\Phi_n^*(z); 0) = 1 - \bar{\beta}z$ ,  $1 \leq j \leq n-1$ . Therefore,  $\mathbf{T}(\Phi_n^*(z); 0) = [1, z - \beta, z(z - \beta), \dots, z^{n-2}(z - \beta)]^T$ . Furthermore, in this case

$$\mathbf{P}_{n-1} = \frac{1}{\sqrt{1 - |\beta|^2}} \mathbf{D}_n^{-1} \mathbf{B}_n,$$

where

$$\mathbf{B}_n = \begin{bmatrix} 1 & \beta & 0 & \cdots & 0 \\ 0 & 1 & \beta & \ddots & \vdots \\ \vdots & \ddots & \ddots & \ddots & 0 \\ \vdots & & \ddots & 1 & \beta \\ 0 & \cdots & \cdots & 0 & 1 \end{bmatrix}. \quad (4.26)$$

Denoting by

$$\mathbf{N}_n = \mathbf{D}_n^{-1} \left( m^{-1} \mathbf{D}_n^{-2} + \mathbf{P}_{n-1} \mathbf{P}_{n-1}^T \right)^{-1} = \left( m^{-1} \mathbf{I}_n + \frac{1}{1 - |\beta|^2} \mathbf{B}_n \mathbf{B}_n^t \right)^{-1} \mathbf{D}_n,$$

and taking into account that  $\Phi_n(0) = [0, 0, \dots, -(n-1)! \beta]^T$ , we get

$$a(z; n) = 1 - \frac{|\beta|^2}{1 - |\beta|^2} (n-1)! n_{n-1, n-1},$$

$$b(z; n) = -\frac{\beta}{1 - |\beta|^2} (n-1)! \left( n_{0, n-1} + (z - \beta) \sum_{i=0}^{n-2} n_{i+1, n-1} z^i \right),$$

where  $n_{i,j}$ ,  $0 \leq i, j \leq n-1$ , are the entries of  $\mathbf{N}_n$ . From this, all elements on (4.23) are known and we can compute  $\Psi_n$ . The corresponding Verblunsky coefficients for  $n \geq 2$  are

$$\Psi_n(0) = b(0; n) = \frac{\beta}{1 - |\beta|^2} (n-1)! (\beta n_{1, n-1} - n_{0, n-1}).$$

The first 20 Verblunsky coefficients for different values of  $m$  and  $\beta$  are shown in Figure 4.1(a), namely  $m = 5$ ,  $\beta = -0.2$  (blue discs) and  $m = 10$ ,  $\beta = -0.5$  (purple square).

---

**Example: Chebyshev polynomials**

Let us consider the measure

$$d\sigma(\theta) = |e^{i\theta} - 1|^2 \frac{d\theta}{2\pi}, \quad (4.27)$$

and its corresponding sequence of monic orthogonal polynomials, given by [178]

$$\Phi_n(z) = \frac{1}{n+1} \sum_{k=0}^n (k+1)z^k, \quad n \geq 0. \quad (4.28)$$

We introduce the perturbation defined in this section and obtain

$$d\tilde{\sigma}(\theta) = d\sigma(\theta) + m \frac{d\theta}{2\pi}. \quad (4.29)$$

We now proceed to get an explicit expression for the sequence of monic polynomials orthogonal with respect to  $\tilde{\sigma}$ . Notice that

$$T_j(\Phi_n(z); 0) = \frac{1}{n+1} \sum_{k=0}^j (k+1)z^k, \quad 0 \leq j \leq n-1,$$

and, as a consequence,

$$T_j^*(\Phi_n(z); 0) = \frac{1}{n+1} \sum_{k=0}^j (k+1)z^{j-k}, \quad 0 \leq j \leq n-1.$$

On the other hand, since

$$\Phi_n^*(z) = \frac{1}{n+1} \sum_{k=0}^n (k+1)z^{n-k}, \quad n \geq 0,$$

we get

$$T_j(\Phi_n^*(z); 0) = \frac{1}{n+1} \sum_{k=0}^j (n+1-k)z^k, \quad 1 \leq j \leq n-1,$$

and, thus,

$$T_j^*(\Phi_n^*(z); 0) = \frac{1}{n+1} \sum_{k=0}^j (n+1-k)z^{j-k}, \quad 1 \leq j \leq n-1.$$

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES




Figure 4.1: Verblunsky coefficients for Bernstein-Szegő and Chebyshev polynomials

Therefore,  $\mathbf{T}(\Phi_n(0); 0) = \frac{1}{n+1} [1, 2, \dots, n]^T$  and  $\mathbf{T}(\Phi_n^*(0); 0) = \frac{1}{n+1} [n+1, n, n-1, \dots, 2]^T$ . Furthermore, since

$$\phi_n(z) = \sqrt{\frac{2}{(n+1)(n+2)}} \sum_{k=0}^n (k+1)z^k, \quad n \geq 0, \quad (4.30)$$

are the orthonormal polynomials with respect to  $\sigma$ , we obtain

$$\mathbf{P}_{n-1} = \mathbf{D}_n^{-1} \mathbf{A}_n \mathbf{\Lambda}_n \quad (4.31)$$

where

$$\mathbf{A}_n = \begin{bmatrix} 1 & 1 & \cdots & 1 \\ 0 & 1 & \ddots & \vdots \\ \vdots & \ddots & 1 & 1 \\ 0 & \cdots & 0 & 1 \end{bmatrix}, \quad \mathbf{\Lambda}_n = \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & p_1 & \ddots & \vdots \\ \vdots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & p_{n-1} \end{bmatrix},$$

and  $p_j = \sqrt{\frac{2}{(j+1)(j+2)}}$ . Finally, we have  $\mathbf{\Phi}_n(0) = \frac{1}{n+1} [1, 2!, \dots, n!]^T$ , and we can obtain  $\Psi_n(z)$  explicitly from (4.23). Figure 4.1(b) shows the behavior of the corresponding Verblunsky coefficients for different values of  $m$ , namely  $m = 10$  (blue discs) and  $m = 100$  (purple square).

This example can be generalized as follows. Let  $\sigma$  be an absolutely continuous measure whose Radon-Nikodym derivative with respect to the Lebesgue measure is  $\sigma' = |z - \alpha|^2$ ,  $z = e^{i\theta}$ , i.e., a positive trigonometric polynomial of degree 1. Applying the transformation introduced in this section, with  $m \in \mathbb{R}_+$ , and assuming  $\alpha \in \mathbb{C}$ , we obtain

---


$$m + |z - \alpha|^2 = m + 1 - \alpha z^{-1} - \bar{\alpha} z + |\alpha|^2, \quad z = e^{i\theta}, \quad (4.32)$$

i.e., another positive trigonometric polynomial that can be represented by  $|\delta z - \gamma|^2$ , where  $\delta \in \mathbb{R}, \gamma \in \mathbb{C}$ . Indeed, as

$$|\delta z - \gamma|^2 = \delta^2 - \delta \gamma z^{-1} - \delta \bar{\gamma} z + |\gamma|^2, \quad z = e^{i\theta},$$

the comparison of the coefficients with (4.32) yields  $1 + |\alpha|^2 + m = \delta^2 + |\gamma|^2$  and  $\alpha = \delta \gamma$ . Thus,

$$1 + |\alpha|^2 + m = \frac{|\alpha|^2}{\delta^2} + \delta^2,$$

so we can get  $\delta$  and  $\gamma$  in terms of  $m$  and  $\alpha$ . In other words, in this case we can express the addition to a Chebyshev measure of a Lebesgue measure (multiplied by a constant  $m$ ) as (4.27).

## 4.2.2 General perturbation of a Toeplitz matrix

In this subsection we generalize the previous perturbation, adding a mass  $m$  to any sub-diagonal of the Toeplitz matrix. Let  $\mathcal{L}_j$  be a linear functional such that its associated bilinear functional satisfies

$$\langle f, g \rangle_{\mathcal{L}_j} = \langle f, g \rangle_{\mathcal{L}} + m \int_{\mathbb{T}} z^j f(z) \overline{g(z)} \frac{dz}{2\pi i z} + \bar{m} \int_{\mathbb{T}} z^{-j} f(z) \overline{g(z)} \frac{dz}{2\pi i z}, \quad m \in \mathbb{C}, \quad j \geq 0. \quad (4.33)$$

Assume  $\mathcal{L}$  is a positive definite linear functional. Then, in terms of the corresponding measures, the above transformation can be expressed as

$$d\bar{\sigma}(z) = d\sigma(z) + 2\Re(mz^j) \frac{dz}{2\pi i z}. \quad (4.34)$$

From (4.33), if  $F_j$  is the  $C$ -function associated with  $\mathcal{L}_j$ , then

$$F_j(z) = F(z) + 2mz^j, \quad (4.35)$$

i.e., a special case of general linear spectral transformation of  $F$ . The infinite Toeplitz matrix  $\tilde{\mathbf{T}}$  associated with  $\mathcal{L}_j$ , is

$$\tilde{\mathbf{T}} = \mathbf{T} + m\mathbf{Z}^j + \bar{m}(\mathbf{Z}^T)^j,$$

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

---

where  $\mathbf{Z}$  is the shift matrix with ones on the first lower-diagonal and zeros on the remaining entries, and  $\mathbf{Z}^T$  is its transpose. Equivalently,

$$\tilde{\mathbf{T}} = \mathbf{T} + \begin{bmatrix} 0 & \cdots & \bar{m} & 0 & \cdots \\ \vdots & 0 & \cdots & \bar{m} & \cdots \\ \mathbf{m} & \vdots & \ddots & \vdots & \ddots \\ 0 & \mathbf{m} & \cdots & 0 & \cdots \\ \vdots & \vdots & \ddots & \vdots & \ddots \end{bmatrix}.$$

Assume that  $\mathcal{L}$  is a positive definite linear functional, and denote by  $\{\Phi_n\}_{n \geq 0}$  its corresponding sequence of monic orthogonal polynomials. We now proceed to determine necessary and sufficient conditions for  $\mathcal{L}_j$  to be a quasi-definite functional, as well as the relation between  $\{\Phi_n\}_{n \geq 0}$  and  $\{\Psi_n\}_{n \geq 0}$ , the sequence of monic orthogonal polynomials with respect to  $\mathcal{L}_j$ .

**Theorem 4.2.3.** *Let  $\mathcal{L}$  be a positive definite moment linear functional and  $\{\Phi_n\}_{n \geq 0}$  its corresponding sequence of monic orthogonal polynomials. Then, the following statements are equivalent:*

- i)  $\mathcal{L}_j$  is a quasi-definite linear functional.
- ii) The matrix  $\mathbf{I}_n + \mathbf{S}_n$  is non-singular, and

$$\tilde{\mathbf{k}}_n = \mathbf{k}_n + \mathbf{W}_n^T(0)(\mathbf{I}_n + \mathbf{S}_n)^{-1} \overline{\mathbf{Y}_n(0)} + \bar{m} \frac{\overline{\Phi_n^{(n-j)}(0)}}{(n-j)!} \neq 0, \quad n \geq 1, \quad (4.36)$$

with

$$\mathbf{Y}_n(0) = \left[ \bar{m} \frac{\Phi_n^{(j)}(0)}{j!}, \dots, \bar{m} \frac{\Phi_n^{(2j)}(0)}{(2j)!} + m \frac{\Phi_n^{(0)}(0)}{(0)!}, \dots, \bar{m} \frac{\Phi_n^{(n)}(0)}{(n)!} + m \frac{\Phi_n^{(n-2j)}(0)}{(n-2j)!}, \dots, m \frac{\Phi_n^{(n-j-1)}(0)}{(n-j-1)!} \right]^T,$$

$$\mathbf{W}_n(0) = [\Phi_n(0) - \bar{m}n! \mathbf{C}_{(0,n-1;n)}], \quad \Phi_n(0) = [\Phi_n(0), \Phi_n'(0), \dots, \Phi_n^{(n-1)}(0)]^T,$$

$$\mathbf{S}_n = \left[ \begin{array}{c|c|c} m\mathbf{A}_{(0,j-1;0,j-1)} & \mathbf{B}_{(0,j-1;j,n-j-1)} & \bar{m}\mathbf{C}_{(0,j-1;n-j,n-1)} \\ \hline m\mathbf{A}_{(j,n-j-1;0,j-1)} & \mathbf{B}_{(j,n-j-1;j,n-j-1)} & \bar{m}\mathbf{C}_{(j,n-j-1;n-j,n-1)} \\ \hline m\mathbf{A}_{(n-j,n-1;0,j-1)} & \mathbf{B}_{(n-j,n-1;j,n-j-1)} & \bar{m}\mathbf{C}_{(n-j,n-1;n-j,n-1)} \end{array} \right],$$



where  $\mathbf{A}$ ,  $\mathbf{B}$ , and  $\mathbf{C}$  are matrices whose elements are given by

$$a_{s,l} = \frac{K_{n-1}^{(s,l+j)}(0,0)}{(l)!(l+j)!}, \quad b_{s,l} = \mathbf{m} \frac{K_{n-1}^{(s,l+j)}(0,0)}{(l)!(l+j)!} + \overline{\mathbf{m}} \frac{K_{n-1}^{(s,l-j)}(0,0)}{(l)!(l-j)!}, \quad c_{s,l} = \frac{K_{n-1}^{(s,l-j)}(0,0)}{(l)!(l-j)!}.$$

Moreover,  $\{\Psi_n\}_{n \geq 0}$ , the corresponding sequence of monic orthogonal polynomials with respect to  $\mathcal{L}_j$ , is given by

$$\Psi_n(z) = A(z;n)\Phi_n(z) + B(z;n)\Phi_n^*(z), \quad n \geq 1, \quad (4.37)$$

with

$$\begin{aligned} A(z;n) &= 1 + \frac{1}{\mathbf{k}_n} \mathbf{W}_n^T(0)(\mathbf{I}_n + \mathbf{S}_n)^{-1} \mathbf{D}_n \mathcal{T}(\Phi_n(z); 0) + \overline{\mathbf{m}} \frac{T_{n-j}^*(\Phi_n(z); 0)}{\mathbf{k}_n}, \\ B(z;n) &= -\frac{1}{\mathbf{k}_n} \mathbf{W}_n^T(0)(\mathbf{I}_n + \mathbf{S}_n)^{-1} \mathbf{D}_n \mathcal{T}(\Phi_n^*(z); 0) - \overline{\mathbf{m}} \frac{T_{n-j}^*(\Phi_n^*(z); 0)}{\mathbf{k}_n}, \\ \mathcal{T}(\Phi_n(z); 0) &= \left[ \mathbf{m} T_j^*(\Phi_n(z); 0), \dots, \mathbf{m} T_{2j}^*(\Phi_n(z); 0) + \overline{\mathbf{m}} T_0^*(\Phi_n(z); 0), \dots, \right. \\ &\quad \left. \mathbf{m} T_{n-1}^*(\Phi_n(z); 0) + \overline{\mathbf{m}} T_{n-2j-1}^*(\Phi_n(z); 0), \dots, \overline{\mathbf{m}} T_{n-j-1}^*(\Phi_n(z); 0) \right]^T. \end{aligned}$$

*Proof.* Let us write  $\Psi_n(z) = \Phi_n(z) + \sum_{k=0}^{n-1} \lambda_{n,k} \Phi_k(z)$ , where, for  $0 \leq k \leq n-1$ ,

$$\lambda_{n,k} = -\frac{\mathbf{m}}{\mathbf{k}_k} \int_{\mathbb{T}} y^j \Psi_n(y) \overline{\Phi_k(y)} \frac{dy}{2\pi i y} - \frac{\overline{\mathbf{m}}}{\mathbf{k}_k} \int_{\mathbb{T}} y^{-j} \Psi_n(y) \overline{\Phi_k(y)} \frac{dy}{2\pi i y}.$$

Therefore,

$$\Psi_n(z) = \Phi_n(z) - \mathbf{m} \int_{\mathbb{T}} y^j \Psi_n(y) K_{n-1}(z, y) \frac{dy}{2\pi i y} - \overline{\mathbf{m}} \int_{\mathbb{T}} y^{-j} \Psi_n(y) K_{n-1}(z, y) \frac{dy}{2\pi i y}.$$

Taking into account that

$$y^j \Psi_n(y) = \sum_{l=0}^n \frac{\Psi_n^{(l)}(0)}{l!} y^{l+j} = \sum_{l=j}^{n+j} \frac{\Psi_n^{(l-j)}(0)}{(l-j)!} y^l,$$

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

and, for  $|y| = 1$ ,  $K_{n-1}(z, y) = \sum_{l=0}^{n-1} \frac{K_{n-1}^{(0,l)}(z, 0)}{l!} \frac{1}{y^l}$ , we obtain

$$\int_{\mathbb{T}} y^j \Psi_n(y) K_{n-1}(z, y) \frac{dy}{2\pi i y} = \sum_{l=j}^{n-1} \frac{\Psi_n^{(l-j)}(0)}{(l-j)!} \frac{K_{n-1}^{(0,l)}(z, 0)}{(l)!} = \sum_{l=0}^{n-j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l+j)}(z, 0)}{(l+j)!}.$$

In an analog way,

$$\int_{\mathbb{T}} y^{-j} \Psi_n(y) K_{n-1}(z, y) \frac{dy}{2\pi i y} = \sum_{l=0}^{n-j} \frac{\Psi_n^{(l+j)}(0)}{(l+j)!} \frac{K_{n-1}^{(0,l)}(z, 0)}{(l)!} = \sum_{l=j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l-j)}(z, 0)}{(l-j)!}.$$

Thus, we get

$$\Psi_n(z) = \Phi_n(z) - \mathbf{m} \sum_{l=0}^{n-j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l+j)}(z, 0)}{(l+j)!} - \overline{\mathbf{m}} \sum_{l=j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l-j)}(z, 0)}{(l-j)!}, \quad (4.38)$$

or, equivalently,

$$\begin{aligned} \Psi_n(z) = \Phi_n(z) - \mathbf{m} \sum_{l=0}^{j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l+j)}(z, 0)}{(l+j)!} - \overline{\mathbf{m}} \sum_{l=n-j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l-j)}(z, 0)}{(l-j)!}, \\ - \sum_{l=j}^{n-j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \left( \mathbf{m} \frac{K_{n-1}^{(0,l+j)}(z, 0)}{(l+j)!} + \overline{\mathbf{m}} \frac{K_{n-1}^{(0,l-j)}(z, 0)}{(l-j)!} \right). \end{aligned} \quad (4.39)$$

In particular, for  $0 \leq s \leq n$ ,

$$\begin{aligned} \Psi_n^{(s)}(0) = \Phi_n^{(s)}(0) - \mathbf{m} \sum_{l=0}^{j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(s,l+j)}(0, 0)}{(l+j)!} - \overline{\mathbf{m}} \sum_{l=n-j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(s,l-j)}(0, 0)}{(l-j)!}, \\ - \sum_{l=j}^{n-j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \left( \mathbf{m} \frac{K_{n-1}^{(s,l+j)}(0, 0)}{(l+j)!} + \overline{\mathbf{m}} \frac{K_{n-1}^{(s,l-j)}(0, 0)}{(l-j)!} \right), \end{aligned}$$

i.e., we obtain a system of  $n+1$  linear equations and  $n+1$  unknowns as follows

$$\Psi_n^{(s)}(0) = \Phi_n^{(s)}(0) - \mathbf{m} \sum_{l=0}^{j-1} a_{s,l} \Psi_n^{(l)}(0) - \sum_{l=j}^{n-j-1} b_{s,l} \Psi_n^{(l)}(0) - \overline{\mathbf{m}} \sum_{l=n-j}^n c_{s,l} \Psi_n^{(l)}(0).$$

Thus, if  $\mathbf{M}_{(s_1, s_2; l_1, l_2)} = [m_{s,i}]_{s_1 \leq s \leq s_2; l_1 \leq l \leq l_2}$ , then

$$(\mathbf{I}_{n+1} + \mathbf{S}_{n+1}) \begin{bmatrix} \Psi_n^{(0)}(0) \\ \vdots \\ \Psi_n^{(n)}(0) \end{bmatrix} = \begin{bmatrix} \Phi_n^{(0)}(0) \\ \vdots \\ \Phi_n^{(n)}(0) \end{bmatrix}.$$

Notice that the entries in the last row of the above matrix vanish, which is consistent with the fact that  $\Psi_n^{(n)}(0) = \Phi_n^{(n)}(0) = n!$ . Therefore, if we denote by

$$\Psi_n(0) = [\Psi_n(0), \Psi_n'(0), \dots, \Psi_n^{(n-1)}(0)]^T,$$

then the above  $(n+1) \times (n+1)$  linear system can be reduced to a  $n \times n$  linear system as follows

$$(\mathbf{I}_n + \mathbf{S}_n)\Psi_n(0) = \mathbf{W}_n(0). \quad (4.40)$$

Since  $\mathcal{L}_j$  is a quasi-definite linear functional, there exists a unique family of monic polynomials orthogonal with respect to  $\mathcal{L}_j$ . Therefore, the matrix  $\mathbf{I}_n + \mathbf{S}_n$  is non-singular, according to the existence and uniqueness of the solution of the above linear system. As a consequence, if

$$\mathbb{K}_{n-1}(z, 0) = \begin{bmatrix} m \frac{K_{n-1}^{(0,j)}(z, 0)}{j!} \\ \vdots \\ m \frac{K_{n-1}^{(0,2j)}(z, 0)}{(2j)!} + \overline{m} \frac{K_{n-1}^{(0,0)}(z, 0)}{(0)!} \\ \vdots \\ m \frac{K_{n-1}^{(0,n-1)}(z, 0)}{(n-1)!} + \overline{m} \frac{K_{n-1}^{(0,n-2j-1)}(z, 0)}{(n-2j-1)!} \\ \vdots \\ \overline{m} \frac{K_{n-1}^{(0,n-j-1)}(z, 0)}{(n-j-1)!} \end{bmatrix}, \quad (4.41)$$

then (4.39) becomes

$$\Psi_n(z) = \Phi_n(z) - \Psi_n^T(0)\mathbf{D}_n\mathbb{K}_{n-1}(z, 0) - \overline{m} \frac{K_{n-1}^{(0,n-j)}(z, 0)}{(n-j)!}.$$

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

Thus, from (4.40) and Theorem 4.2.2, (4.37) holds. Furthermore,

$$\begin{aligned}
 0 &\neq \langle \Psi_n, \Phi_n \rangle_{\mathcal{L}_j} \\
 &= \mathbf{k}_n + \mathbf{m} \sum_{l=0}^{n-j} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{\overline{\Phi_n^{(l+j)}(0)}}{(l+j)!} + \overline{\mathbf{m}} \sum_{l=j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{\overline{\Phi_n^{(l-j)}(0)}}{(l-j)!} \\
 &= \mathbf{k}_n + \Psi_n^T(0) \overline{\mathbf{Y}_n(0)} + \overline{\mathbf{m}} \frac{\overline{\Phi_n^{(n-j)}(0)}}{(n-j)!},
 \end{aligned}$$

so (4.36) follows. For the converse, assume  $\mathbf{I}_n + \mathbf{S}_n$  is non-singular for every  $n \geq 1$  and define  $\{\Psi_n\}_{n \geq 0}$  as in (4.37). We show that  $\{\Psi_n\}_{n \geq 0}$  is orthogonal with respect to  $\mathcal{L}_j$ . Indeed, for  $0 \leq k \leq n-1$  and taking into account (4.38), we get

$$\begin{aligned}
 \langle \Psi_n, \Phi_k \rangle_{\mathcal{L}_j} &= \langle \Phi_n, \Phi_k \rangle_{\mathcal{L}} - \mathbf{m} \left\langle \sum_{l=0}^{n-j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l+j)}(z,0)}{(l+j)!}, \Phi_k(z) \right\rangle_{\mathcal{L}} \\
 &\quad - \overline{\mathbf{m}} \left\langle \sum_{l=j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{K_{n-1}^{(0,l-j)}(z,0)}{(l-j)!}, \Phi_k(z) \right\rangle_{\mathcal{L}} \\
 &\quad + \mathbf{m} \sum_{l=0}^{n-j-1} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{\overline{\Phi_k^{(l+j)}(0)}}{(l+j)!} + \overline{\mathbf{m}} \sum_{l=j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{\overline{\Phi_k^{(l-j)}(0)}}{(l-j)!} = 0.
 \end{aligned}$$

On the other hand,

$$\begin{aligned}
 \widetilde{\mathbf{k}}_n &= \langle \Psi_n, \Phi_n \rangle_{\mathcal{L}_j} \\
 &= \mathbf{k}_n + \mathbf{m} \sum_{l=0}^{n-j} \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{\overline{\Phi_n^{(l+j)}(0)}}{(l+j)!} + \overline{\mathbf{m}} \sum_{l=j}^n \frac{\Psi_n^{(l)}(0)}{(l)!} \frac{\overline{\Phi_n^{(l-j)}(0)}}{(l-j)!} \\
 &= \mathbf{k}_n + \Psi_n^T(0) \overline{\mathbf{Y}_n(0)} + \overline{\mathbf{m}} \frac{\overline{\Phi_n^{(n-j)}(0)}}{(n-j)!} \neq 0,
 \end{aligned}$$

since (4.36) is assumed. Thus, we conclude that  $\mathcal{L}_j$  is quasi-definite.  $\square$

**Remark 4.2.1.** The case  $\mathcal{L}_0$  ( $j = 0$ ) reduces to the linear functional analyzed in the previous subsection, with mass  $\mathfrak{K}\mathbf{m}$ . In such a case,  $\widetilde{\mathbf{k}}_0 = \mathbf{k}_0 + \mathfrak{K}\mathbf{m}$ . On the other hand, for  $j \geq 1$ , it follows from (4.33) that  $\widetilde{\mathbf{k}}_l = \mathbf{k}_l$  for  $0 \leq l \leq j-1$ . In other words, we only need (4.36) for  $n \geq j$ . Notice that for a given  $j$ , the polynomials of degree  $n < j$  remain unchanged. In such a case, (4.36) and (4.37) still hold, with the convention that the

---

negative derivatives are zero.

Finally, applying the Szegő transformation to (4.35), we get

$$\widetilde{S}(x) = S(x) + 2\mathbf{m} \frac{(x - \sqrt{x^2 - 1})^j}{\sqrt{x^2 - 1}},$$

and thus  $\widetilde{S}$ , the Stieltjes function for the corresponding perturbed measure on the real line, can not be expressed as a linear spectral transform of  $S$ , since square roots appear for any value of  $j$ . Therefore, we conclude that a perturbation on the moments of a measure supported on  $\mathbb{T}$  does not yield a general linear spectral transformation of the corresponding Stieltjes function. Conversely, if we consider a similar perturbation of the moments of a measure on the real line, then

$$\widetilde{S}(x) = S(x) + \frac{\mathbf{m}}{x^{j+1}}.$$

Applying the Szegő transformation,

$$F_0(z) = F(z) + \mathbf{m} \frac{1 - z^2}{2z(x)^{j+1}} = F(z) + 2^j \mathbf{m} \frac{(1 - z^2)z^j}{(z^2 + 1)^{j+1}},$$

which is a general linear spectral transformation of  $F$ . In the special case when  $j = 0$ ,

$$F_0(z) = F(z) - \mathbf{m} \frac{z^2 - 1}{z^2 + 1}.$$

As a conclusion, the study of general linear spectral transformations on the unit circle is far more complicated than the real line case.

### Example: Lebesgue polynomials

We present an example of the previous transformation when  $\sigma$  is the normalized Lebesgue measure and  $j = 1$ , i.e., the transformation

$$\langle f, g \rangle_{\mathcal{L}_1} = \int_{\mathbb{T}} f(z) \overline{g(z)} \frac{dz}{2\pi iz} + \mathbf{m} \int_{\mathbb{T}} z f(z) \overline{g(z)} \frac{dz}{2\pi iz} + \overline{\mathbf{m}} \int_{\mathbb{T}} z f(z) \overline{g(z)} \frac{dz}{2\pi iz}, \quad \mathbf{m} \in \mathbb{C}, \quad (4.42)$$

where  $\mathbf{m} \in \mathbb{C}$ . Our purpose is to obtain necessary and sufficient conditions for  $\mathcal{L}_1$  to be a positive definite (quasi-definite) functional. Consequently, we deduce its corresponding family of orthogonal polynomials, as well as the sequence of Verblunsky

#### 4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES

---

coefficients. Notice that in this case,  $\Phi_n(z) = z^n$ ,  $n \geq 0$ , as well as  $\mathbf{k}_n = 1$ ,  $n \geq 0$ . Thus,

$$K_{n-1}^{(0,l)}(z, 0) = l!z^l, \quad 0 \leq l \leq n-1.$$

So,  $\mathbb{K}_{n-1}(z, 0) = [\mathbf{m}z, \mathbf{m}z^2 + \bar{\mathbf{m}}z^0, \dots, \mathbf{m}z^{n-1} + \bar{\mathbf{m}}z^{n-3}, \bar{\mathbf{m}}z^{n-2}]^T$  and  $\Phi_n(0) = \mathbf{0}^T$ ,  $n \geq 1$ . On the other hand,

$$K_{n-1}^{(s,l)}(0, 0) = \begin{cases} s!l! & \text{if } s = l, \\ 0 & \text{otherwise,} \end{cases}$$

and, therefore,

$$\begin{aligned} a_{s,l} &= \frac{K_{n-1}^{(s,l+1)}(0, 0)}{(l)!(l+1)!} = \delta_{s,s-1}, \\ b_{s,l} &= \mathbf{m} \frac{K_{n-1}^{(s,l+1)}(0, 0)}{(l)!(l+1)!} + \bar{\mathbf{m}} \frac{K_{n-1}^{(s,l-1)}(0, 0)}{(l)!(l-1)!} = \mathbf{m}\delta_{s,s-1} + \bar{\mathbf{m}}\delta_{s,s+1}, \\ c_{s,l} &= \frac{K_{n-1}^{(s,l-1)}(0, 0)}{(l)!(l-1)!} = \delta_{s,s+1}, \end{aligned}$$

where  $\delta_{s,l}$  is the Kronecker's delta. Thus,

$$\mathbf{I}_n + \mathbf{S}_n = \begin{bmatrix} 1 & \bar{\mathbf{m}} & & & \\ \mathbf{m} & 1 & \ddots & & \\ & \ddots & \ddots & \bar{\mathbf{m}} & \\ & & \mathbf{m} & 1 & \bar{\mathbf{m}} \\ & & & \mathbf{m} & 1 \end{bmatrix}, \quad n \geq 2.$$

Notice that for  $n \geq 2$ ,  $\mathbf{I}_n + \mathbf{S}_n$  is  $\tilde{\mathbf{T}}_{n-1}$ , the  $n \times n$  Toeplitz matrix associated with  $\mathcal{L}_1$ . We thus need to establish the conditions on  $m$  for  $\tilde{\mathbf{T}}_{n-1}$  be non-singular. Since  $\tilde{\mathbf{T}}_{n-1}$  is hermitian, their eigenvalues  $\{\lambda_k\}_{k=1}^n$  are real numbers. Moreover,  $\tilde{\mathbf{T}}_{n-1}$  is quasi-definite if and only if  $\lambda_k \neq 0$ , for every  $1 \leq k \leq n$  [92]. From Theorem 2.4 in [18], the eigenvalues of  $\tilde{\mathbf{T}}_{n-1}$  are

$$\lambda_k = 1 + 2|\mathbf{m}| \cos \frac{\pi k}{n+1}, \quad k = 1, \dots, n.$$

Thus,  $\mathcal{L}_1$  is a quasi-definite linear functional if and only if

$$|\mathbf{m}| \neq -\left(2 \cos \frac{\pi k}{n+1}\right)^{-1}, \quad k = 1, \dots, n, \quad n \geq 1,$$

or, equivalently,

$$\frac{\pi k}{\cos^{-1}\left(-\frac{1}{2|\mathbf{m}|}\right)} \notin \mathbb{N}. \quad (4.43)$$

Assuming that (4.43) holds,  $\{\Psi_n\}_{n \geq 0}$  can be obtained using (4.37), since all elements are known. Since  $\mathbb{K}_{n-1}(0,0) = (0, \overline{\mathbf{m}}, 0, \dots, 0)^T$  and  $\mathbf{W}_n = [0, \dots, 0, -\overline{\mathbf{m}}(n-1)!]$ , the sequence of Verblunsky coefficients can be computed using (A.1.2). It is not difficult to see that

$$\Psi_n(0) = -\overline{\mathbf{m}}^2(n-1)!\ell_{n,2},$$

where  $\ell_{i,j} = (\mathbf{I}_n + \mathbf{S}_n)_{i,j}^{-1}$ . An explicit expression for  $\ell_{n,2}$  can be obtained using the method described in [192]. Indeed,

$$\ell_{n,2} = \frac{(-1)^{n+2} \mathbf{m}^{n-2}}{\theta_n}$$

where  $\theta_n$  is the solution of the recurrence relation

$$\theta_i = \theta_{i-1} - |\mathbf{m}|^2 \theta_{i-2}, \quad i = 2, \dots, n,$$

with initial conditions  $\theta_0 = \theta_1 = 1$ . Thus,

$$\theta_n = \left( 2^{-(n+1)} + \frac{2^{-(n+1)}}{\sqrt{1-|\mathbf{m}|^2}} \right) \left( (1 - \sqrt{1-|\mathbf{m}|^2})^n + (1 + \sqrt{1-|\mathbf{m}|^2})^n \right),$$

and therefore we get

$$\Psi_n(0) = \frac{(-1)^{n+3} \mathbf{m}^{n-2} \overline{\mathbf{m}}(n-1)!}{\left( 2^{-(n+1)} + \frac{2^{-(n+1)}}{\sqrt{1-|\mathbf{m}|^2}} \right) \left( (1 - \sqrt{1-|\mathbf{m}|^2})^n + (1 + \sqrt{1-|\mathbf{m}|^2})^n \right)}. \quad (4.44)$$

From (4.44) notice that  $|\Psi_n(0)|$  grows as  $n$  increases, and it grows faster for values  $|\mathbf{m}| > 1$ . Using small values of  $\mathbf{m}$ , the first Verblunsky coefficients are small (close to zero), but then begin to grow as  $n$  increases.

#### **4. SPECTRAL TRANSFORMATIONS OF MOMENT MATRICES**

---



## Chapter 5

# Spectral transformations associated with mass points

... then we get a  $\mu$  (measure) for which mass points do not attract zeros ...

— E. B. Saff and V. Totik. [171]

In Chapter 4 we consider the addition of a Lebesgue measure to the hermitian linear functional  $\mathcal{L}$  defined in (2.23). In other words, we perturb the main diagonal of the Toeplitz matrix (2.24). The generalization of the previous perturbations to affect any sub-diagonal of the Toeplitz matrix (2.24) is also considered. As we see, they are general linear spectral transformations of the corresponding  $C$ -function (2.39).

One of the goals of this chapter is to show two new examples of linear spectral transformations associated with the first derivative of a complex Dirac's linear functional. The first one appears when the support of the Dirac's linear functional is a point on the unit circle. The second one corresponds to a Dirac's linear functional supported in two symmetric points with respect to the unit circle. Necessary and sufficient conditions for the quasi-definiteness of the new linear functional are obtained. Outer relative asymptotics for the new sequence of monic orthogonal polynomials in terms of the original ones are studied. We also prove that this spectral transformation can be decomposed as an iteration of particular cases of the canonical spectral transformations (2.44) and (2.45).

The last part of this chapter is devoted to the study of a relevant family of orthogonal

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

polynomials associated with perturbations of the original orthogonality measure by means of mass points: discrete Sobolev orthogonal polynomials. We compare the discrete Sobolev orthogonal polynomials with the original ones. Finally, we analyze the behavior of their zeros and provide some numerical examples to illustrate it. An analogous inner product for measures supported on the real line is study in Appendix A.

### 5.1 Adding the derivative of a Dirac's delta

Let  $\mathcal{L}$  be a hermitian linear functional given by (2.23). Its derivative  $D\mathcal{L}$  [189] is defined by

$$\langle D\mathcal{L}, f \rangle = -i\langle \mathcal{L}, zf'(z) \rangle, \quad f \in \Lambda.$$

In this section we first consider a perturbation of a linear functional  $\mathcal{L}$  by the addition of a derivative of a Dirac's delta, i.e.,

$$\langle \mathcal{L}_1, f \rangle = \langle \mathcal{L}, f \rangle + m\langle D\delta_\alpha, f \rangle, \quad m \in \mathbb{R}, \quad |\alpha| = 1. \quad (5.1)$$

Let  $\mathcal{L}_U$  be a linear functional such that

$$\langle \mathcal{L}_U, f \rangle = \langle \mathcal{L}, f \rangle + mf(\alpha), \quad m \in \mathbb{R}, \quad |\alpha| = 1.$$

We say that  $\mathcal{L}_U$  is the Uvarov spectral transformation of the linear functional  $\mathcal{L}$  [54]. The connection between the corresponding sequences of monic orthogonal polynomials as well as the associated GGT matrices using LU and QR factorization has been studied in [54]. The iteration of Uvarov transformations has been considered in [78; 117]; see also Appendix B. Asymptotic properties for the corresponding sequences of orthogonal polynomials have been studied in [201]. Notice that the addition of a Dirac's delta derivative (on a point of the unit circle) to a linear functional can be considered as the limit case of two Uvarov spectral transformations with equal masses and opposite sign, located on two nearby points on the unit circle  $\alpha_1 = e^{i\theta_1}$  and  $\alpha_2 = e^{i\theta_2}$ ,  $0 \leq \theta_1, \theta_2 \leq 2\pi$ , when  $\theta_1 \rightarrow \theta_2$ , but the difficulties to deal with them yield a different approach.

---

### 5.1.1 Mass point on the unit circle

In terms of the associated bilinear functional (5.1) becomes

$$\langle f, g \rangle_{\mathcal{L}_1} = \langle f, g \rangle_{\mathcal{L}} - im \left( \alpha f'(\alpha) \overline{g(\alpha)} - \bar{\alpha} f(\alpha) \overline{g'(\alpha)} \right). \quad (5.2)$$

In the next theorem we obtain necessary and sufficient conditions for  $\mathcal{L}_1$  to be a quasi-definite linear functional, as well as an expression for its corresponding family of orthogonal polynomials.

**Theorem 5.1.1.** *Let us assume  $\mathcal{L}$  is a quasi-definite linear functional and denote by  $\{\Phi_n\}_{n \geq 0}$  its corresponding sequence of monic orthogonal polynomials. Let consider  $\mathcal{L}_1$  as in (5.2). Then, the following statements are equivalent:*

- i)  $\mathcal{L}_1$  is quasi-definite.
- ii) The matrix  $\mathbf{D}(\alpha) + m\mathbb{K}_{n-1}(\alpha, \alpha)$ , with

$$\mathbb{K}_{n-1}(\alpha, \alpha) = \begin{bmatrix} K_{n-1}(\alpha, \alpha) & K_{n-1}^{(0,1)}(\alpha, \alpha) \\ K_{n-1}^{(1,0)}(\alpha, \alpha) & K_{n-1}^{(1,1)}(\alpha, \alpha) \end{bmatrix}, \quad \mathbf{D}(\alpha) = \begin{bmatrix} 0 & -i\alpha \\ i\alpha^{-1} & 0 \end{bmatrix},$$

is non-singular, and

$$\mathbf{k}_n + m\Phi_n(\alpha)^H (\mathbf{D}(\alpha) + m\mathbb{K}_{n-1}(\alpha, \alpha))^{-1} \Phi_n(\alpha) \neq 0, \quad n \geq 1. \quad (5.3)$$

Furthermore, the sequence  $\{\Psi_n\}_{n \geq 0}$  of monic orthogonal polynomials associated with  $\mathcal{L}_1$  is given by

$$\Psi_n(z) = \Phi_n(z) - m \begin{bmatrix} K_{n-1}(z, \alpha) \\ K_{n-1}^{(0,1)}(z, \alpha) \end{bmatrix}^T (\mathbf{D}(\alpha) + m\mathbb{K}_{n-1}(\alpha, \alpha))^{-1} \Phi_n(\alpha), \quad (5.4)$$

where  $\Phi_n(\alpha) = [\Phi_n(\alpha), \Phi_n'(\alpha)]^T$ .

*Proof.* Assume  $\mathcal{L}_1$  is quasi-definite and denote by  $\{\Psi_n\}_{n \geq 0}$  its corresponding family of monic orthogonal polynomials. Let us consider the Fourier expansion

$$\Psi_n(z) = \Phi_n(z) + \sum_{k=0}^{n-1} \lambda_{n,k} \Phi_k(z),$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

where, for  $n \geq 1$ ,

$$\lambda_{n,k} = \frac{\langle \Psi_n(z), \Phi_k(z) \rangle_{\mathcal{L}}}{\mathbf{k}_k} = \frac{im \left( \alpha \Psi'_n(\alpha) \overline{\Phi_k(\alpha)} - \bar{\alpha} \Psi_n(\alpha) \overline{\Phi'_k(\alpha)} \right)}{\mathbf{k}_k}, \quad 0 \leq k \leq n-1.$$

Thus,

$$\begin{aligned} \Psi_n(z) &= \Phi_n(z) + \sum_{k=0}^{n-1} \frac{im \left( \alpha \Psi'_n(\alpha) \overline{\Phi_k(\alpha)} - \bar{\alpha} \Psi_n(\alpha) \overline{\Phi'_k(\alpha)} \right)}{\mathbf{k}_k} \Phi_k(z), \\ &= \Phi_n(z) + im \left( \alpha \Psi'_n(\alpha) K_{n-1}(z, \alpha) - \bar{\alpha} \Psi_n(\alpha) K_{n-1}^{(0,1)}(z, \alpha) \right). \end{aligned} \quad (5.5)$$

Taking the derivative with respect to  $z$  in the previous expression and evaluating at  $z = \alpha$ , we obtain the linear system

$$\Psi_n^{(i)}(\alpha) = \Phi_n^{(i)}(\alpha) + im \left( \alpha \Psi'_n(\alpha) K_{n-1}^{(i,0)}(\alpha, \alpha) - \bar{\alpha} \Psi_n(\alpha) K_{n-1}^{(i,1)}(\alpha, \alpha) \right), \quad i = 0, 1,$$

which yields

$$\begin{bmatrix} \Phi_n(\alpha) \\ \Phi'_n(\alpha) \end{bmatrix} = \begin{bmatrix} 1 + im\bar{\alpha} K_{n-1}^{(0,1)}(\alpha, \alpha) & -im\alpha K_{n-1}(\alpha, \alpha) \\ im\bar{\alpha} K_{n-1}^{(1,1)}(\alpha, \alpha) & 1 - im\alpha K_{n-1}^{(1,0)}(\alpha, \alpha) \end{bmatrix} \begin{bmatrix} \Psi_n(\alpha) \\ \Psi'_n(\alpha) \end{bmatrix},$$

and denoting  $\mathbf{Q} = [Q, Q']^T$ , we get

$$\Phi_n(\alpha) = (\mathbf{I}_2 + m\mathbb{K}_{n-1}(\alpha, \alpha)\mathbf{D}(\alpha)) \Psi_n(\alpha).$$

Thus, the necessary condition for regularity is that  $\mathbf{I}_2 + m\mathbb{K}_{n-1}(\alpha, \alpha)\mathbf{D}(\alpha)$  must be non-singular. Taking into account  $\mathbf{D}^{-1}(\alpha) = \mathbf{D}(\alpha)$  we have the first part of our statement. Furthermore, from (5.5),

$$\begin{aligned} \Psi_n(z) &= \Phi_n(z) + m \left( K_{n-1}(z, \alpha), K_{n-1}^{(0,1)}(z, \alpha) \right) \begin{bmatrix} 0 & i\alpha \\ -i\bar{\alpha} & 0 \end{bmatrix} \begin{bmatrix} \Psi_n(\alpha) \\ \Psi'_n(\alpha) \end{bmatrix} \\ &= \Phi_n(z) - m \begin{bmatrix} K_{n-1}(z, \alpha) \\ K_{n-1}^{(0,1)}(z, \alpha) \end{bmatrix}^T (\mathbf{D}(\alpha) + m\mathbb{K}_{n-1}(\alpha, \alpha))^{-1} \Phi_n(\alpha). \end{aligned}$$

This yields (5.4). Conversely, if  $\{\Psi_n\}_{n \geq 0}$  is given by (5.5), then, for  $0 \leq k \leq n-1$ ,

$$\begin{aligned} \langle \Psi_n, \Psi_k \rangle_{\mathcal{L}_1} &= \left\langle \Phi_n(z) + im \left( \alpha \Psi'_n(\alpha) K_{n-1}(z, \alpha) - \bar{\alpha} \Psi_n(\alpha) K_{n-1}^{(0,1)}(z, \alpha) \right), \Psi_k(z) \right\rangle_{\mathcal{L}_1} \\ &= \left\langle \Phi_n(z) + im \left( \alpha \Psi'_n(\alpha) K_{n-1}(z, \alpha) - \bar{\alpha} \Psi_n(\alpha) K_{n-1}^{(0,1)}(z, \alpha) \right), \Psi_k(z) \right\rangle_{\mathcal{L}} \\ &\quad - im \left( \alpha \Psi'_n(\alpha) \overline{\Psi_k(\alpha)} - \bar{\alpha} \Psi_n(\alpha) \overline{\Psi'_k(\alpha)} \right) = 0. \end{aligned}$$

On the other hand, for  $n \geq 1$ ,

$$\begin{aligned} \widetilde{\mathbf{k}}_n &= \langle \Psi_n(z), \Psi_n(z) \rangle_{\mathcal{L}_1} = \langle \Psi_n(z), \Phi_n(z) \rangle_{\mathcal{L}_1} \\ &= \left\langle \Phi_n(z) + im \left( \alpha \Psi'_n(\alpha) K_{n-1}(z, \alpha) - \bar{\alpha} \Psi_n(\alpha) K_{n-1}^{(0,1)}(z, \alpha) \right), \Phi_n(z) \right\rangle_{\mathcal{L}} \\ &\quad - im \left( \alpha \Psi'_n(\alpha) \overline{\Phi_n(\alpha)} - \bar{\alpha} \Psi_n(\alpha) \overline{\Phi'_n(\alpha)} \right) \\ &= \mathbf{k}_n + m \Phi_n(\alpha)^H (\mathbf{D}(\alpha) + m \mathbb{K}_{n-1}(\alpha, \alpha))^{-1} \Phi_n(\alpha) \neq 0, \end{aligned}$$

where we are using the reproducing property (2.34). As a conclusion,  $\{\Psi_n\}_{n \geq 0}$  is the sequence of monic orthogonal polynomials with respect to  $\mathcal{L}_1$ .  $\square$

Using the Christoffel-Darboux formula (2.33), another way to express (5.4) is the following.

**Corollary 5.1.1.** *Let  $\{\Psi_n\}_{n \geq 0}$  be the sequence of monic orthogonal polynomials associated to  $\mathcal{L}_1$  defined as in (5.2). Then,*

$$(z - \alpha)^2 \Psi_n(z) = A(z, n, \alpha) \Phi_n(z) + B(z, n, \alpha) \Phi_n^*(z), \quad (5.6)$$

where  $A(z, n, \alpha)$  and  $B(z, n, \alpha)$  are polynomials of degree 2 and 1, respectively, in the variable  $z$ , given by

$$\begin{aligned} A(z, n, \alpha) &= (z - \alpha)^2 - \frac{m\alpha}{\mathbf{k}_n \Delta_{n-1}} \left( (Y_{1,1} \Phi_n(\alpha) + Y_{1,2} \Phi'_n(\alpha)) \overline{\Phi_n(\alpha)} (z - \alpha) \right. \\ &\quad \left. + (Y_{2,1} \Phi_n(\alpha) + Y_{2,2} \Phi'_n(\alpha)) (\Phi_n(\alpha) (z - \alpha) + \alpha \Phi_n(\alpha) z) \right), \\ B(z, n, \alpha) &= \frac{m\alpha}{\mathbf{k}_n \Delta_{n-1}} \left( (Y_{1,1} \Phi_n(\alpha) + Y_{1,2} \Phi'_n(\alpha)) \overline{\Phi_n^*(\alpha)} \right. \\ &\quad \left. + (Y_{2,1} \Phi_n(\alpha) + Y_{2,2} \Phi'_n(\alpha)) (\overline{\Phi_n^*(\alpha)} (z - \alpha) + \alpha \overline{\Phi_n^*(\alpha)} z) \right), \end{aligned}$$

where  $Y_{1,1} = m K_{n-1}^{(1,1)}(\alpha, \alpha)$ ,  $Y_{1,2} = im \alpha K_{n-1}^{(0,1)}(\alpha, \alpha)$ ,  $Y_{2,1} = -im \bar{\alpha} K_{n-1}^{(1,0)}(\alpha, \alpha)$ ,  $Y_{2,2} = m \alpha K_{n-1}(\alpha, \alpha)$ , and  $\Delta_{n-1}$  is the determinant of the matrix  $\mathbf{D}(\alpha) + im \mathbb{K}_{n-1}(\alpha, \alpha)$ .

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

### 5.1.1.1 Outer relative asymptotics

In this subsection we assume  $\mathcal{L}$  is a positive definite linear functional, with an associated positive Borel measure  $\sigma$ . We are interested in the asymptotic behavior of the orthogonal polynomials associated with the addition of the derivative of a Dirac's delta on the unit circle given in (5.6). We assume that  $\sigma$  is regular in the sense of Stahl and Totik [181], so that

$$\lim_{n \rightarrow \infty} \kappa_n^{1/n} = 1.$$

Regularity is a necessary and sufficient condition for the existence of  $n$ -th root asymptotics, i.e.,  $\lim_{n \rightarrow \infty} |\phi_n|^{1/n} < \infty$ . It is easy to see that the existence of the ratio asymptotics  $\lim_{n \rightarrow \infty} \phi_n / \phi_{n-1}$  implies the existence of the root asymptotics, and, in general, the converse is not true. Therefore, ratio asymptotics does not must hold for regular measures.

In particular, we study its ratio asymptotics with respect to  $\{\Phi_n\}_{n \geq 0}$ . First, we state some results that are useful in our study.

**Theorem 5.1.2.** [115] *Let  $\sigma$  be a regular finite positive Borel measure supported on  $(-\pi, \pi]$ . Let  $J \subset (-\pi, \pi)$  be a compact subset such that  $\sigma$  is absolutely continuous in an open set containing  $J$ . Assume that  $\sigma'$  is positive and continuous at each point of  $J$ . Let  $i, j$  be non-negative integers. Then, uniformly for  $\theta \in J$ ,  $z = e^{i\theta}$ ,*

$$\lim_{n \rightarrow \infty} \frac{z^{i-j} K_n^{(i,j)}(z, z)}{n^{i+j} K_n(z, z)} = \frac{1}{i + j + 1}.$$

**Lemma 5.1.1.** [85] *Let  $f, g$  be two polynomials in  $\mathbb{P}$  with degree at least  $j$ . Then*

$$\frac{f^{(j)}(z)}{g^{(j)}(z)} = \frac{g^{(j-1)}(z)}{g^{(j)}(z)} \left( \frac{f^{(j-1)}(z)}{g^{(j-1)}(z)} \right)' + \frac{f^{(j-1)}(z)}{g^{(j-1)}(z)}.$$

Using the previous lemma, the outer ratio asymptotics for the derivatives of orthonormal polynomials are deduced.

**Lemma 5.1.2.** *Assume  $\mathcal{L}$  is a positive definite linear functional, with an associated positive Borel measure  $\sigma$  and denote by  $\{\phi_n\}_{n \geq 0}$  its corresponding sequence of orthonormal polynomials. If  $\sigma \in \mathcal{N}$ , then uniformly in  $\mathbb{C} \setminus \overline{\mathbb{D}}$*

$$\lim_{n \rightarrow \infty} \frac{\phi_{n+1}^{(j)}(z)}{\phi_n^{(j)}(z)} = z, \quad \lim_{n \rightarrow \infty} \frac{\phi_n^{(j)}(z)}{\phi_n^{(j+1)}(z)} = 0, \quad j \geq 0.$$

*Proof.* According to Lemma 5.1.1,

$$\frac{\phi_{n+1}^{(j)}(z)}{\phi_n^{(j)}(z)} = \frac{\phi_n^{(j-1)}(z)}{\phi_n^{(j)}(z)} \left( \frac{\phi_{n+1}^{(j-1)}(z)}{\phi_n^{(j-1)}(z)} \right)' + \frac{\phi_{n+1}^{(j-1)}(z)}{\phi_n^{(j-1)}(z)}. \quad (5.7)$$

Using induction in  $j$ , we get uniformly in  $\mathbb{C} \setminus \overline{\mathbb{D}}$ ,

$$\lim_{n \rightarrow \infty} \left( \frac{\phi_{n+1}^{(j-1)}(z)}{\phi_n^{(j-1)}(z)} \right)' = 1, \quad \lim_{n \rightarrow \infty} \frac{\phi_n^{(j-1)}(z)}{\phi_n^{(j)}(z)} = 0.$$

Therefore, if  $n$  tends to infinity in (5.7), the result follows.  $\square$

**Corollary 5.1.2.** *If  $\sigma \in \mathcal{N}$ , then uniformly in  $\mathbb{C} \setminus \overline{\mathbb{D}}$*

$$\lim_{n \rightarrow \infty} \frac{\phi_n^{*(j)}(z)}{\phi_n^{(j)}(z)} = 0, \quad \lim_{n \rightarrow \infty} \frac{K_{n-1}^{(l,r)}(z,y)}{\phi_n^{(i)}(z)\phi_n^{(j)}(y)} = 0, \quad 0 \leq l < i, 0 \leq r < j.$$

*Proof.* From Lemma 5.1.1, we have

$$\frac{\phi_n^{*(j)}(z)}{\phi_n^{(j)}(z)} = \frac{\phi_n^{(j-1)}(z)}{\phi_n^{(j)}(z)} \left( \frac{\phi_n^{*(j-1)}(z)}{\phi_n^{(j-1)}(z)} \right)' + \frac{\phi_n^{*(j-1)}(z)}{\phi_n^{(j-1)}(z)}. \quad (5.8)$$

Using a similar argument as in the proof of the previous lemma, the first statement follows. The second statement is a straightforward consequence of the first part of this corollary and Lemma 5.1.2.  $\square$

**Theorem 5.1.3.** *Let  $\mathcal{L}$  be a positive definite linear functional, whose associated measure  $\sigma$  satisfies the conditions of Theorem 5.1.2. Let  $\{\Psi_n\}_{n \geq 0}$  the sequence of monic orthogonal polynomials associated with  $\mathcal{L}_1$  defined as in (5.2). Then, uniformly in  $\mathbb{C} \setminus \overline{\mathbb{D}}$ ,*

$$\lim_{n \rightarrow \infty} \frac{\Psi_n(z)}{\Phi_n(z)} = 1.$$

*Proof.* From the expression (5.6),

$$\frac{\Psi_n(z)}{\Phi_n(z)} = \frac{A(z, n, \alpha)}{(z - \alpha)^2} + \frac{B(z, n, \alpha)}{(z - \alpha)^2} \frac{\Phi_n^*(z)}{\Phi_n(z)}.$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

Since, for  $z \in \mathbb{C} \setminus \overline{\mathbb{D}}$  by Corollary 5.1.2,

$$\lim_{n \rightarrow \infty} \frac{\Phi_n^*(z)}{\Phi_n(z)} = 0,$$

it suffices to show that, for  $|\alpha| = 1$ ,

$$\lim_{n \rightarrow \infty} \frac{A(z, n, \alpha)}{(z - \alpha)^2} = 1.$$

Notice that  $\lim_{n \rightarrow \infty} \Phi_n(\alpha) = O(1)$ ,  $\lim_{n \rightarrow \infty} \Phi_n'(\alpha) = O(n)$ ,  $\lim_{n \rightarrow \infty} \Phi_n^*(\alpha) = O(1)$ ,  $\lim_{n \rightarrow \infty} \Phi_n^{*\prime}(\alpha) = O(n)$ , and  $\lim_{n \rightarrow \infty} K_n(\alpha, \alpha) = O(n)$ .

On the other hand, dividing the numerator and denominator of  $\frac{A(z, n, \alpha)}{(z - \alpha)^2} - 1$  by  $n^2 K_{n-1}(\alpha, \alpha)$ , and using Theorem 5.1.2, we obtain

$$\begin{aligned} \lim_{n \rightarrow \infty} \frac{\Phi_n(\alpha) Y_{2,1}}{n^2 K_{n-1}(\alpha, \alpha)} &= O(1/n), & \lim_{n \rightarrow \infty} \frac{\Phi_n'(\alpha) Y_{2,2}}{n^2 K_{n-1}(\alpha, \alpha)} &= O(1/n), \\ \lim_{n \rightarrow \infty} \frac{\Phi_n(\alpha) Y_{1,1}}{n^2 K_{n-1}(\alpha, \alpha)} &= O(1), & \lim_{n \rightarrow \infty} \frac{\Phi_n'(\alpha) Y_{1,2}}{n^2 K_{n-1}(\alpha, \alpha)} &= O(1), \end{aligned}$$

so that the numerator of  $\frac{A(z, n, \alpha)}{(z - \alpha)^2} - 1$  behaves as  $O(1)$ . Similarly, one can show that the denominator behaves as  $O(n)$  and, therefore,

$$\lim_{n \rightarrow \infty} \frac{A(z, n, \alpha)}{(z - \alpha)^2} = 1.$$

The same arguments can be applied to  $B(z, n, \alpha)$ , what ensures the result.  $\square$

### Example: Lebesgue polynomials

We now study one example that illustrates the behavior of the Verblunsky coefficients associated with the Lebesgue measure  $d\sigma(\theta) = \frac{d\theta}{2\pi}$  and to the perturbation (5.2) given by

$$d\bar{\sigma}(\theta) = \frac{d\theta}{2\pi} + m\delta'_\alpha,$$

where  $m \in \mathbb{R}$  and  $|\alpha| = 1$ . It is very well known that  $\Phi_n(z) = z^n$  is the  $n$ -th monic orthogonal polynomial with respect to  $d\sigma$ , and thus  $\Psi_n(z)$ , the  $n$ -th monic orthogonal polynomial with respect to  $d\bar{\sigma}$ , can be obtained using (5.6). Indeed, evaluating these



---

polynomials at  $z = 0$ , for the special case  $\alpha = 1$ , is not difficult to show that

$$\Psi_n(0) = \left( \frac{n(n-1)(n+1)}{6} - \frac{in}{m} \right) \left( \frac{n^2(n-1)(n+1)}{12} - \frac{1}{m^2} \right)^{-1}. \quad (5.9)$$

From the last expression we are able to obtain the regularity condition in terms of the mass, by setting  $|\Psi_n(0)| \neq 1$ ,  $n \geq 1$ . Notice that  $|\Psi_n(0)| \rightarrow 0$ , as it can be seen from (5.9). Thus, there exists a non-negative integer  $n_0$ , depending on  $m$ , such that  $|\Psi_n(0)| < 1$  for  $n \geq n_0$ , but some of the preceding Verblunsky coefficients will be of modulus greater than 1, destroying the positivity of the perturbed functional. Indeed, from (5.3), we obtain that the positivity condition for this perturbation is

$$m^2 < \frac{12}{n^2(n^2 - 1)}, \quad n \geq 2.$$

Since the right side is a positive monotonic decreasing sequence, we only have a positive definite case if  $m = 0$ .

### 5.1.2 Mass points outside the unit circle

Now, consider a hermitian linear functional  $\mathcal{L}_2$  such that its associated bilinear functional satisfies

$$\begin{aligned} \langle f, g \rangle_{\mathcal{L}_2} &= \langle f, g \rangle_{\mathcal{L}} + im \left( \alpha^{-1} f(\alpha) \overline{g'(\bar{\alpha}^{-1})} - \alpha f'(\alpha) \overline{g(\bar{\alpha}^{-1})} \right) \\ &\quad + i\bar{m} \left( \bar{\alpha} f(\bar{\alpha}^{-1}) \overline{g'(\alpha)} - \bar{\alpha}^{-1} p'(\bar{\alpha}^{-1}) \overline{q(\alpha)} \right), \end{aligned} \quad (5.10)$$

with  $m, \alpha \in \mathbb{C}$ ,  $|\alpha| \neq 0$ , and  $|\alpha| \neq 1$ . As in the previous section, we are interested in the regularity conditions for this linear functional and the corresponding family of orthogonal polynomials. Assuming that  $\mathcal{L}_2$  is a quasi-definite linear functional and following the method used in the proof of Theorem 5.1.1, we get

$$\begin{aligned} \Psi_n(z) &= \Phi_n(z) + im \left( \alpha \Psi'_n(\alpha) K_{n-1}(z, \bar{\alpha}^{-1}) - \alpha^{-1} \Psi_n(\alpha) K_{n-1}^{(0,1)}(z, \bar{\alpha}^{-1}) \right) \\ &\quad + i\bar{m} \left( \bar{\alpha}^{-1} \Psi'_n(\bar{\alpha}^{-1}) K_{n-1}(z, \alpha) - \bar{\alpha} \Psi_n(\bar{\alpha}^{-1}) K_{n-1}^{(0,1)}(z, \alpha) \right). \end{aligned} \quad (5.11)$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

Evaluating the above expression and its first derivative in  $\alpha$  and  $\bar{\alpha}^{-1}$ , we get the following linear systems

$$\begin{aligned} \begin{bmatrix} \Phi_n(\alpha) \\ \Phi'_n(\alpha) \end{bmatrix} &= \begin{bmatrix} 1 + im\alpha^{-1}K_{n-1}^{(0,1)}(\alpha, \bar{\alpha}^{-1}) & -im\alpha K_{n-1}(\alpha, \bar{\alpha}^{-1}) \\ im\alpha^{-1}K_{n-1}^{(1,1)}(\alpha, \bar{\alpha}^{-1}) & 1 - im\alpha K_{n-1}^{(1,0)}(\alpha, \bar{\alpha}^{-1}) \end{bmatrix} \begin{bmatrix} \Psi_n(\alpha) \\ \Psi'_n(\alpha) \end{bmatrix} \\ &+ \begin{bmatrix} im\alpha K_{n-1}^{(0,1)}(\alpha, \alpha) & -im\alpha^{-1}K_{n-1}(\alpha, \alpha) \\ im\alpha K_{n-1}^{(1,1)}(\alpha, \alpha) & -im\alpha^{-1}K_{n-1}^{(1,0)}(\alpha, \alpha) \end{bmatrix} \begin{bmatrix} \Psi_n(\bar{\alpha}^{-1}) \\ \Psi'_n(\bar{\alpha}^{-1}) \end{bmatrix}, \end{aligned} \quad (5.12)$$

$$\begin{aligned} \begin{bmatrix} \Phi_n(\bar{\alpha}^{-1}) \\ \Phi'_n(\bar{\alpha}^{-1}) \end{bmatrix} &= \begin{bmatrix} im\alpha^{-1}K_{n-1}^{(0,1)}(\bar{\alpha}^{-1}, \bar{\alpha}^{-1}) & -im\alpha K_{n-1}(\bar{\alpha}^{-1}, \bar{\alpha}^{-1}) \\ im\alpha^{-1}K_{n-1}^{(1,1)}(\bar{\alpha}^{-1}, \bar{\alpha}^{-1}) & -im\alpha K_{n-1}^{(1,0)}(\bar{\alpha}^{-1}, \bar{\alpha}^{-1}) \end{bmatrix} \begin{bmatrix} \Psi_n(\alpha) \\ \Psi'_n(\alpha) \end{bmatrix} \\ &+ \begin{bmatrix} 1 + im\alpha K_{n-1}^{(0,1)}(\bar{\alpha}^{-1}, \alpha) & -im\alpha^{-1}K_{n-1}(\bar{\alpha}^{-1}, \alpha) \\ im\alpha K_{n-1}^{(1,1)}(\bar{\alpha}^{-1}, \alpha) & 1 - im\alpha^{-1}K_{n-1}^{(1,0)}(\bar{\alpha}^{-1}, \alpha) \end{bmatrix} \begin{bmatrix} \Psi_n(\bar{\alpha}^{-1}) \\ \Psi'_n(\bar{\alpha}^{-1}) \end{bmatrix}, \end{aligned} \quad (5.13)$$

which yields into the system of 4 linear equations with 4 unknowns

$$\begin{bmatrix} \Phi_n(\alpha) \\ \Phi_n(\bar{\alpha}^{-1}) \end{bmatrix} = \begin{bmatrix} \mathbf{I}_2 + m\mathbb{K}_{n-1}(\alpha, \bar{\alpha}^{-1})\mathbf{D}(\alpha) & \bar{m}\mathbb{K}_{n-1}(\alpha, \alpha)\mathbf{D}(\bar{\alpha}^{-1}) \\ m\mathbb{K}_{n-1}(\bar{\alpha}^{-1}, \bar{\alpha}^{-1})\mathbf{D}(\alpha) & \mathbf{I}_2 + \bar{m}\mathbb{K}_{n-1}(\bar{\alpha}^{-1}, \alpha)\mathbf{D}(\bar{\alpha}^{-1}) \end{bmatrix} \begin{bmatrix} \Psi_n(\alpha) \\ \Psi_n(\bar{\alpha}^{-1}) \end{bmatrix},$$

where  $(\mathbf{Q}, \mathbf{R})^T = (Q, Q', R, R')^T$ . Thus, in order  $\mathcal{L}_2$  to be a quasi-definite linear functional, we need that the  $4 \times 4$  matrix defined as above must be non-singular. On the other hand,

$$\begin{bmatrix} \Psi_n(\alpha) \\ \Psi_n(\bar{\alpha}^{-1}) \end{bmatrix} = \begin{bmatrix} \mathbf{I}_2 + m\mathbb{K}_{n-1}(\alpha, \bar{\alpha}^{-1})\mathbf{D}(\alpha) & \bar{m}\mathbb{K}_{n-1}(\alpha, \alpha)\mathbf{D}(\bar{\alpha}^{-1}) \\ m\mathbb{K}_{n-1}(\bar{\alpha}^{-1}, \bar{\alpha}^{-1})\mathbf{D}(\alpha) & \mathbf{I}_2 + \bar{m}\mathbb{K}_{n-1}(\bar{\alpha}^{-1}, \alpha)\mathbf{D}(\bar{\alpha}^{-1}) \end{bmatrix}^{-1} \begin{bmatrix} \Phi_n(\alpha) \\ \Phi_n(\bar{\alpha}^{-1}) \end{bmatrix}.$$

As a consequence, from (5.11), we get

$$\Psi_n(z) = \Phi_n(z) - m \begin{bmatrix} K_{n-1}(z, \bar{\alpha}^{-1}) \\ K_{n-1}^{(0,1)}(z, \bar{\alpha}^{-1}) \end{bmatrix}^T \mathbf{D}(\alpha) \Psi_n(\alpha) - \bar{m} \begin{bmatrix} K_{n-1}(z, \alpha) \\ K_{n-1}^{(0,1)}(z, \alpha) \end{bmatrix}^T \mathbf{D}(\bar{\alpha}^{-1}) \Psi_n(\bar{\alpha}^{-1}), \quad (5.14)$$

where  $\Psi_n(\alpha)$  and  $\Psi_n(\bar{\alpha}^{-1})$  can be obtained from the above linear system. Assuming that the regularity conditions hold, and following the method used in the proof of Theorem 5.1.1, it is not difficult to show that  $\{\Psi_n\}_{n \geq 0}$ , defined as in (5.14), is the sequence of monic orthogonal polynomials with respect to  $\mathcal{L}_2$ .

---

### 5.1.2.1 Outer relative asymptotics

The following result was proved in [70] using a different method, and it has been generalized for rectifiable Jordan curves or arcs in [20]. We show here another proof of the same result.

**Lemma 5.1.3.** *If  $\sigma \in \mathcal{N}$ , then uniformly in  $\mathbb{C} \setminus \overline{\mathbb{D}}$ ,*

$$\lim_{n \rightarrow \infty} \frac{K_{n-1}^{(i,j)}(z,y)}{\phi_n^{(i)}(z)\overline{\phi_n^{(j)}(y)}} = \frac{1}{z\bar{y}-1}, \quad i, j \geq 0.$$

*Proof.* From the Christoffel-Darboux formula (2.33), we obtain

$$\overline{\phi_n^*(z)\phi_n^{*(j)}(y)} - \overline{\phi_n(z)\phi_n^{(j)}(y)} = (1 - z\bar{y})K_{n-1}^{(0,j)}(z,y) - jzK_{n-1}^{(0,j-1)}(z,y),$$

and, as a consequence,

$$\begin{aligned} \overline{\phi_n^{*(i)}(z)\phi_n^{*(j)}(y)} - \overline{\phi_n^{(i)}(z)\phi_n^{(j)}(y)} &= (1 - z\bar{y})K_{n-1}^{(i,j)}(z,y) - k\bar{y}K_{n-1}^{(i-1,j)}(z,y) \\ &\quad - j\left(zK_{n-1}^{(i,j-1)}(z,y) + kK_{n-1}^{(k-1,j-1)}(z,y)\right). \end{aligned}$$

Thus, dividing by  $\overline{\phi_n^{(i)}(z)\phi_n^{(j)}(y)}$  and using Corollary 5.1.2 when  $n$  tends to infinity, the result follows.  $\square$

**Remark 5.1.1.** *Notice that*

$$\lim_{n \rightarrow \infty} \frac{K_{n-1}^{(j,j)}(\alpha, \alpha)}{|\phi_n^{(j)}(\alpha)|^2} = \frac{1}{|\alpha|^2 - 1}, \quad |\alpha| > 1, \quad j \geq 1.$$

It is possible to obtain a generalization of Theorem 5.1.3 for the sequence of monic orthogonal polynomials associated with (5.10). As above, we can express (5.14) as in (5.6). Using the Christoffel-Darboux formula (2.33), we obtain

$$\Psi_n(z) = \left(1 + \widetilde{A}(z, n, \alpha)\right)\Phi_n(z) + \widetilde{B}(z, n, \alpha)\Phi_n^*(z),$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

with

$$\begin{aligned}\widetilde{A}(z, n, \alpha) &= im\bar{\alpha}^{-1} \frac{\overline{\Phi'_n(\bar{\alpha}^{-1})(1-\alpha^{-1}z) + z\overline{\Phi_n(\bar{\alpha}^{-1})}}}{\mathbf{k}_n(1-\alpha^{-1})^2} \Psi_n(\alpha) - im\alpha \frac{\overline{\Phi_n(\bar{\alpha}^{-1})}}{\mathbf{k}_n(1-\alpha^{-1}z)} \Psi'_n(\alpha) \\ &\quad + i\bar{m}\alpha \frac{\overline{\Phi'_n(\alpha)(1-\bar{\alpha}z) + z\overline{\Phi_n(\alpha)}}}{\mathbf{k}_n(1-\bar{\alpha})^2} \Psi_n(\bar{\alpha}^{-1}) - i\bar{m}\alpha^{-1} \frac{\overline{\Phi_n(\alpha)}}{\mathbf{k}_n(1-\bar{\alpha}z)} \Psi'_n(\bar{\alpha}^{-1}), \\ \widetilde{B}(z, n, \alpha) &= im\alpha \frac{\overline{\Phi_n(\bar{\alpha}^{-1})}}{\mathbf{k}_n(1-\alpha^{-1}z)} \Psi'_n(\alpha) - im\bar{\alpha}^{-1} \frac{\overline{\Phi_n^*(\bar{\alpha}^{-1})(1-\alpha^{-1}z) + z\overline{\Phi_n^*(\bar{\alpha}^{-1})}}}{\mathbf{k}_n(1-\alpha^{-1})^2} \Psi_n(\alpha) \\ &\quad + i\bar{m}\alpha^{-1} \frac{\overline{\Phi_n^*(\alpha)}}{\mathbf{k}_n(1-\bar{\alpha}z)} \Psi'_n(\bar{\alpha}^{-1}) - i\bar{m}\alpha \frac{\overline{\Phi_n^*(\alpha)(1-\bar{\alpha}z) + z\overline{\Phi_n^*(\alpha)}}}{\mathbf{k}_n(1-\bar{\alpha})^2} \Psi_n(\bar{\alpha}^{-1}),\end{aligned}$$

where the values of  $\Psi_n(\alpha)$ ,  $\Psi'_n(\alpha)$ ,  $\Psi_n(\bar{\alpha}^{-1})$ , and  $\Psi'_n(\bar{\alpha}^{-1})$  can be obtained by solving the  $4 \times 4$  linear system shown above. Denoting the entries of the  $2 \times 2$  matrices in (5.12) - (5.13) by  $\{b_{i,j}\}$ ,  $\{c_{i,j}\}$ ,  $\{a_{i,j}\}$  and  $\{d_{i,j}\}$ , respectively, we get

$$\begin{aligned}\Psi_n(\alpha) &= (d_{1,1}\Phi_n(\alpha) + d_{1,2}\Phi'_n(\alpha) + c_{1,1}\Phi_n(\bar{\alpha}^{-1}) + c_{1,2}\Phi'_n(\bar{\alpha}^{-1})) / \Delta, \\ \Psi'_n(\alpha) &= (d_{2,1}\Phi_n(\alpha) + d_{2,2}\Phi'_n(\alpha) + c_{2,1}\Phi_n(\bar{\alpha}^{-1}) + c_{2,2}\Phi'_n(\bar{\alpha}^{-1})) / \Delta, \\ \Psi_n(\bar{\alpha}^{-1}) &= (a_{1,1}\Phi_n(\alpha) + a_{1,2}\Phi'_n(\alpha) + b_{1,1}\Phi_n(\bar{\alpha}^{-1}) + b_{1,2}\Phi'_n(\bar{\alpha}^{-1})) / \Delta, \\ \Psi_n(\bar{\alpha}^{-1}) &= (a_{2,1}\Phi_n(\alpha) + a_{2,2}\Phi'_n(\alpha) + b_{2,1}\Phi_n(\bar{\alpha}^{-1}) + b_{2,2}\Phi'_n(\bar{\alpha}^{-1})) / \Delta,\end{aligned}$$

where  $\Delta$  is the determinant of the  $4 \times 4$  matrix. To get the asymptotic result, it suffices to show that  $\widetilde{A}(z, n, \alpha) \rightarrow 0$  and  $\widetilde{B}(z, n, \alpha) \rightarrow 0$  as  $n \rightarrow \infty$ . First, notice that applying the corresponding derivatives to the Christoffel-Darboux formula (2.33), we obtain

$$\begin{aligned}K_{n-1}^{(0,1)}(z, y) &= \frac{\overline{\Phi_n^{*'}(y)\Phi_n^*(z) - \Phi'_n(y)\Phi_n(z)}}{\mathbf{k}_n(1-\bar{y}z)} + \frac{zK_{n-1}(z, y)}{1-\bar{y}z}, \\ K_{n-1}^{(1,0)}(z, y) &= \frac{\overline{\Phi_n^*(y)\Phi_n^{*'}(z) - \Phi_n(y)\Phi'_n(z)}}{\mathbf{k}_n(1-\bar{y}z)} + \frac{\bar{y}K_{n-1}(z, y)}{1-\bar{y}z}, \\ K_{n-1}^{(1,1)}(z, y) &= \frac{\overline{\Phi_n^{*'}(y)\Phi_n^{*'}(z) - \Phi'_n(y)\Phi'_n(z)}}{\mathbf{k}_n(1-\bar{y}z)} + \frac{zK_{n-1}^{(1,0)}(z, y) + \bar{y}K_{n-1}^{(0,1)}(z, y) + K_{n-1}(z, y)}{1-\bar{y}z}.\end{aligned}$$

On the other hand, if  $\mathcal{L}$  is positive definite, and its corresponding measure  $\sigma \in \mathcal{N}$ , then by Corollary 5.1.2 (see also [137]) we have  $\Phi_n(\alpha) = O(\alpha^n)$ ,  $\Phi'_n(\alpha) = O(n\alpha^n)$ , and

$$\lim_{n \rightarrow \infty} \frac{\Phi_n(\alpha)}{\Phi_n^*(\alpha)} = 0, \quad |\alpha| < 1, \quad \lim_{n \rightarrow \infty} \frac{\Phi_n^*(\alpha)}{\Phi_n(\alpha)} = 0, \quad |\alpha| > 1.$$

---

Assume, without loss of generality, that  $|\alpha| < 1$ . If  $|\alpha| < 1$  and  $\sigma \in \mathcal{S}$ , notice that  $\Phi_n(\alpha)$  and  $\Phi_n^*(\alpha)$  are  $\mathcal{O}(\alpha^n)$ , then  $\lim_{n \rightarrow \infty} K_n(\alpha, \alpha) < \infty$  and  $K_n(\bar{\alpha}^{-1}, \bar{\alpha}^{-1}) = \mathcal{O}(|\alpha|^{-2n})$ , as well as  $\overline{K_n(\alpha, \bar{\alpha}^{-1})} = K_n(\bar{\alpha}^{-1}, \alpha) = \mathcal{O}(n)$ . Observe that, except for the entries containing  $K_{n-1}(\alpha, \alpha)$  and their derivatives, all other entries of the  $4 \times 4$  matrix diverge, and thus its determinant diverges much faster than any other term in the expressions for  $\Psi_n(\alpha)$ ,  $\Psi_n'(\alpha)$ ,  $\Psi_n(\bar{\alpha}^{-1})$  and  $\Psi_n'(\bar{\alpha}^{-1})$ , so that  $\widetilde{A}(z, n, \alpha) \rightarrow 0$  and  $\widetilde{B}(z, n, \alpha) \rightarrow 0$  as  $n$  tends to  $\infty$ . As a consequence,

**Theorem 5.1.4.** *Let  $\mathcal{L}$  be a positive definite linear functional, whose associated measure  $\sigma \in \mathcal{S}$ . Let  $\{\Psi_n\}_{n \geq 0}$  the sequence of monic orthogonal polynomials associated to  $\mathcal{L}_2$  defined as in (5.10). Then, uniformly in  $\mathbb{C} \setminus \mathbb{T}$ ,*

$$\lim_{n \rightarrow \infty} \frac{\Psi_n(z)}{\Phi_n(z)} = 1.$$

### 5.1.3 C-functions and linear spectral transformations

First, we assume that  $|\alpha| = 1$ . Let us consider the moments associated with  $\mathcal{L}_1$ . Notice that  $\widetilde{c}_0 = c_0$ . For  $k \geq 1$ , we have  $\widetilde{c}_k = \langle z^k, 1 \rangle_{\mathcal{L}_1} = c_k - imk\alpha^k$ . In a similar way,  $\widetilde{c}_{-k} = c_{-k} + imk\bar{\alpha}^k$ . Therefore,

$$\begin{aligned} F_1(z) &= \widetilde{c}_0 + 2 \sum_{k=1}^{\infty} \widetilde{c}_{-k} z^k = c_0 + 2 \sum_{k=1}^{\infty} (c_{-k} + imk\bar{\alpha}^k) z^k = F(z) + 2im \sum_{k=1}^{\infty} k\bar{\alpha}^k z^k \\ &= F(z) + \frac{2im\alpha}{z-\alpha} + \frac{2im\alpha^2}{(z-\alpha)^2}. \end{aligned}$$

This means that the resulting  $C$ -function is a perturbation of  $F$  by the addition of a rational function with a double pole at  $z = \alpha$ .

Now, we assume  $|\alpha| > 1$ , and let consider the moments associated with  $\mathcal{L}_2$ . Notice that  $\widehat{c}_0 = c_0$ . For  $k \in \mathbb{N}$ , we have from (5.10),

$$\widehat{c}_k = c_k - imk\alpha^k - imk\bar{\alpha}^{-k}, \quad \widehat{c}_{-k} = c_{-k} + imk\bar{\alpha}^k + imk\alpha^{-k},$$

and, as a consequence,

$$F_2(z) = \widehat{c}_0 + 2 \sum_{k=1}^{\infty} \widehat{c}_{-k} z^k = F(z) - \frac{2im\alpha}{z-\alpha} + \frac{2im\alpha^2}{(z-\alpha)^2} - \frac{2im\bar{\alpha}^{-1}}{z-\bar{\alpha}^{-1}} + \frac{2im\bar{\alpha}^{-2}}{(z-\bar{\alpha}^{-1})^2}. \quad (5.15)$$

This means that the resulting  $C$ -function is a perturbation of the initial one by the

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

addition of a rational function with two double poles at  $\alpha$  and  $\bar{\alpha}^{-1}$ .

### Connection with special linear transformations

We show that perturbations  $\mathcal{L}_2$  can be expressed in terms of special cases of the spectral transformations (2.44) and (2.45). Let  $\mathcal{F}_C(\alpha)$  and  $\mathcal{F}_G(\alpha, m)$  be linear spectral transformations associated with the modification of the original functional  $\mathcal{L}$  given by

$$\langle \mathcal{L}_C, f \rangle = \langle \mathcal{L}, |z - \alpha|^2 f(z) \rangle \quad (5.16)$$

and

$$\langle \mathcal{L}_G, |z - \alpha|^2 f(z) \rangle = \langle \mathcal{L}, f \rangle + m\delta_\alpha, \quad |\alpha| = 1, \quad m \in \mathbb{R}, \quad (5.17)$$

respectively [81; 82]. The polynomial coefficients associated with  $\mathcal{F}_C(\alpha)$  and  $\mathcal{F}_G(\alpha, m)$  are

$$\begin{aligned} A_C(z) = D_G(z) &= (z - \alpha)(1 - \bar{\alpha}z), & D_C(z) = A_G(z) &= z, \\ B_C(z) &= -\bar{\alpha}c_0z^2 + (\alpha c_{-1} - \bar{\alpha}c_1)z + \alpha c_0, & B_G(z) &= \bar{\alpha}\tilde{c}_0z^2 + 2i\Im(q_0)z - \alpha\tilde{c}_0, \end{aligned}$$

where  $q_0$  is a free parameter that depends on the mass used in (5.17). Now, consider the following product of transformations

$$\mathcal{F}_D = \mathcal{F}_{G_2}(\alpha, m_2) \circ \mathcal{F}_{G_1}(\alpha, m_1) \circ \mathcal{F}_{C_2}(\alpha) \circ \mathcal{F}_{C_1}(\alpha). \quad (5.18)$$

It is not difficult to show that  $F_D$ , the  $C$ -function associated with  $\mathcal{F}_D$ , is given by

$$\begin{aligned} F_D(z) &= F(z) + \frac{B_{C_1}(z)}{D_{G_1}(z)} + \frac{B_{G_2}(z)}{D_{G_2}(z)} + \frac{B_{C_2}(z)A_{G_1}(z)}{D_{G_1}(z)D_{G_2}(z)} + \frac{B_{G_1}(z)A_{G_2}(z)}{D_{G_1}(z)D_{G_2}(z)} \\ &= F(z) + \frac{B_{C_1}(z) + B_{G_2}(z)}{(z - \alpha)(1 - \bar{\alpha}z)} + \frac{z(B_{C_2}(z) + B_{G_1}(z))}{(z - \alpha)^2(1 - \bar{\alpha}z)^2}. \end{aligned}$$

Assuming that all transformations are normalized, i.e., all of the first moments are equal to 1, and denoting  $K_1 = \alpha c_{-1} - \bar{\alpha}c_1 + 2i\Im(q_0^{(1)})$  and  $K_2 = \alpha c_{-1} - \bar{\alpha}c_1 + 2i\Im(q_0^{(2)})$ , where  $q_0^{(1)}$  and  $q_0^{(2)}$  are the free parameters associated with  $\mathcal{F}_{G_1}$  and  $\mathcal{F}_{G_2}$ , respectively,

we obtain

$$\begin{aligned}
F_D(z) &= F(z) + \frac{K_2 z}{(z-\alpha)(1-\bar{\alpha}z)} + \frac{K_1 z^2}{(z-\alpha)^2(1-\bar{\alpha}z)^2} \\
&= F(z) + \frac{L_1}{(z-\alpha)} + \frac{L_2}{(z-\alpha)^2} + \frac{L_3}{(z-\bar{\alpha}^{-1})} + \frac{L_4}{(z-\bar{\alpha}^{-1})^2}, \tag{5.19}
\end{aligned}$$

for some constants  $L_1, L_2, L_3$ , and  $L_4$ , satisfying

$$\begin{aligned}
-\bar{\alpha}K_2 &= L_1 + L_3, \\
(1+|\alpha|^2)K_2 + K_1 &= -(\alpha + 2\bar{\alpha}^{-1})L_1 + L_2 - (2\alpha + \bar{\alpha}^{-1})L_3 + L_4, \\
-\alpha K_2 &= (\bar{\alpha}^{-2} + 2\alpha\bar{\alpha}^{-1})L_1 - 2\bar{\alpha}^{-1}L_2 + (\alpha^2 + 2\bar{\alpha}^{-1}\alpha)L_3 - 2\alpha L_4, \\
0 &= -\alpha\bar{\alpha}^{-2}L_1 + \bar{\alpha}^{-2}L_2 - \alpha^2\bar{\alpha}^{-1}L_3 + \alpha^2L_4.
\end{aligned}$$

Furthermore, comparing (5.15) and (5.19), we have  $L_2 = -\alpha L_1$  and  $L_4 = -\bar{\alpha}^{-1}L_3$ . Solving the above system we arrive at

$$L_1 = \frac{\alpha|\alpha|^2}{1-|\alpha|^2}K_2, \quad L_3 = -\frac{\alpha}{1-|\alpha|^2}K_2,$$

and thus, we conclude that transformation (5.18) is equivalent to  $\mathcal{F}_2(\alpha^{-1}, m)$ , the transformation associated with (5.10), with

$$m = \frac{|\alpha|^2}{2i(1-|\alpha|^2)}K_2.$$

## 5.2 Non-standard inner products

In the last few years, some attention has been paid to the asymptotic properties of orthogonal polynomials with respect to non-standard inner products. In particular, the algebraic and analytic properties of orthogonal polynomials associated with a Sobolev inner product have attracted the interest of many researchers, see [132] for an updated overview with more than 300 references.

A discrete Sobolev inner product in  $\mathbb{C} \setminus \bar{\mathbb{D}}$  is given by

$$\langle f, g \rangle_S = \int_{\mathbb{T}} f(z)\overline{g(z)}d\sigma(z) + \mathbf{f}(Z) \mathbf{A} \mathbf{g}(Z)^H, \tag{5.20}$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

where

$$\mathbf{f}(Z) = \left( f(\alpha_1), \dots, f^{(l_1)}(\alpha_1), \dots, f(\alpha_m), \dots, f^{(l_m)}(\alpha_m) \right),$$

$\mathbf{A}$  is an  $M \times M$  positive semi-definite hermitian matrix, with  $M = l_1 + \dots + l_m + m$ , and  $|\alpha_i| > 1$ ,  $i = 1, \dots, m$ . Since  $\mathbf{A}$  is positive semi-definite, the inner product (5.20) is positive definite. Therefore, there exists a sequence of polynomials  $\{\psi_n\}_{n \geq 0}$ ,

$$\psi_n(z) = \gamma_n z^n + (\text{lower degree terms}), \quad \gamma_n > 0,$$

which is orthonormal with respect to (5.20). We are interested in the outer relative asymptotic behavior of  $\{\psi_n\}_{n \geq 0}$  with respect to the sequence  $\{\phi_n\}_{n \geq 0}$  of orthonormal polynomials with respect to  $\sigma$ . We show that if  $\sigma \in \mathcal{N}$  and  $\mathbf{A}$  is positive definite, then this outer relative asymptotics follows. Similar results have been obtained for the case when the measure is supported on a bounded interval of the real line [118; 134].

### 5.2.1 Outer relative asymptotics

In [70; 116; 126], the relative asymptotic behavior of orthogonal polynomials with respect to a discrete Sobolev inner product on the unit circle was studied. In this section, we propose a slightly modified outline.

The nondiagonal structure of the matrix  $\mathbf{A}$  makes the analysis of the situation much more difficult. First of all, let us prove an important result which gives precise information about the matrix  $\mathbf{A}$ .

**Lemma 5.2.1.** *The outer relative asymptotic behavior of orthogonal polynomials with respect to the inner product (5.20) does not depend on the matrix  $\mathbf{A}$ .*

*Proof.* Let  $\{\tilde{\psi}_n\}_{n \geq 0}$  be the sequence of orthonormal polynomials with respect to the inner product

$$\langle f, g \rangle_{\tilde{\sigma}} = \int_{\mathbb{T}} f(z) \overline{g(z)} d\sigma(z) + \mathbf{f}(Z) \mathbf{B} \mathbf{g}(Z)^H,$$

where  $\mathbf{B}$  is an arbitrary positive definite hermitian matrix of order  $M$ . Expanding  $\psi_n$  in terms of  $\{\phi_n\}_{n \geq 0}$ , we have

$$\psi_n(z) = \frac{\gamma_n}{\kappa_n} \phi_n(z) + \sum_{k=0}^{n-1} \lambda_{n,k} \phi_k(z) \tag{5.21}$$



where

$$\lambda_{n,k} = \int_{\mathbb{T}} \psi_n(z) \overline{\phi(z)} d\sigma(z) = -\psi_n(Z) \mathbf{A} \phi_n(Z).$$

Substituting this expression in (5.21), we obtain

$$\psi_n(z) = \frac{\gamma_n}{\kappa_n} \phi_n(z) - \psi_n(Z) \mathbf{A} \mathbf{K}_n(z, Z)^T, \quad (5.22)$$

where  $\mathbf{K}_n(z, Z) = (K_n(z, \alpha_1), \dots, K_n^{(0, l_1)}(z, \alpha_1), \dots, K_n(z, \alpha_m), \dots, K_n^{(0, l_m)}(z, \alpha_m))$  and  $K_n^{(i, j)}(z, y)$  denotes the  $i$ -th (resp.  $j$ -th) partial derivative of  $K_n(z, y)$  with respect to the variable  $z$  (resp.  $y$ ). In an analogous way, we get

$$\widetilde{\psi}_n(z) = \frac{\widetilde{\gamma}_n}{\kappa_n} \phi_n(z) - \widetilde{\psi}_n(Z) \mathbf{B} \mathbf{K}_n(z, Z)^T, \quad (5.23)$$

where  $\widetilde{\gamma}_n$  is the leading coefficient of  $\psi_n$ . From (5.22) and (5.23) and following the method used in the proof of Theorem 5.1.1, we get [70; 116]

$$\begin{aligned} \frac{\widetilde{\gamma}_n \widetilde{\psi}_n(z)}{\gamma_n \psi_n(z)} &= \frac{\det(\mathbf{I} + \mathbf{A} \mathbb{T}_n) \det(\mathbf{I} + \mathbf{B} \mathbb{K}_n)}{\det(\mathbf{I} + \mathbf{B} \mathbb{T}_n) \det(\mathbf{I} + \mathbf{A} \mathbb{K}_n)}, \\ \left( \frac{\widetilde{\gamma}_n}{\gamma_n} \right)^2 &= \frac{\det(\mathbf{I} + \mathbf{B} \mathbb{K}_n) \det(\mathbf{I} + \mathbf{A} \mathbb{K}_{n+1})}{\det(\mathbf{I} + \mathbf{A} \mathbb{K}_n) \det(\mathbf{I} + \mathbf{B} \mathbb{K}_{n+1})}, \end{aligned}$$

where  $\mathbb{K}_n$  is a positive definite matrix of order  $M$ ,  $n \geq M$ , which can be described by blocks. The  $r, s$  block of  $\mathbb{K}_n$  is the  $(l_r + 1) \times (l_s + 1)$  matrix

$$\left( K_n^{(i, j)}(z_r, \overline{z_s}) \right)_{i=0, \dots, l_r}^{j=0, \dots, l_s}, \quad r, s = 0, \dots, m.$$

$\mathbb{T}_n$  is obtained through the following equation  $\mathbb{T}_n = \mathbb{K}_n + \mathbb{V}_n$ , where  $\mathbb{V}_n = -\frac{1}{\phi_n(z)} \mathbf{K}_n(z, Z)^T \phi_n(Z)$ . Since [70; 126]

$$\lim_{n \rightarrow \infty} \frac{\det(\mathbf{I} + \mathbf{A} \mathbb{K}_n)}{\det(\mathbf{I} + \mathbf{B} \mathbb{K}_n)} = \lim_{n \rightarrow \infty} \frac{\det(\mathbf{I} + \mathbf{A} \mathbb{T}_n)}{\det(\mathbf{I} + \mathbf{B} \mathbb{T}_n)} = \frac{\det \mathbf{A}}{\det \mathbf{B}},$$

we can deduce that

$$\lim_{n \rightarrow \infty} \frac{\widetilde{\gamma}_n \widetilde{\psi}_n(z)}{\gamma_n \psi_n(z)} = 1, \quad \lim_{n \rightarrow \infty} \left( \frac{\widetilde{\gamma}_n}{\gamma_n} \right)^2 = 1,$$

and the lemma is proved.  $\square$

For the discrete Sobolev inner product with a single mass point associated with

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

(5.20),

$$\langle f, g \rangle_{S_1} = \int_{\mathbb{T}} f(z) \overline{g(z)} d\sigma(z) + \lambda f^{(j)}(\alpha) \overline{g^{(j)}(\alpha)}, \quad |\alpha| > 1, \quad (5.24)$$

we have

**Lemma 5.2.2.** *Let  $\{\psi_{n;1}\}_{n \geq 0}$ ,  $\psi_{n;1} = \gamma_{n;1} z^n + (\text{lower degree terms})$  be the sequence of orthonormal polynomials with respect to (5.24). If  $\sigma \in \mathcal{N}$ , then*

$$\lim_{n \rightarrow \infty} \frac{\gamma_{n;1}}{\kappa_n} = \frac{1}{|\alpha|}.$$

*Proof.* From (5.22) we have

$$\psi_{n;1}(z) = \frac{\gamma_{n;1}}{\kappa_n} \phi_n(z) - \lambda \psi_{n;1}^{(j)}(\alpha) K_{n-1}^{(0,j)}(z, \alpha). \quad (5.25)$$

Taking derivatives in (5.25) and evaluating at  $z = \alpha$ , we get

$$\psi_{n;1}^{(j)}(\alpha) = \frac{\gamma_{n;1} / \kappa_n \phi_n^{(j)}(\alpha)}{1 + \lambda K_{n-1}^{(j,j)}(\alpha, \alpha)}. \quad (5.26)$$

Thus, (5.26) yields

$$\left( \frac{\beta_n}{\alpha_n} \right)^2 = \frac{1 + \lambda K_{n-1}^{(j,j)}(\alpha, \alpha)}{1 + \lambda K_n^{(j,j)}(\alpha, \alpha)}.$$

Using the previous identity and Lemma 5.1.2,

$$\lim_{n \rightarrow \infty} \frac{\gamma_{n;1}^2}{\kappa_n^2} = \lim_{n \rightarrow \infty} \frac{1 + \lambda K_{n-1}^{(j,j)}(\alpha, \alpha)}{1 + \lambda K_n^{(j,j)}(\alpha, \alpha)} = \lim_{n \rightarrow \infty} \frac{|\phi_{n-1}^{(j)}(\alpha)|^2}{|\phi_n^{(j)}(\alpha)|^2} = \frac{1}{|\alpha|^2},$$

and the lemma is proved. □

Using the previous lemma, we prove the relative asymptotics in  $\mathbb{C} \setminus \overline{\mathbb{D}}$ .

**Theorem 5.2.1.** *If  $\sigma \in \mathcal{N}$ , then uniformly in  $\mathbb{C} \setminus \overline{\mathbb{D}}$*

$$\lim_{n \rightarrow \infty} \frac{\psi_{n;1}(z)}{\phi_n(z)} = B(\alpha), \quad B(\alpha) = \frac{\overline{\alpha}(z - \alpha)}{|\alpha|(\overline{\alpha}z - 1)}. \quad (5.27)$$

*Proof.* From (5.25), we have

$$\frac{\psi_{n;1}(z)}{\phi_n(z)} = \frac{\gamma_{n;1}}{\kappa_n} - \lambda \psi_{n;1}^{(j)}(\alpha) \overline{\phi_n^{(j)}(\alpha)} \frac{K_{n-1}^{(0,j)}(z, \alpha)}{\phi_n(z) \phi_n^{(j)}(\alpha)}. \quad (5.28)$$

Using (5.26), we obtain

$$\lim_{n \rightarrow \infty} \lambda \psi_{n;1}^{(j)}(\alpha) \overline{\phi_n^{(j)}(\alpha)} = \left( |\alpha| - \frac{1}{|\alpha|} \right). \quad (5.29)$$

The outer relative asymptotics (5.27) follows letting  $n$  tends to infinity in (5.28), using Lemma 5.2.2, Lemma 5.1.3, and (5.29).  $\square$

From Theorem 5.2.1 we can see that the outer relative asymptotic behavior of orthogonal polynomials associated with (5.24) does not depend on the specific choice of  $j$  and  $\lambda$ .

**Lemma 5.2.3.**  $\sigma \in \mathcal{N}$ , then  $S_1 \in \mathcal{N}$ .

*Proof.* Assume, without loss of generality, that  $j = 0$  and  $\lambda = 1$ . From (5.25) and (5.26) we get

$$\psi_{n;1}(z) = \frac{\gamma_{n;1}}{\kappa_n} \phi_n(z) - \frac{\phi_n(\alpha)}{1 + K_{n-1}(\alpha, \alpha)} K_{n-1}(z, \alpha). \quad (5.30)$$

The evaluation at  $z = 0$  of this last expression yields

$$\frac{\psi_{n;1}(0)}{\gamma_{n;1}} = \frac{\phi_n(0)}{\kappa_n} - \frac{|\phi_n(\alpha)|^2}{1 + K_{n-1}(\alpha, \alpha)} \frac{K_{n-1}(0, \alpha)}{\gamma_{n;1} \overline{\phi_n(\alpha)}},$$

and using the Christoffel-Darboux formula (2.33), we obtain

$$\frac{K_{n-1}(0, \alpha)}{\gamma_{n;1} \overline{\phi_n(\alpha)}} = \frac{\kappa_n}{\gamma_{n;1}} \left( \frac{\overline{\phi_n^*(\alpha)}}{\phi_n(\alpha)} - \frac{\phi_n(0)}{\kappa_n} \right). \quad (5.31)$$

From Corollary 5.1.2, under our conditions, the following limit holds  $\lim_{n \rightarrow \infty} \frac{\overline{\phi_n^*(\alpha)}}{\phi_n(\alpha)} = 0$ . Since  $K_n(\alpha, \alpha)$  is an increasing sequence and  $\lim_{n \rightarrow \infty} \frac{1}{\phi_n(\alpha)} = 0$ , applying the Stolz-Césaro criterion, we have

$$\lim_{n \rightarrow \infty} \frac{|\phi_n(\alpha)|^2}{1 + K_n(\alpha, \alpha)} = \left( 1 - \frac{1}{|\alpha|^2} \right). \quad (5.32)$$

On the other hand, from (5.30) we can deduce the following identity

$$\frac{|\phi_n(\alpha)|^2}{1 + K_n(\alpha, \alpha)} = 1 - \frac{1 + K_{n-1}(\alpha, \alpha)}{1 + K_n(\alpha, \alpha)} = 1 - \left( \frac{\gamma_{n;1}}{\kappa_n} \right)^2. \quad (5.33)$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

Thus,

$$\lim_{n \rightarrow \infty} \frac{K_{n-1}(0, \alpha)}{\gamma_{n;1} \phi_n(\alpha)} = 0$$

and the result follows. □

We are now in a position to summarize the results obtained above, in the following statement.

**Theorem 5.2.2.** *Let  $\{\psi_n\}_{n \geq 0}$  be the sequence of monic orthogonal polynomials associated with the inner product (5.20). Then, uniformly in  $\mathbb{C} \setminus \mathbb{D}$ ,*

$$\lim_{n \rightarrow \infty} \frac{\psi_n(z)}{\phi_n(z)} = \prod_{i=1}^m B(\alpha_i)^{l_i+1}.$$

*Proof.* First of all, we prove the result for

$$\mathbf{f}(Z) = \mathbf{f}_m(Z) = (f^{(l_1)}(\alpha_1), \dots, f^{(l_m)}(\alpha_m)),$$

and  $\mathbf{A}_m$  a positive definite hermitian matrix of order  $m$ . Let  $\{\psi_{n;m}\}_{n \geq 0}$  be the sequence of orthonormal polynomials with respect to (5.20) for  $\mathbf{f}(Z) = \mathbf{f}_m(Z)$ . We can assume, without loss of generality,  $\mathbf{A}_m = \mathbf{I}_m$  by Lemma 5.2.1. Therefore, the relative asymptotics can be written as follows

$$\lim_{n \rightarrow \infty} \frac{\psi_{n;m}(z)}{\phi_n(z)} = \lim_{n \rightarrow \infty} \frac{\psi_{n;1}(z)}{\phi_n(z)} \prod_{i=2}^m \frac{\psi_{n;i}(z)}{\psi_{n;i-1}(z)},$$

which, using Lemma 5.2.3 and Theorem 5.2.1, immediately yields

$$\lim_{n \rightarrow \infty} \frac{\psi_{n;m}(z)}{\phi_n(z)} = \prod_{i=1}^m B(\alpha_i).$$

Finally, the proof for a general  $\mathbf{f}(Z)$  is a straightforward consequence of the previous analysis. □

### 5.2.2 Zeros

In this subsection we study the asymptotic behavior of the zeros of orthogonal polynomials associated with the discrete Sobolev inner product (5.24). In contrast with the

real line case [4; 23; 61; 62; 130; 157] (see also Appendix A), there is not a well developed theory for zeros of discrete Sobolev orthogonal polynomials on the unit circle.

The monic version of (5.28) is

$$\Psi_n(z) = \Phi_n(z) - \frac{\lambda \Phi_n^{(j)}(\alpha)}{1 + \lambda K_{n-1}^{(j,j)}(\alpha, \alpha)} K_{n-1}^{(0,j)}(z, \alpha). \quad (5.34)$$

Thus,

$$\begin{bmatrix} \Psi_0(z) \\ \Psi_1(z) \\ \vdots \\ \Psi_{n-1}(z) \end{bmatrix} = \mathbf{L}_n \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix},$$

where  $\mathbf{L}_n$  is a  $n \times n$  lower triangular matrix with 1 as entries in the main diagonal, and the remaining entries are given by (5.34), i.e.,

$$l_{m,k} = -\frac{1}{\|\Phi_k\|_\sigma^2} \frac{\lambda \Phi_m^{(j)}(\alpha) \overline{\Phi_k^{(j)}(\alpha)}}{(1 + \lambda K_{m-1}^{(j,j)}(\alpha, \alpha))}, \quad 1 \leq m \leq n, \quad 0 \leq k \leq m-1.$$

### GGT matrices

One of our aim is to find a relation between  $\mathbf{H}_\Psi$ , the Hessenberg matrix associated with the monic orthogonal polynomials  $\{\Psi_n\}_{n \geq 0}$ , and  $\mathbf{H}_\sigma$ . In particular, we get

$$z \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix} = (\mathbf{H}_\sigma)_n \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix} + \Phi_n(z) \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix},$$

and, on the other hand,

$$z \begin{bmatrix} \Psi_0(z) \\ \Psi_1(z) \\ \vdots \\ \Psi_{n-1}(z) \end{bmatrix} = (\mathbf{H}_\Psi)_n \begin{bmatrix} \Psi_0(z) \\ \Psi_1(z) \\ \vdots \\ \Psi_{n-1}(z) \end{bmatrix} + \Psi_n(z) \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}. \quad (5.35)$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

Substituting in (5.35), we obtain

$$z\mathbf{L}_n \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix} = (\mathbf{H}_\Psi)_n \mathbf{L}_n \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix} + \Phi_n(z) \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix} + \mathbf{A}_n \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix},$$

where

$$\mathbf{A}_n = \begin{bmatrix} 0 & \dots & \dots & 0 \\ \vdots & & & \vdots \\ 0 & \dots & \dots & 0 \\ l_{n,0} & \dots & \dots & l_{n,n-1} \end{bmatrix}.$$

As a consequence,

$$z \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix} = (\mathbf{L}_n^{-1}(\mathbf{H}_\Psi)_n \mathbf{L}_n + \mathbf{L}_n^{-1} \mathbf{A}_n) \begin{bmatrix} \Phi_0(z) \\ \Phi_1(z) \\ \vdots \\ \Phi_{n-1}(z) \end{bmatrix} + \Phi_n(z) \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix},$$

so

$$(\mathbf{H}_\sigma)_n = \mathbf{L}_n^{-1}(\mathbf{H}_\Psi)_n \mathbf{L}_n + \mathbf{L}_n^{-1} \mathbf{A}_n$$

and therefore, since

$$\mathbf{L}_n^{-1} \mathbf{A}_n = \mathbf{A}_n,$$

we have

**Theorem 5.2.3.** *Let  $(\mathbf{H}_\sigma)_n$  and  $(\mathbf{H}_\Psi)_n$  be the  $n \times n$  truncated GGT matrices associated with  $\{\Phi_n\}_{n \geq 0}$  and  $\{\Psi_n\}_{n \geq 0}$ , respectively. Then,*

$$(\mathbf{H}_\Psi)_n = \mathbf{L}_n((\mathbf{H}_\sigma)_n - \mathbf{A}_n)\mathbf{L}_n^{-1}.$$

*As a consequence, the zeros of  $\Psi_{n+1}$  are the eigenvalues of the matrix  $(\mathbf{H}_\sigma)_n - \mathbf{A}_n$ , a rank one perturbation of the matrix  $(\mathbf{H}_\sigma)_n$ .*

In the previous theorem we have characterized the eigenvalues of the GGT matrix associated with the discrete Sobolev polynomials as the eigenvalues of a rank one perturbation of the GGT matrix associated with the measure.

Notice that  $\mathbf{A}_n = (0, \dots, 0, 1)^T (l_{n,0}, l_{n,1}, \dots, l_{n,n-1})$  and, since  $l_{n,k} = 0$  for  $k < j$ , then

$$\mathbf{A}_n = \frac{\lambda \Phi_n^{(j)}(\alpha)}{1 + \lambda K_{n-1}^{(j,j)}(\alpha, \alpha)} \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix} \begin{bmatrix} 0, \dots, 0, \frac{\overline{\Phi_j^{(j)}(\alpha)}}{\|\Phi_j\|^2}, \dots, \frac{\overline{\Phi_{n-1}^{(j)}(\alpha)}}{\|\Phi_{n-1}\|^2} \end{bmatrix}.$$

As an example, if  $d\sigma(\theta) = \frac{d\theta}{2\pi}$  is the Lebesgue measure, it is not difficult to see that in such a case, if  $\alpha = 0$ , then  $\mathbf{A}_n = 0$ ,  $n \neq j$ , and

$$\mathbf{A}_j = \frac{\lambda(j!)^2}{1 + \lambda(j!)^2} \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix} [0, \dots, 0, 1, 0, \dots, 0],$$

where the one is in the position  $j$ . On the other hand, if  $\alpha = 1$ , then for  $n \geq j$ ,

$$\mathbf{A}_n = \frac{\lambda \frac{(n)!}{(n-j)!}}{1 + \lambda \sum_{k=j}^{n-1} \left( \frac{k!}{(k-j)!} \right)^2} \begin{bmatrix} 0 \\ \vdots \\ 0 \\ 1 \end{bmatrix} \left[ 0, \dots, 0, j!, (j+1)!, \dots, \frac{(n-1)!}{(n-j-1)!} \right].$$

### Asymptotic behavior

First, denote by  $\{\phi_n(\cdot; d\sigma_{j+1})\}_{n \geq 0}$  the corresponding sequence of orthonormal polynomial with respect to

$$d\sigma_j(z) = |z - \alpha|^{2(j+1)} d\sigma(z), \quad j \geq 0,$$

i.e., the product of  $j+1$  transformations as (5.16). For any  $j \geq 0$ , the relation between  $\phi_n(\cdot; d\sigma_{j+1})$  and  $\phi_n(\cdot; d\sigma)$  is given by [126]

$$(z - \alpha)^{j+1} \phi_{n-j-1}(z, d\sigma_{j+1}) = \frac{\eta_{n-j-1}}{\alpha_n} \left( \phi_n(z) - \sum_{k=0}^j \gamma_{n,k} K_{n-1}^{(0,k)}(z, \alpha) \right), \quad (5.36)$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

---

where  $\eta_n$  is the leading coefficient of  $\phi_n(\cdot, d\sigma_{j+1})$ , and  $\gamma_{n,k}$  is the  $k$ -th component of the vector

$$\begin{bmatrix} \phi_n(\alpha) & \phi'_n(\alpha) & \dots & \phi_n^{(j)}(\alpha) \end{bmatrix} \begin{bmatrix} K_{n-1}(\alpha, \alpha) & K_{n-1}^{(0,1)}(\alpha, \alpha) & \dots & K_{n-1}^{(0,j)}(\alpha, \alpha) \\ K_{n-1}^{(1,0)}(\alpha, \alpha) & K_{n-1}^{(1,1)}(\alpha, \alpha) & \dots & K_{n-1}^{(1,j)}(\alpha, \alpha) \\ \vdots & \vdots & \ddots & \vdots \\ K_{n-1}^{(j,0)}(\alpha, \alpha) & K_{n-1}^{(j,1)}(\alpha, \alpha) & \dots & K_{n-1}^{(j,j)}(\alpha, \alpha) \end{bmatrix}^{-1}.$$

If  $\sigma \in \mathcal{N}$ , then [126]

$$\lim_{n \rightarrow \infty} \frac{\phi_n(z; d\sigma_{j+1})}{\phi_{n+j+1}(z)} = \left( \frac{\bar{\alpha}}{|\alpha|} \frac{1}{\bar{\alpha}z - 1} \right)^{j+1}, \quad (5.37)$$

holds uniformly in  $|z| > 1$  if  $|\alpha| \geq 1$ , and in  $|z| \geq 1$  if  $|\alpha| > 1$ .

On the other hand, by Theorem 5.2.1, for  $|\alpha| > 1$ ,

$$\lim_{n \rightarrow \infty} \frac{\psi_n(z)}{\phi_n(z)} = \frac{\bar{\alpha}}{|\alpha|} \frac{z - \alpha}{\bar{\alpha}z - 1}, \quad (5.38)$$

uniformly on every compact subset of  $|z| > 1$ . From (5.37) and (5.38), we have

$$\left( \frac{|\alpha|}{\bar{\alpha}} (\bar{\alpha}z - 1) \right)^j (z - \alpha) \lim_{n \rightarrow \infty} \frac{\phi_n(z; d\sigma_{j+1})}{\phi_{n+j+1}(z)} = \lim_{n \rightarrow \infty} \frac{\psi_{n+j+1}(z)}{\phi_{n+j+1}(z)}.$$

Hence,

$$\lim_{n \rightarrow \infty} \frac{\psi_n(z)}{\phi_{n-j-1}(z; d\sigma_{j+1})} = \left( \frac{|\alpha|}{\bar{\alpha}} (\bar{\alpha}z - 1) \right)^j (z - \alpha),$$

uniformly  $|z| \geq 1$ . The following result follows immediately from Hurwitz's Theorem [49].

**Theorem 5.2.4.** *There is a positive integer  $n_0$  such that, for  $n \geq n_0$ , the  $n$ -th Sobolev monic orthogonal polynomial  $\Psi_n$  defined by (5.24), with  $|\alpha| > 1$ , has exactly 1 zero in  $\mathbb{C} \setminus \bar{\mathbb{D}}$  accumulating in  $\alpha$ , while the remaining zeros belong to  $\mathbb{D}$ .*

This result is analogous to the well known result of Meijer [140] for Sobolev orthogonal polynomials on the real line; see also Appendix A. We now turn our attention to the case when  $\lambda$  tends to infinity. For a fixed  $n$ ,  $j = 0$  and  $\lambda$  tends to infinity,  $n - 1$  zeros of  $\psi_n$  tend to the zeros of  $\phi_{n-1}(z, d\sigma_1)$ , and the remaining zero tends to  $z = \alpha$ . On the other hand, for  $j = 1$ , the zeros of  $\psi_n$  tend to the zeros of a linear combination of



---

$\Phi_n(z)$ ,  $(z - \alpha)\Phi_{n-1}(z, d\sigma_1)$ , and  $(z - \alpha)^2\Phi_{n-2}(z, d\sigma_2)$  when  $\lambda \rightarrow \infty$ . This result can be generalized for arbitrary  $j$ . Indeed, from (5.36), notice that

$$-\gamma_{n,j}K_{n-1}^{(0,j)}(z, \alpha) = (z - \alpha)^{j+1}\phi_{n-j-1}(z, d\sigma_{j+1}) - \frac{\eta_{n-j-1}}{\alpha_n}\phi_n(z) + \sum_{k=0}^{j-1}\gamma_{n,k}K_{n-1}^{(0,k)}(z, \alpha).$$

Applying the last formula recursively for  $k = 0, 1, \dots, j - 1$ , we obtain

**Theorem 5.2.5.** *Let  $\{\psi_n\}_{n \geq 0}$  be the sequence of orthonormal polynomials with respect to (5.24), with  $j \geq 0$ . Then  $\psi_n(z)$  is a linear combination of  $\phi_n(z)$ ,  $(z - \alpha)\phi_{n-1}(z, d\sigma_1)$ ,  $\dots$ ,  $(z - \alpha)^{j+1}\phi_{n-j-1}(z, d\sigma_{j+1})$ . As a consequence, the zeros of  $\psi_n(z)$  tend to the zeros of such a linear combination when  $\lambda \rightarrow \infty$ .*

Now, we provide an extremal characterization for the limit discrete Sobolev polynomials, when the mass tends to infinity. Notice that when  $\lambda$  tends to infinity in (5.34), we get the limit polynomial

$$\widetilde{\Psi}_n(z) = \Phi_n(z) - \frac{\Phi_n^{(j)}(\alpha)}{K_{n-1}^{(j,j)}(\alpha, \alpha)}K_{n-1}^{(0,j)}(z, \alpha). \quad (5.39)$$

It is easily seen that  $\widetilde{\Psi}_n^{(j)}(\alpha) = 0$ , as well as  $\widetilde{\Psi}_n$  is orthogonal to the linear space  $\text{span}\{1, z - \alpha, \dots, (z - \alpha)^{j-1}, (z - \alpha)^{j+1}, \dots, (z - \alpha)^{n-1}\} \in \mathbb{P}_n$ . Assume that  $\widehat{\Psi}_n$  is a monic polynomial of degree  $n$  such that  $\widehat{\Psi}_n^{(j)}(\alpha) = 0$ . Then, we can write

$$\widehat{\Psi}_n(z) = \Phi_n(z) + \sum_{k=0}^{n-1}\lambda_{n,k}\phi_k(z), \quad (5.40)$$

for some (unique) complex numbers  $\lambda_{n,k}$ , and therefore

$$\Phi_n^{(j)}(\alpha) + \sum_{k=0}^{n-1}\lambda_{n,k}\phi_k^{(j)}(\alpha) = 0.$$

On the other hand, from Cauchy-Schwarz inequality, we get

$$\left|\Phi_n^{(j)}(\alpha)\right|^2 \leq \sum_{k=0}^{n-1}|\lambda_{n,k}|^2 \sum_{k=0}^{n-1}|\phi_k^{(j)}(\alpha)|^2 = K_{n-1}^{(j,j)}(\alpha, \alpha) \sum_{k=0}^{n-1}|\lambda_{n,k}|^2,$$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS

and taking norms with respect to  $\mu$  in (5.40), we obtain

$$\|\widehat{\Psi}_n\|_{\sigma}^2 = \|\Phi_n\|_{\sigma}^2 + \sum_{k=0}^{n-1} |\lambda_{n,k}|^2.$$

Thus,

$$\|\widehat{\Psi}_n\|_{\sigma}^2 = \|\Phi_n\|_{\sigma}^2 + \sum_{k=0}^{n-1} |\lambda_{n,k}|^2 \geq \|\Phi_n\|_{\sigma}^2 + \frac{|\Phi_n^{(j)}(\alpha)|^2}{K_{n-1}^{(j,j)}(\alpha, \alpha)}.$$

But the term in the right hand side is precisely  $\|\widetilde{\Psi}_n\|_{\sigma}^2$ . As a consequence, we have proved the following extremal characterization for the limit polynomial  $\widetilde{\Psi}_n$ .

**Theorem 5.2.6.** *Let  $\widetilde{\Psi}_n$  be the limit monic polynomial of degree  $n$  defined by (5.39), then*

$$\|\widetilde{\Psi}_n\|_{\sigma}^2 = \min \left\{ \int_{\mathbb{T}} |\widetilde{\Psi}_n(z)|^2 d\mu(z); \widetilde{\Psi}_n(z) = z^n + (\text{lower degree terms}), \widetilde{\Psi}_n^{(j)}(\alpha) = 0 \right\}.$$

### Example: Lebesgue polynomials

For the normalized Lebesgue measure, it is very well known that its corresponding monic orthogonal polynomial sequence is  $\Phi_n(z) = z^n$ ,  $n \geq 0$ . Thus,

$$\begin{aligned} \Phi_n^{(j)}(\alpha) &= \frac{n!}{(n-j)!} \alpha^{n-j}, & K_{n-1}^{(0,j)}(z, \alpha) &= \sum_{k=j}^{n-1} \frac{k!}{(k-j)!} z^k \bar{\alpha}^{k-j}, \\ K_n^{(j,j)}(\alpha, \alpha) &= \sum_{k=j}^n \left( \frac{k!}{(k-j)!} \right)^2 |\alpha|^{2(k-j)}. \end{aligned}$$

When  $n$ ,  $j$ , and  $\lambda$  are fixed and  $\alpha$  varies, we were able to identify some 'critical' values of  $\alpha$  that change the behavior of the zeros of  $\psi_n$ . Of course, such values of  $\alpha$  will depend on  $n$ ,  $j$ , and  $\lambda$ . The following table illustrates such a situation when  $n = 30$ ,  $j = 2$ , and  $\lambda = 10$ .

Figures 5.1(a) illustrates the information in Table 5.1, showing the location of the zeros of  $\psi_{30}$  for several values of  $\alpha$ . Namely, the zeros corresponding to  $\alpha = 0.4 + 0.4i$  (blue discs),  $\alpha = 0.7 + 0.7i$  (purple square),  $\alpha = 1.05 + 1.05i$  (yellow diamonds), and  $\alpha = 1.6 + 1.6i$  (green triangles) were plotted. On the other hand, Figure 5.1(b) illustrates the behavior of the zeros of  $\psi_n$  as  $n \rightarrow \infty$ . The zeros corresponding to  $n = 10$ ,  $n = 20$ ,  $n = 30$ , and  $n = 100$  were plotted. Notice that one of the zeros approaches the value of

Table 5.1: Critical values for  $n = 30$ ,  $j = 2$ , and  $\lambda = 10$

$\alpha$	Behavior of zeros
$0 <  \alpha  < 0.8202$	All zeros approximately aligned and increase with $\alpha$
$ \alpha  \sim 0.8202$	One of the zeros ( $z_i$ ) breaks the pattern
$0.8202 <  \alpha  < 1.3194$	$z_i$ increases with $\alpha$
$ \alpha  \sim 0.1.3194$	$z_i$ changes sign
$1.1394 <  \alpha  < 1.6263$	$z_i$ decreases with $\alpha$
$ \alpha  \sim 1.6263$	$z_i$ goes back to the aligned pattern
$ \alpha  > 1.6263$	All zeros approximately aligned and decrease with $\alpha$

$\alpha$  as  $n$  increases, as stated in Theorem 5.2.4.




Figure 5.1: Zeros for Lebesgue polynomials with  $j = 2$ ,  $\lambda = 10$

### Example: Bernstein-Szegő polynomials

In order to analyze the behavior of the zeros according to the location of  $\alpha$ , we present some numerical computations of such zeros for the orthogonal polynomials associated with perturbations of the form (5.24) for one special case of probability measures on the unit circle: the Bernstein-Szegő measures.

For the Bernstein-Szegő measure, see example in Section 4.2.1, we have  $\phi_n(z) =$

## 5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS




Figure 5.2: Zeros of Bernstein-Szegő polynomials with  $b = 0.8 + 0.8i$ ,  $j = 2$ , and  $\lambda = 1$

$z^{n-1}(z - \bar{b})$ ,  $n \geq 1$ . Thus, we get

$$\begin{aligned}\phi_n^{(j)}(\alpha) &= \frac{n!}{(n-j)!} \alpha^{n-j} - \bar{b} \frac{(n-1)!}{(n-j-1)!} \alpha^{n-j-1}, \\ K_{n-1}^{(0,j)}(z, \alpha) &= \sum_{k=j}^{n-1} (z^k - \bar{b} z^{k-1}) \left( \frac{k!}{(k-j)!} \bar{\alpha}^{k-j} - \bar{b} \frac{(k-1)!}{(k-j-1)!} \bar{\alpha}^{k-j-1} \right), \\ K_n^{(j,j)}(\alpha, \alpha) &= \sum_{k=j}^n \left| \frac{k!}{(k-j)!} \bar{\alpha}^{k-j} - \bar{b} \frac{(k-1)!}{(k-j-1)!} \bar{\alpha}^{k-j-1} \right|^2,\end{aligned}$$

and, we obtain again the expression of  $\psi_n$  using (5.25). We perform a similar numerical analysis of the zeros of  $\psi_{30}$  as a function of  $\alpha$  as in the Lebesgue case. Figure 5.2(a) shows the behavior of the zeros of  $\psi_{30}$  for fixed  $j$ ,  $\lambda$ , and  $b$ , and several values of  $\alpha$ , namely  $\alpha = 0.4 + 0.4i$  (blue discs),  $\alpha = 0.7 + 0.7i$  (purple square),  $\alpha = 1.08 + 1.08i$  (yellow diamonds), and  $\alpha = 2.2 + 2.2i$  (green triangles). As before, the behavior of the zeros as  $n \rightarrow \infty$  is illustrated in Figure 5.2(b), using the same values of  $n$  plotted in the Lebesgue case.

On the other hand, the behavior of the zeros of  $\psi_n$  when  $\lambda \rightarrow \infty$  is shown in Figure 5.2.2. According to Theorem 5.2.5, the zeros of  $\psi_n$  tend to a linear combination of  $\phi_n(z)$ ,  $(z - \alpha)\phi_{n-1}(z, d\mu_1)$  and  $(z - \alpha)\phi_{n-2}(z, d\mu_2)$ , when  $j = 1$ . We computed the zeros of such polynomials for the Lebesgue (Fig. 5.3(a)) and Bernstein-Szegő (Fig. 5.3(b)) cases.




Figure 5.3: Zeros of Lebesgue and Bernstein-Szegő polynomials with  $n = 8, j = 1$

The points on the outer diameter correspond to the zeros of the above mentioned linear combination. Notice that when  $\lambda$  increases, the zeros of  $\psi_n$  approach the outer diameter, as per Theorem 5.2.5.

## **5. SPECTRAL TRANSFORMATIONS ASSOCIATED WITH MASS POINTS**

## Chapter 6

# Generators of rational spectral transformations for $C$ -functions

...Stieltjes function(s)<sup>1</sup> with polynomial coefficients can be presented as a finite superposition of four fundamental elementary transforms ...

— A. Zhedanov. [202]

In this chapter we deal with transformations of sequences of orthogonal polynomials associated with the linear functional  $\mathcal{L}$  using spectral transformations of the corresponding  $C$ -function  $F$ . First, we study the modifications obtained by multiplying a hermitian functional by a polynomial of any degree, in short, polynomial modifications. We characterize when two functionals are related by a polynomial modification. We are interested in those modifications which preserve their hermitian character. The possibility of considering modifications that do not preserve the hermitian character of the functional leads to left and right orthogonality [13]. Hence, the preservation of the hermiticity calls for the use of Laurent polynomials as perturbations. Laurent polynomial modifications of hermitian linear functional have been previously considered in [28; 54; 72; 81; 82; 124; 183].

Next, we distinguish two related problems: to characterize the quasi-definiteness character of the direct polynomial modification (2.44) and the inverse polynomial mod-

---

<sup>1</sup>S- functions.

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

ification (2.45) from the original functional. Due to the non-uniqueness of the inverse problem we pay special attention to this one. Finally, we show that a linear spectral transformation of  $F$  can be obtained as a finite composition of spectral transformations (2.44)-(2.45), and also that any rational spectral transformations can be obtained as a finite composition of linear and  $\pm k$  associated spectral transformations.

### 6.1 Hermitian polynomial transformation

Let  $\mathcal{L}$  be the hermitian linear functional introduced in (2.23). The polynomial modification  $f\mathcal{L}$  is defined by

$$\langle f\mathcal{L}, g \rangle = \langle \mathcal{L}, fg \rangle, \quad f, g \in \Lambda.$$

This polynomial modification is hermitian if and only if  $f$  is a hermitian Laurent polynomial, i.e.,  $f = f_*$ , which is equivalent to state that  $f = p + p_*$ ,  $p \in \mathbb{P}$ . Such a polynomial  $p$  can be uniquely determined by  $f$  simply requiring  $p(0) \in \mathbb{R}$ .

Féjer-Riesz's Theorem [66; 166] states that any Laurent polynomial  $f$  which is non-negative on  $\mathbb{T}$  can be factorized  $f(e^{i\theta}) = |p(e^{i\theta})|^2$ , where  $p$  is a polynomial whose zeros all lie in  $\overline{\mathbb{D}}$  or, in other words, one can find  $p$  whose zeros are all in  $\mathbb{C} \setminus \mathbb{D}$ . By analyticity,  $f(z) = p(z)\overline{p(1/\bar{z})}$ . Thus,  $f\mathcal{L}$  is positive definite for a positive definite linear functional  $\mathcal{L}$  if  $f$  also satisfies Féjer-Riesz's condition. Its analog on the real line, that is,  $P(x) \geq 0$  on  $\mathbb{R}$  implies  $P(x) = Q(x)\overline{Q(\bar{x})}$  where  $Q$  has all its zeros in  $\overline{\mathbb{C}}_+$  it is well known.

Another way of characterizing a hermitian Laurent polynomial modification is through the polynomial  $g = z^{\deg p} f$  of degree  $2\deg p$ . The condition  $f = f_*$  means that  $g$  is self-reciprocal, i.e.,  $g = g^*$ . Therefore, hermitian polynomial modifications are related with self-reciprocal polynomials of even degree. The zeros  $z_i$  of a self-reciprocal polynomial lie on the unit circle or appear in symmetric pairs  $z_i, 1/\bar{z}_i$ . Indeed, this property characterizes the self-reciprocal polynomials up to numerical factors. This implies that any self-reciprocal polynomial of even degree factorizes into a product of self-reciprocal polynomials of degree 2. As a consequence, an arbitrary hermitian polynomial modification  $f\mathcal{L}$  can be factorized as a product of elementary ones of degree one. In the sequel we assume for simplicity that the polynomial modification  $f$  is a monic hermitian Laurent polynomial from which we can deduce immediately the more general case.



We denote by  $W^{\perp n}$ <sup>1</sup> the orthogonal complement in  $\mathbb{P}_n$  of a subspace  $W \subset \mathbb{P}_n$ .

**Theorem 6.1.1.** [189] *Let  $\mathcal{L}$  be a hermitian linear functional such that the corresponding sequence of monic orthogonal polynomials  $\{\Phi_n\}_{n \geq 0}$  exists. Then,*

$$\left\{z^k \Phi_n(z)\right\}_{k=1}^r \cup \left\{z^k \Phi_n^*(z)\right\}_{k=1}^r$$

*is a basis of  $(z^r \mathbb{P}_{n-r-1})^{\perp n+r}$  for  $n \geq r \geq 1$  and a generator system of  $\mathbb{P}_{n+r}$  for  $r > n \geq 0$ .*

As an immediate consequence of Theorem 6.1.1 we have the following result.

**Corollary 6.1.1.** [30] *Let  $\mathcal{L}$  be a hermitian linear functional such that there exists the corresponding sequence of monic orthogonal polynomials  $\{\Phi_n\}_{n \geq 0}$ . Then, every polynomial  $\Psi_n(z) \in (z^r \mathbb{P}_{n-r-1})^{\perp n+r}$  has a unique decomposition,  $\Psi_n(z) = C(z)\Phi_n(z) + D(z)\Phi_n^*(z)$ ,  $C \in \mathbb{P}_r$ ,  $D \in \mathbb{P}_{r-1}$ , for  $n \geq r \geq 1$ , and every polynomial  $\Psi_n \in \mathbb{P}_{n+r}$  has infinitely many decompositions for  $r > n \geq 0$ .*

To study the hermitian polynomial modifications we have the next theorem.

**Theorem 6.1.2.** [30] *Let  $\mathcal{L}$  and  $\tilde{\mathcal{L}}$  be hermitian linear functionals with finite segments of monic orthogonal polynomials  $\{\Phi_j\}_{j=0}^n$ ,  $\{\Psi_j\}_{j=0}^{n+r}$ , respectively, and let  $f(z) = p(z) + p_*(z) = z^{-r}g(z)$  with  $f$  be a polynomial of degree  $r$ . Then, the following statements are equivalent:*

i)  $\tilde{\mathcal{L}} = f\mathcal{L}$  in  $\mathbb{P}_n$ .

ii) *There exist  $C_j \in \mathbb{P}_r$ ,  $D_j \in \mathbb{P}_{r-1}$  with  $C_j(0) \neq 0$  such that*

$$g(z)\Psi_j = C_j(z)\Phi_{j+r}(z) + D_j(z)\Phi_{j+r}^*(z), \quad j \geq 0.$$

iii) *There exist  $C_j \in \mathbb{P}_r$ ,  $D_j \in \mathbb{P}_{r-1}$  with  $C_j(0) \neq 0$  such that*

$$g(z)\Psi_j^*(z) = zD_j^*(z)\Phi_{j+r}(z) + C_j^*(z)\Phi_{j+r}^*(z), \quad D_j^*(z) = D_j^{*r-1}(z), \quad j \geq 0.$$

*The polynomials  $C_j \in \mathbb{P}_r$ ,  $D_j \in \mathbb{P}_{r-1}$  satisfying ii) or iii) are unique,  $\deg C_j = r$ ,  $C_j(0) \in \mathbb{R}$  and  $C_j^*(0) = g(0)$ .*

The above theorem has the following consequence for quasi-definite functionals.

<sup>1</sup>The orthogonal complement  $W^{\perp n}$  of a subspace  $W$  of an inner product space  $V$  is the set of all vectors in  $V$  that are orthogonal to every vector in  $W$ .

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

**Corollary 6.1.2.** [30] *Let  $\mathcal{L}$  and  $\tilde{\mathcal{L}}$  be quasi-definite hermitian linear functionals with sequence of monic orthogonal polynomials  $\{\Phi_n\}_{n \geq 0}$ ,  $\{\Psi_n\}_{n \geq 0}$ , respectively, and let  $f(z) = p(z) + p(z)_* = z^{-r}g(z)$  with  $f$  be a polynomial of degree  $r$ . Then,  $\mathcal{L} = p\tilde{\mathcal{L}}$  if and only if there exist polynomials  $C_n \in \mathbb{P}_r$ ,  $D_n \in \mathbb{P}_{r-1}$  with  $C_n(0) \neq 0$  such that*

$$g(z)\Psi_n(z) = C_n(z)\Phi_{n+r}(z) + D_n(z)\Phi_{n+r}^*(z), \quad n \geq 0,$$

or, equivalently,

$$g(z)\Psi_n^*(z) = zD_n^*(z)\Phi_{n+r}(z) + C_n^*(z)\Phi_{n+r}^*(z), \quad n \geq 0.$$

### 6.2 Laurent polynomial transformation: Direct problem

The direct problem deals with the case where we suppose that a hermitian linear functional  $\mathcal{L}$  associated with the sequence of orthogonal polynomials  $\{\Phi_n\}_{n \geq 0}$ , and a hermitian polynomial of degree 1 are given, i.e., we consider the spectral transformation (2.44). Then, we obtain information about the functional  $\mathcal{L}_R$  and its sequence of monic orthogonal polynomials  $\{\Psi_n\}_{n \geq 0}$ .

From (2.45), we get

$$\tilde{c}_{-k} = c_{-(k+1)} + c_{-(k-1)} - (\alpha + \bar{\alpha})c_{-k}, \quad (6.1)$$

where  $\{\tilde{c}_n\}_{n \geq 0}$  is the sequence of moments associated with  $\mathcal{L}_R$ . Notice that if  $a = -2\Re(\alpha)$ , and using (6.1), we obtain

$$\tilde{\mathbf{T}} = \mathbf{Z}\mathbf{T} + a\mathbf{T} + \mathbf{T}\mathbf{Z}^T,$$

where  $\tilde{\mathbf{T}}$  is the Toeplitz matrix associated with  $\mathcal{L}_R$ , and  $\mathbf{Z}$  is the shift matrix with ones on the first upper-diagonal and zeros on the remaining entries.

#### 6.2.1 Regularity conditions and Verblunsky coefficients

If  $\mathcal{L}$  is quasi-definite, necessary and sufficient conditions for  $\mathcal{L}_R$  to be also quasi-definite have been studied. Moreover, the explicit expression for sequences of monic polynomials orthogonal with respect to  $\mathcal{L}_R$  have been obtained [28; 183].

---

**Theorem 6.2.1.** [183]

i) If  $|\Re\alpha| \neq 1$ , and  $b_1, b_2$  are zeros of the polynomial  $z^2 - (\alpha + \bar{\alpha})z + 1$ , then  $\mathcal{L}_R$  is quasi-definite if and only if  $K_n^*(b_1, b_2) \neq 0$ ,  $n \geq 0$ . In addition, if  $\{\Psi_n\}_{n \geq 0}$  denotes the sequence of monic polynomials orthogonal with respect to  $\mathcal{L}_R$ , then

$$\Psi_{n-1}(z) = \frac{\Phi_n(z)K_{n-1}^*(b_1, b_2) - K_{n-1}^*(z, b_2)\Phi_n(b_1)}{K_{n-1}^*(b_1, b_2)(z - b_1)}, \quad n \geq 1, \quad (6.2)$$

and

$$\Psi_{n-1}(0) = \frac{\Phi_n(b_1)\Phi_{n-1}(b_2) - \Phi_n(b_2)\Phi_{n-1}(b_1)}{K_{n-1}^*(b_1, b_2)(b_1 - b_2)\mathbf{k}_{n-1}}, \quad n \geq 1. \quad (6.3)$$

ii) If  $|\Re\alpha| = 1$ , and  $b$  is the double zero of the polynomial  $z^2 - (\alpha + \bar{\alpha})z + 1$ , then  $\mathcal{L}_R$  is quasi-definite if and only if  $K_n^*(b, b) \neq 0$ ,  $n \geq 0$ . In addition,

$$\Psi_{n-1}(z) = \frac{\Phi_n(z)K_{n-1}^*(b, b) - K_{n-1}^*(z, b)\Phi_n(b)}{K_{n-1}^*(b, b)(z - b)}, \quad n \geq 1, \quad (6.4)$$

and

$$\Psi_{n-1}(0) = -b \frac{\Phi_n(0)K_{n-1}^*(b, b)\mathbf{k}_{n-1} - \Phi_{n-1}(b)\Phi_n(b)}{K_{n-1}^*(b, b)\mathbf{k}_{n-1}}, \quad n \geq 1. \quad (6.5)$$

According to Corollary 6.1.2, (6.2) and (6.4) can be written as follows.

**Corollary 6.2.1.**

i) If  $|\Re\alpha| \neq 1$ , and  $b_1, b_2$  are the zeros of the polynomial  $z^2 - (\alpha + \bar{\alpha})z + 1$ , then

$$\begin{aligned} (z - b_1)(z - b_2)\Psi_n(z) &= z \left( z - b_2 - \frac{\Phi_{n+1}(b_1)\Phi_n^*(b_2)}{\mathbf{k}_n K_n^*(b_1, b_2)} \right) \Phi_n(z) \\ &\quad + \left( (z - b_2)\Phi_{n+1}(0) + \frac{b_2\Phi_{n+1}(b_1)\Phi_n(b_2)}{\mathbf{k}_n K_n^*(b_1, b_2)} \right) \Phi_n^*(z). \end{aligned}$$

ii) If  $|\Re\alpha| = 1$ , and  $b$  is the double zero of the polynomial  $z^2 - (\alpha + \bar{\alpha})z + 1$ , then

$$\begin{aligned} (z - b)^2\Psi_n(z) &= z \left( z - b - \frac{\Phi_{n+1}(b)\Phi_n^*(b)}{\mathbf{k}_n K_n^*(b, b)} \right) \Phi_n(z) \\ &\quad + \left( (z - b)\Phi_{n+1}(0) + \frac{b\Phi_{n+1}(b)\Phi_n(b)}{\mathbf{k}_n K_n^*(b, b)} \right) \Phi_n^*(z). \end{aligned}$$

There is another equivalent condition for the quasi-definiteness of  $\mathcal{L}_R$  and, consequently, an expression for the corresponding Verblunsky coefficients.

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

**Theorem 6.2.2.** [28] *The linear functional  $\mathcal{L}_R$  is quasi-definite if and only if  $\Pi_n(b_1) \neq 0$ ,  $n \geq 0$ , where*

$$\Pi_n(x) = \begin{vmatrix} x\Phi_n(x) & \Phi_n^*(x) \\ x^{-1}\Phi_n(x^{-1}) & \Phi_n^*(x^{-1}) \end{vmatrix}.$$

Moreover, the families of Verblunsky coefficients  $\{\Psi_n(0)\}_{n \geq 1}$  are given by

$$Y_n(0) = (b_1 - b_1^{-1}) \frac{\Phi_n(b_1)\Phi_n(b_1^{-1})}{\Pi_n(b_1)}, \quad n \geq 1. \quad (6.6)$$

In [72] it was obtained the relation between the corresponding GGT matrices as well as an explicit expression for the Verblunsky coefficients of associated with this perturbation is given. Thus, the invariance of the Szegő class of bounded variation measures follows.

### 6.2.2 C-functions

From the relation between the moments (6.1) we get that the C-function associated with  $\mathcal{L}_R$  is

$$F_R(z) = \frac{1}{2} \left( (z + z^{-1} - (\alpha + \bar{\alpha})) F(z) + c_0 (z - z^{-1}) + c_1 - c_{-1} \right). \quad (6.7)$$

In a more general situation, if we consider a finite composition of  $\mathcal{L}_R$  with order  $k \geq 0$  defined by

$$\mathcal{L}_{R^{(k)}} = \mathfrak{R} \left( \prod_{i=1}^k (z - \alpha_i) \right) \mathcal{L}, \quad \alpha_i \in \mathbb{C}, \quad (6.8)$$

we can prove the following result.

**Lemma 6.2.1.** *A spectral transformation given by  $\widetilde{F}(z) = A(z)F(z) + B(z)$  such that  $A$  and  $B$  provide a true asymptotic behavior to (2.43) is equivalent to (6.8). Furthermore,*

$$A(z) = \frac{1}{2} \mathfrak{R} p(z), \quad B(z) = P(z) - P_*(z), \quad (6.9)$$

where  $P$  is the polynomial of second kind associated with

$$p(z) = \prod_{i=1}^k (z - \alpha_i)$$

with respect to the linear functional (6.8).

---

*Proof.* From (6.7) we get that  $\mathcal{L}_R$  is a spectral transformation with  $C = 0$  and  $D = 1$ . On the other hand, if we start with a generic spectral transformation where  $C = 0$ ,  $B$  is a hermitian Laurent polynomial of degree one, and  $D = 1$ ,  $A$  should be a hermitian Laurent polynomial of degree one (see Section 6.4), and the only choice for this spectral transformation is (6.7) containing one free parameter  $\alpha$ .

If we consider the linear functional (6.8), from (6.7) we have

$$\widetilde{F}(z) = A(z)F(z) + B(z), \quad (6.10)$$

where

$$A(z) = (z + z^{-1} - (\alpha_1 + \bar{\alpha}_1)) \cdots (z + z^{-1} - (\alpha_k + \bar{\alpha}_k))$$

is a hermitian Laurent polynomial of degree  $k$ . This transformation contains  $k$  free parameters  $\alpha_k$ .  $B$  is a polynomial of the same type and degree as  $A$ .

Conversely, if we start with a generic spectral transformation where

$$A = (z + z^{-1} - (\alpha_1 + \bar{\alpha}_1)) \cdots (z + z^{-1} - (\alpha_k + \bar{\alpha}_k))$$

is a hermitian Laurent polynomial of degree  $k$ ,  $C \equiv 0$ , and  $D \equiv 1$ , then  $B$  should be a hermitian Laurent polynomial of same degree as  $A$ , in order to satisfy  $A \equiv A_*$  (see Section 6.4). Moreover, it is easily seen that for (6.8) the polynomial  $B$  is uniquely determined by means of the sequence of moments  $\{c_n\}_{n \in \mathbb{Z}}$  associated with the linear functional  $\mathcal{L}$ . Furthermore, (6.9) follows immediately from the definition of (6.8).  $\square$

### 6.3 Laurent polynomial transformations: Inverse problem

In this section we study the inverse polynomial modification given by (2.45). More precisely, given a hermitian functional  $\mathcal{L}$  whose corresponding sequence of monic orthogonal polynomials is denoted by  $\{\Phi_n\}_{n \geq 0}$  and a hermitian Laurent polynomial of degree 1, we obtain information about the hermitian solutions  $\mathcal{L}_{R^{(-1)}}$  of (2.45) and their sequence of monic orthogonal polynomials  $\{\Psi_n\}_{n \geq 0}$ . If  $\mathcal{L}$  is a positive definite linear functional, necessary and sufficient conditions in order  $\mathcal{L}_{R^{(-1)}}$  to be a quasi-definite linear functional are given. The relation between the corresponding sequences of monic orthogonal polynomials is presented. We also obtain the relations between the corre-

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

sponding  $C$ -functions in such a way that a linear spectral transformation appears.

We can describe the hermitian solutions of  $\mathcal{L}_{R^{(-1)}}$  starting from a particular one  $\mathcal{L}_{R_0^{(-1)}}$  [30]. This approach shows that the inverse problem is related to the study of the influence of Dirac's deltas and their derivatives on the quasi-definiteness and the sequence of monic orthogonal polynomials of a hermitian functional.

We are only interested in those values of  $\alpha$  such that  $0 < |\Re(\alpha)| < 1$ . However, in this case the zeros  $b$  and  $\bar{b}$  of  $z^2 - (\alpha + \bar{\alpha})z + 1$  are complex conjugate and, furthermore,  $|b| = 1$ . We denote by  $\sigma$  and  $\sigma_R^{(-1)}$  the measures associated with the positive definite case of  $\mathcal{L}$  and  $\mathcal{L}_{R^{(-1)}}$ , respectively, i.e.,

$$d\sigma_{R^{(-1)}}(z) = \frac{d\sigma(z)}{2\Re(z - \alpha)} + m_1\delta(z - b) + m_2\delta(z - \bar{b}), \quad m_1, m_2 \in \mathbb{R}.$$

Here,  $\sigma$  is a non-trivial probability measure supported on  $\mathbb{T}$ , which can be decomposed as in (2.37). Thus, if  $\sigma_s = 0$ , then the integral

$$\tilde{c}_n = \int_0^{2\pi} \frac{e^{in\theta} \omega(\theta)}{z + z^{-1} - (\alpha + \bar{\alpha})} \frac{d\theta}{2\pi} = \frac{1}{2\pi i} \int_{\mathbb{T}} \frac{z^n \omega(z)}{z^2 - (\alpha + \bar{\alpha})z + 1} dz,$$

has singularities in  $z = b$  and  $z = \bar{b}$ . These singularities can be removed if we consider

$$\begin{aligned} \tilde{c}_n &= \frac{1}{2\pi i} \int_{\mathbb{T}} \frac{z^n \omega(\theta)}{z^2 - (\alpha + \bar{\alpha})z + 1} dz \\ &= \frac{1}{2\pi i(b - \bar{b})} \left( \int_{\mathbb{T}} \frac{z^n (\omega(z) - \omega(b))}{z - b} dz - \int_{\mathbb{T}} \frac{z^n (\omega(z) - \omega(\bar{b}))}{z - \bar{b}} dz + \frac{1}{2} b^n \omega(b) - \frac{1}{2} \bar{b}^n \omega(\bar{b}) \right), \end{aligned} \quad (6.11)$$

assuming that  $\omega \in C^{2+}$ ; that is,  $\omega$  satisfies the Lipschitz condition<sup>1</sup> of order  $\tau$  ( $0 < \tau \leq 1$ ) on  $\mathbb{T}$  [195]. Notice that this is also valid if  $\sigma_s \neq 0$ , as long as  $\sigma_s$  has a finite number of mass points different from  $b$  and  $\bar{b}$ .

From (2.45) we get

$$c_{-k} = \tilde{c}_{-(k+1)} + \tilde{c}_{-(k-1)} - (\alpha + \bar{\alpha})\tilde{c}_{-k}. \quad (6.12)$$

If  $a = -2\Re(\alpha)$ , and using (6.12), we obtain

$$\mathbf{T} = \mathbf{Z}\tilde{\mathbf{T}} + a\tilde{\mathbf{T}} + \tilde{\mathbf{T}}\mathbf{Z}^T$$

<sup>1</sup> $|\omega(z) - \omega(y)| \leq C|x - y|^\tau$ ,  $C > 0$ ,  $z, y \in \mathbb{T}$ .

Furthermore, the hermitian Toeplitz matrices can be characterized as  $\mathbf{T} = \mathbf{T}^*$  together with  $\mathbf{Z}\mathbf{T}\mathbf{Z}^T = \mathbf{T}$ , and, therefore,

$$\mathbf{T}\mathbf{Z}^T = \widetilde{\mathbf{T}}\mathbf{B},$$

where  $\mathbf{B} = \mathbf{I} + a\mathbf{Z}^T + (\mathbf{Z}^T)^2$  is an infinite lower triangular matrix with ones in the main diagonal, with the following structure

$$\mathbf{B} = \left[ \begin{array}{c|c|c} \mathbf{A} & 0 & \dots \\ \hline \mathbf{A}^T & \mathbf{A} & \ddots \\ \hline 0 & \ddots & \ddots \end{array} \right],$$

where  $\mathbf{A} = \begin{bmatrix} 1 & 0 \\ a & 1 \end{bmatrix}$ . On the other hand, is not difficult to show that

$$\mathbf{B}^{-1} = \left[ \begin{array}{c|c|c|c} \mathbf{A}_1 & 0 & 0 & \dots \\ \hline \mathbf{A}_2 & \mathbf{A}_1 & 0 & \ddots \\ \hline \mathbf{A}_3 & \mathbf{A}_2 & \mathbf{A}_1 & \ddots \\ \hline \vdots & \ddots & \ddots & \ddots \end{array} \right],$$

where  $\mathbf{A}_1 = \mathbf{A}^{-1}$ ,  $\mathbf{A}_k = (-1)^{k-1} \mathbf{A}^{-1} \mathbf{M}^{k-1}$ ,  $k \geq 2$ , and

$$\mathbf{M} = \mathbf{A}^{-1} \mathbf{A}^T = \begin{bmatrix} 1 & a \\ -a & 1 - a^2 \end{bmatrix}.$$

In other words,  $\mathbf{B}^{-1}$  is a lower triangular block matrix, with Toeplitz structure. Finally,

$$\mathbf{T}\mathbf{S} = \widetilde{\mathbf{T}},$$

where  $\mathbf{S}$  is given by

$$\mathbf{S} = \mathbf{Z}^T \mathbf{B}^{-1} = \left[ \begin{array}{c|c|c|c} \mathbf{Z}_1^T & 0 & 0 & \dots \\ \hline \mathbf{Z}_1 & \mathbf{Z}_1^T & 0 & \ddots \\ \hline 0 & \mathbf{Z}_1 & \mathbf{Z}_1^T & \ddots \\ \hline \vdots & \ddots & \ddots & \ddots \end{array} \right] \left[ \begin{array}{c|c|c|c} \mathbf{A}_1 & 0 & 0 & \dots \\ \hline \mathbf{A}_2 & \mathbf{A}_1 & 0 & \ddots \\ \hline \mathbf{A}_3 & \mathbf{A}_2 & \mathbf{A}_1 & \ddots \\ \hline \vdots & \ddots & \ddots & \ddots \end{array} \right],$$

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

with  $\mathbf{Z}_1 = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ , i.e.,  $\mathbf{S}$  is also a lower triangular block matrix with Toeplitz structure.

### 6.3.1 Regularity conditions and Verblunsky coefficients

Assume that  $\mathcal{L}_{R^{(-1)}}$  is quasi-definite and let  $\{\Psi_n\}_{n \geq 0}$  be its corresponding sequence of monic orthogonal polynomials with leading coefficients  $\tilde{k}_n$ . We next state the relation between  $\{\Psi_n\}_{n \geq 0}$  and  $\{\Phi_n\}_{n \geq 0}$ .

**Theorem 6.3.1.** *Let  $\mathcal{L}$  be a positive definite linear functional. If  $\mathcal{L}_{R^{(-1)}}$ , given as in (2.45), is a quasi-definite linear functional, then  $\Psi_n$ , the  $n$ -th monic polynomial orthogonal with respect to  $\mathcal{L}_{R^{(-1)}}$ , is*

$$\Psi_n(z) = \left( z + \frac{\tilde{k}_n}{k_{n-1}} \right) \Phi_{n-1}(z) + \left( \Phi_n(0) - \frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0) \right) \Phi_{n-1}^*(z). \quad (6.13)$$

Conversely, if  $\{\Psi_n\}_{n \geq 0}$  is given by (6.13) and assuming that  $|\Psi_n(0)| \neq 1$ ,  $n \geq 1$ , then  $\{\Psi_n\}_{n \geq 0}$  is the sequence of monic polynomials orthogonal with respect to  $\mathcal{L}_{R^{(-1)}}$ .

*Proof.* Let

$$\Psi_n(z) = \Phi_n(z) + \sum_{m=0}^{n-1} \lambda_{n,m} \Phi_m(z). \quad (6.14)$$

Multiplying the above expression by  $\overline{\Phi_m}$  and applying  $\mathcal{L}$ , for  $0 \leq m \leq n-1$ , we get

$$\langle \mathcal{L}, \Psi_n \overline{\Phi_m} \rangle = \lambda_{n,m} k_m,$$

or, equivalently,

$$\langle \mathcal{L}_{R^{(-1)}}, (z + z^{-1} - (\alpha + \bar{\alpha})) \Psi_n(z) \overline{\Phi_m(z)} \rangle = \lambda_{n,m} k_m.$$

Thus,

$$\lambda_{n,m} = \frac{1}{k_m} \langle \mathcal{L}_{R^{(-1)}}, (z + z^{-1} - (\alpha + \bar{\alpha})) \Psi_n(z) \overline{\Phi_m(z)} \rangle, \quad 0 \leq m \leq n-1.$$



If  $m = n - 1$ , then

$$\begin{aligned}\lambda_{n,n-1} &= \frac{1}{k_{n-1}} \left( \langle \mathcal{L}_{R^{(-1)}}, z \Psi_n(z) \overline{\Phi_{n-1}(z)} \rangle + \langle \mathcal{L}_{R^{(-1)}}, z^{-1} \Psi_n(z) \overline{\Phi_{n-1}(z)} \rangle \right) \\ &= \frac{\tilde{k}_n}{k_{n-1}} \left( 1 - \Psi_{n+1}(0) \overline{\Phi_{n-1}(0)} \right).\end{aligned}$$

On the other hand, for  $0 \leq m \leq n - 2$ ,

$$\lambda_{n,m} = \frac{1}{k_m} \langle \mathcal{L}_{R^{(-1)}}, z \Psi_n(z) \overline{\Phi_m(z)} \rangle = -\frac{\tilde{k}_n}{k_m} \Psi_{n+1}(0) \overline{\Phi_m(0)}.$$

Substituting these values into (6.14), we obtain

$$\begin{aligned}\Psi_n(z) &= \Phi_n(z) + \frac{\tilde{k}_n}{k_{n-1}} \Phi_{n-1}(z) - \tilde{k}_n \Psi_{n+1}(0) \sum_{m=0}^{n-1} \frac{\overline{\Phi_m(0)} \Phi_m(z)}{k_m} \\ &= \Phi_n(z) + \frac{\tilde{k}_n}{k_{n-1}} \Phi_{n-1}(z) - \frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0) \Phi_{n-1}^*(z).\end{aligned}\tag{6.15}$$

Using the recurrence relation, we get

$$\begin{aligned}\Psi_n(z) &= z \Phi_{n-1}(z) + \Phi_n(0) \Phi_{n-1}^*(z) + \frac{\tilde{k}_n}{k_{n-1}} \Phi_{n-1}(z) - \frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0) \Phi_{n-1}^*(z) \\ &= \left( z + \frac{\tilde{k}_n}{k_{n-1}} \right) \Phi_{n-1}(z) + \left( \Phi_n(0) - \frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0) \right) \Phi_{n-1}^*(z),\end{aligned}\tag{6.16}$$

which proves the first statement of the theorem.

Notice that evaluating (6.15) at  $z = 0$ , we get

$$\Psi_n(0) = \frac{\tilde{k}_n}{k_{n-1}} \Phi_{n-1}(0) + \Phi_n(0) - \frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0),\tag{6.17}$$

and thus (6.16) becomes

$$\Psi_n(z) = \left( z + \frac{\tilde{k}_n}{k_{n-1}} \right) \Phi_{n-1}(z) + \left( \Psi_n(0) - \frac{\tilde{k}_n}{k_{n-1}} \Phi_{n-1}(0) \right) \Phi_{n-1}^*(z).$$

If we denote  $\nu_n = \tilde{k}_{n+1}/k_n$  and  $l_n = \Psi_{n+1}(0) - \nu_n \Phi_n(0)$ , considering the reversed

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

polynomial of  $\Psi_{n+1}$ , we obtain the following linear transfer equation

$$\begin{bmatrix} \Psi_{n+1}(z) \\ \Psi_{n+1}^*(z) \end{bmatrix} = \begin{bmatrix} z + \nu_n & l_n \\ \bar{l}_n z & \nu_n z + 1 \end{bmatrix} \begin{bmatrix} \Phi_n(z) \\ \Phi_n^*(z) \end{bmatrix}.$$

Notice that the determinant of the above transfer matrix is

$$\begin{aligned} (z + \nu_n)(\nu_n z + 1) - |l_n|^2 z &= \nu_n z^2 + (\nu_n^2 + 1 - |l_n|^2)z + \nu_n \\ &= \nu_n(z^2 + 1) + (\nu_n^2(1 - |\Phi_n(0)|^2) + 1 - |\Psi_{n+1}(0)|^2)z \\ &\quad + \nu_n(\Psi_{n+1}(0)\overline{\Phi_n(0)} + \overline{\Psi_{n+1}(0)}\Phi_n(0))z \\ &= \nu_n(z^2 - (\alpha + \bar{\alpha})z + 1), \end{aligned}$$

where the last equality becomes clear looking at (6.18) in the following theorem. Furthermore, we get

$$\Phi_n(z) = \frac{(\nu_n z + 1)\Psi_{n+1}(z) - l_n \Psi_{n+1}^*(z)}{\nu_n(z^2 - (\alpha + \bar{\alpha})z + 1)}, \quad \Phi_n^*(z) = \frac{(z + \nu_n)\Psi_{n+1}^*(z) - \bar{l}_n z \Psi_{n+1}(z)}{\nu_n(z^2 - (\alpha + \bar{\alpha})z + 1)},$$

and thus we obtain the following alternative expression that relates both sequences of polynomials

$$(z^2 - (\alpha + \bar{\alpha})z + 1)\Phi_n(z) = \Psi_{n+2}(z) + \nu_n^{-1}(\Psi_{n+1}(z) - \Phi_{n+1}(0)\Psi_{n+1}^*(z)).$$

We now prove that the sequence of monic polynomials  $\{\Psi_n\}_{n \geq 0}$  given in (6.13) is orthogonal with respect to  $\mathcal{L}_{R^{(-1)}}$ . Notice that  $\Psi_{n+1}(z) - \Psi_{n+1}(0)\Psi_n^*(z)$  is a polynomial of degree  $n+1$  that vanishes at  $z=0$  and, thus,  $\Psi_{n+1}(z) - \Psi_{n+1}(0)\Psi_n^*(z) = zp(z)$ , where  $p(z)$  is a polynomial of degree  $n$ . Then,

$$\begin{aligned} zp(z) &= (z + \nu_n)\Phi_n(z) + l_n \Phi_n^*(z) - \Psi_{n+1}(0)(\overline{l_{n-1} z \Phi_{n-1}(z)} + (\nu_{n-1} z + 1)\Phi_{n-1}^*(z)) \\ &= z(z + \nu_{n-1})\Phi_{n-1}(z) + z l_{n-1} \Phi_{n-1}^*(z) = z\Psi_n(z), \end{aligned}$$

where the fourth equality follows from (2.30) and (6.17). That is,  $\{\Psi_n\}_{n \geq 0}$  satisfies a recurrence relation like (2.25), and it is therefore an orthogonal sequence with respect

to some linear functional  $\widetilde{\mathcal{L}}$ . We will prove that  $\widetilde{\mathcal{L}} = \mathcal{L}_{R^{(-1)}}$ . For  $0 \leq k \leq n-1$ , consider

$$\begin{aligned} \langle \widetilde{\mathcal{L}}, (z+z^{-1} - (\alpha + \bar{\alpha}))\Phi_n(z)\bar{z}^k \rangle &= \langle \widetilde{\mathcal{L}}, (z^2 - (\alpha + \bar{\alpha})z + 1)\Phi_n(z)\bar{z}^{k+1} \rangle \\ &= \langle \widetilde{\mathcal{L}}, \Psi_{n+2}(z)\bar{z}^{k+1} \rangle + \nu_n^{-1} \langle \widetilde{\mathcal{L}}, (\Psi_{n+1}(z) - \Phi_{n+1}(0)\Psi_{n+1}^*(z))\bar{z}^{k+1} \rangle = 0. \end{aligned}$$

On the other hand, for  $k = n$  we get

$$\langle \widetilde{\mathcal{L}}, \Psi_{n+2}(z) + \nu_n^{-1} (\Psi_{n+1}(z) - \Phi_{n+1}(0)\Psi_{n+1}^*(z))\bar{z}^{k+1} \rangle = \nu_n^{-1} \widetilde{k}_{n+1} = k_n.$$

Thus,  $\{\Phi_n\}_{n \geq 0}$  is the sequence of monic polynomials orthogonal with respect to  $(z+z^{-1} - (\alpha + \bar{\alpha}))\widetilde{\mathcal{L}}$ . But then  $(z+z^{-1} - (\alpha + \bar{\alpha}))\widetilde{\mathcal{L}} = \mathcal{L}$  and, therefore,  $\widetilde{\mathcal{L}} = \mathcal{L}_{R^{(-1)}}$ .  $\square$

**Theorem 6.3.2.** *Let  $\mathcal{L}$  be a positive definite linear functional and  $\sigma$  its associated measure. If  $\mathcal{L}_{R^{(-1)}}$  is a quasi-definite linear functional, then*

- i)  $(\mathfrak{I}(\widetilde{c}_1))^2 \neq (1 - (\Re(\alpha))^2)\widetilde{c}_0^2 - \Re(\alpha)\widetilde{c}_0 - \frac{1}{4}$ .
- ii)  $(1 - |\Phi_n(0)|^2)\nu_n^2 + A_{n+1}\nu_n + 1 - |\Psi_{n+1}(0)|^2 = 0$ ,  $n \geq 1$ ,

where  $A_n = \overline{\Psi_n(0)}\Phi_{n-1}(0) + \Psi_n(0)\overline{\Phi_{n-1}(0)} + \alpha + \bar{\alpha}$ .

*Proof.* From (6.12), for  $k = 0$  and assuming that  $c_0 = 1$ , we have

$$\Re(\widetilde{c}_1) = \frac{1}{2} + \Re(\alpha)\widetilde{c}_0.$$

In addition, in order to be  $\mathcal{L}_{R^{(-1)}}$  a quasi-definite functional, we need

$$\det \widetilde{\mathbf{T}}_1 = \begin{vmatrix} \widetilde{c}_0 & \widetilde{c}_1 \\ \widetilde{c}_{-1} & \widetilde{c}_0 \end{vmatrix} = \widetilde{c}_0^2 - (\Re(\widetilde{c}_1))^2 - (\mathfrak{I}(\widetilde{c}_1))^2 \neq 0,$$

where  $\widetilde{\mathbf{T}}$  is the Toeplitz matrix associated with  $\mathcal{L}_{R^{(-1)}}$ , and  $\widetilde{\mathbf{T}}_n$  is its corresponding  $(n+1) \times (n+1)$  leading principal submatrix. Therefore, for the choice of  $\alpha$ , we get

$$(\mathfrak{I}(\widetilde{c}_1))^2 \neq \widetilde{c}_0^2 - \left( \frac{1}{2} + \Re(\alpha)\widetilde{c}_0 \right)^2,$$

which is i). Thus,  $\widetilde{c}_0$  and  $\mathfrak{I}(\widetilde{c}_1)$  are free parameters, while  $\Re(\widetilde{c}_1)$  is determined by  $\widetilde{c}_0$  and the choice of  $\alpha$ .

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

Furthermore, we have

$$\begin{aligned} k_n &= \langle \Phi_n, \Phi_n \rangle_{\mathcal{L}} = \langle \Psi_n, \Phi_n \rangle_{\mathcal{L}} = \left\langle \left( z + z^{-1} - (\alpha + \bar{\alpha}) \right) \Psi_n(z), \Phi_n(z) \right\rangle_{\mathcal{L}_{R^{(-1)}}} \\ &= - \left( \Psi_{n+1}(0) \overline{\Phi_n(0)} + \alpha + \bar{\alpha} \right) \tilde{k}_n + \langle \Psi_n(z), z \Phi_n(z) \rangle_{\mathcal{L}_{R^{(-1)}}}. \end{aligned}$$

On the other hand, from (6.15),

$$\langle \Psi_n(z), z \Phi_n(z) \rangle_{\mathcal{L}_{R^{(-1)}}} = -\overline{\Psi_{n+1}(0)} \Psi_n(0) \tilde{k}_n - \frac{\tilde{k}_n}{k_{n-1}} \tilde{k}_n + \frac{\tilde{k}_n}{k_{n-1}} \overline{\Psi_{n+1}(0)} \Phi_{n-1}(0) \tilde{k}_n,$$

and from (6.17),

$$\begin{aligned} \langle \Psi_n(z), z \Phi_n(z) \rangle_{\mathcal{L}_{R^{(-1)}}} &= -\overline{\Psi_{n+1}(0)} \Psi_n(0) \tilde{k}_n - \frac{\tilde{k}_n}{k_{n-1}} \tilde{k}_n \\ &\quad + \left( \Psi_n(0) - \Phi_n(0) + \frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0) \right) \overline{\Psi_{n+1}(0)} \tilde{k}_n \\ &= -\overline{\Psi_{n+1}(0)} \Phi_n(0) \tilde{k}_n - \frac{\tilde{k}_n}{k_{n-1}} \tilde{k}_n + \frac{\tilde{k}_n}{k_{n-1}} |\Psi_{n+1}(0)|^2 \tilde{k}_n. \end{aligned}$$

Thus, if  $A_{n+1} = \overline{\Psi_{n+1}(0)} \Phi_n(0) + \Psi_{n+1}(0) \overline{\Phi_n(0)} + \alpha + \bar{\alpha}$ , then

$$\begin{aligned} k_n &= -A_{n+1} \tilde{k}_n + \left( |\Psi_{n+1}(0)|^2 - 1 \right) \frac{\tilde{k}_n}{k_{n-1}} \tilde{k}_n, \\ 1 - |\Phi_n(0)|^2 &= -A_{n+1} \frac{\tilde{k}_n}{k_{n-1}} + \left| \frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0) \right|^2 - \left( \frac{\tilde{k}_n}{k_{n-1}} \right)^2. \end{aligned}$$

Since, from (6.17),  $\frac{\tilde{k}_n}{k_{n-1}} \Psi_{n+1}(0) = \frac{\tilde{k}_n}{k_{n-1}} \Phi_{n-1}(0) + \Phi_n(0) - \Psi_n(0)$ , we obtain

$$1 - |\Psi_n(0)|^2 = -A_n \frac{\tilde{k}_n}{k_{n-1}} - \left( 1 - |\Phi_{n-1}(0)|^2 \right) \left( \frac{\tilde{k}_n}{k_{n-1}} \right)^2.$$

Therefore,

$$\left( 1 - |\Phi_{n-1}(0)|^2 \right) \left( \frac{\tilde{k}_n}{k_{n-1}} \right)^2 + A_n \frac{\tilde{k}_n}{k_{n-1}} + 1 - |\Psi_n(0)|^2 = 0, \quad (6.18)$$

which is *ii*). □

---

Now, from (6.17),

$$\begin{aligned}\Psi_{n+1}(0) &= \Phi_{n-1}(0) + [\Phi_n(0) - \Psi_n(0)] \frac{k_{n-1}}{\tilde{k}_n} \\ &= \frac{(\Phi_n(0) - \Psi_n(0)) \prod_{k=1}^{n-1} (1 - |\Phi_k(0)|^2)}{\prod_{k=1}^n (1 - |\Psi_k(0)|^2) \tilde{c}_0} + \Phi_{n-1}(0), \quad n \geq 1.\end{aligned}$$

We can thus build an algorithm to compute recursively the sequence  $\{\Psi_{n+1}(0)\}_{n \geq 1}$ , starting from  $\Psi_1(0)$ ; see Algorithm 6.1.

---

**Algorithm 6.1** Check Regularity

---

**input**  $\alpha, \tilde{c}_0, \{\Phi_n(0)\}_{n \geq 1}$

1:  $\Re(\tilde{c}_1) = \frac{1}{2} + \Re(\alpha) \tilde{c}_0$

2: **if** Theorem 6.3.2 *i* **then**

3:  $\Psi_1(0) = -\frac{\tilde{c}_1}{\tilde{c}_0}$

4: **end if**

5: **for**  $n = 1, 2, \dots$  **do**

6:  $\Psi_{n+1}(0) = \frac{(\Phi_n(0) - \Psi_n(0)) \prod_{k=1}^{n-1} (1 - |\Phi_k(0)|^2)}{\prod_{k=1}^n (1 - |\Psi_k(0)|^2) \tilde{c}_0} + \Phi_{n-1}(0)$

7: **if**  $|\Psi_{n+1}(0)| = 1$  **then**

8: **break**

9: **end if**

10: **end for**

---

**Example: Chebyshev polynomials**

Let  $d\sigma(\theta)$  the measure associated with the Chebyshev polynomials defined in (4.27). It is well known [178] that the family of Verblunsky coefficients associated with  $d\sigma$  is

$$\Phi_n(0) = \frac{1}{n+1}, \quad n \geq 1.$$

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS




Figure 6.1: Verblunsky coefficients for Chebyshev polynomials with  $\Re\alpha = 0.8$  and  $a = 0.5i$

Now, let us consider the perturbation

$$d\tilde{\sigma}(\theta) = \frac{|z-1|^2}{z+z^{-1}-(\alpha+\bar{\alpha})} \frac{d\theta}{2\pi}, \quad |z|=1,$$

where  $\Re(\alpha) = 0.6$ . Notice that  $b = 0.6 + 0.8i$ . Then, according to (6.11),

$$\tilde{c}_0 = \frac{1}{1.6i} \left( \int_0^{2\pi} \frac{|e^{i\theta}-1|^2-0.8}{1-(0.6+0.8i)e^{-i\theta}} \frac{d\theta}{2\pi} - \int_0^{2\pi} \frac{|e^{i\theta}-1|^2-0.8}{1-(0.6-0.8i)e^{-i\theta}} \frac{d\theta}{2\pi} \right) = -1,$$

and

$$\begin{aligned} \tilde{c}_1 &= \frac{1}{1.6i} \left( \int_0^{2\pi} \frac{(|e^{i\theta}-1|^2-0.8)e^{i\theta}}{1-(0.6+0.8i)e^{-i\theta}} \frac{d\theta}{2\pi} - \int_0^{2\pi} \frac{(|e^{i\theta}-1|^2-0.8)e^{i\theta}}{1-(0.6-0.8i)e^{-i\theta}} \frac{d\theta}{2\pi} \right. \\ &\quad \left. + \frac{1}{2}(0.6+0.8i)(0.8) - \frac{1}{2}(0.6-0.8i)(0.8) \right) = 0.4. \end{aligned}$$

Observe that part *i*) of Theorem 6.3.2 holds. Applying the algorithm, the first 500 Verblunsky coefficients are shown in Figure 6.1(a). Notice that all of the new Verblunsky coefficients are real. They are distributed on both sides of the origin, in nearly symmetric intervals. If we repeat the computation for  $n = 1500$ , the values accumulate over such intervals. This is shown in Figure 6.1(b).

### Example: Geronimus polynomials

In this subsection we consider a linear functional such that the corresponding measure is supported on an arc of the unit circle. Such a situation appears [78; 178] when  $\Phi_n(0) = a$ ,  $n \geq 1$ , with  $0 < |a| < 1$ . Here the measure  $\sigma$  associated with  $\{\Phi_n(0)\}_{n \geq 1}$  is

supported on the arc

$$\Delta_\nu = \{e^{i\theta} : \nu \leq \theta \leq 2\pi - \nu\},$$

with  $\cos(\nu/2) = \sqrt{1 - |a|^2}$ , but it can have a mass point located on  $\mathbb{T}$ . The orthogonality measure  $\sigma$  is given by

$$d\sigma(\theta) = \frac{\sqrt{\sin\left(\frac{\theta+\nu}{2}\right)\sin\left(\frac{\theta-\nu}{2}\right)}}{2\pi\sin\left(\frac{\theta-\tau}{2}\right)} d\theta + \mathbf{m}_\tau \delta(z - e^{i\tau}), \quad (6.19)$$

where  $e^{i\tau} = \frac{1-a}{1-\bar{a}}$  and

$$\mathbf{m}_\tau = \begin{cases} \frac{2|a|^2 - a - \bar{a}}{|1-a|}, & \text{if } |1-2a| > 1, \\ 0, & \text{if } |1-2a| \leq 1. \end{cases}$$

Moreover, the orthonormal polynomials associated with  $\sigma$  are given by

$$\varphi_n(z) = \frac{1}{(1-|a|^2)^{n/2}} \left( (z+a) \frac{z_1^n - z_2^n}{z_1 - z_2} - z(1-|a|^2) \frac{z_1^{n-1} - z_2^{n-1}}{z_1 - z_2} \right), \quad n \in \mathbb{N},$$

with

$$z_1 = \frac{z+1 + \sqrt{(z-e^{i\nu})(z-e^{-i\nu})}}{2}, \quad z_2 = \frac{z+1 - \sqrt{(z-e^{i\nu})(z-e^{-i\nu})}}{2}.$$

Consider a perturbation of (6.19) given by

$$d\tilde{\sigma}(z) = \frac{d\sigma(z)}{z + z^{-1} - (\alpha + \bar{\alpha})},$$

with  $\Re\alpha = 0.8$  and  $a = 0.5i$ . Notice that in this case  $b = 0.8 + 0.6i$  and thus  $b \notin \Delta_\nu$ . Then,

$$\begin{aligned} \tilde{c}_0 &= \int_{\frac{\pi}{3}}^{\frac{5\pi}{3}} \frac{\sqrt{\sin\left(\frac{1}{2}\theta + \frac{1}{6}\pi\right)\sin\left(\frac{1}{2}\theta - \frac{1}{6}\pi\right)}}{2(\cos\theta - 0.8)\pi\sin(\theta/2)} d\theta = -0.45876, \\ \tilde{c}_1 &= \int_{\frac{\pi}{3}}^{\frac{5\pi}{3}} \frac{(\cos\theta + i\sin\theta) \sqrt{\sin\left(\frac{1}{2}\theta + \frac{1}{6}\pi\right)\sin\left(\frac{1}{2}\theta - \frac{1}{6}\pi\right)}}{2(\cos\theta - 0.8)\pi\sin(\theta/2)} d\theta = 0.13299, \end{aligned}$$

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

and  $i$ ) holds. In such a situation, the algorithm becomes

$$\Psi_{n+1}(0) = \frac{[a - \Psi_n(0)](1 - |a|^2)^{n-1}}{\prod_{k=1}^n (1 - |\Psi_k(0)|^2) \tilde{c}_0} + a, \quad n \geq 1,$$

and the first 500 Verblunsky coefficients are shown in Figure 6.2(a).




Figure 6.2: Verblunsky coefficients for Geronimus polynomials with  $\Re \alpha = 0.8$  and  $a = 0.5i$

As it is shown in this figure, the Verblunsky coefficients associated with the modified measure have the same argument with respect to a certain point (the value of  $a$ ). That is, they are located on a straight line, on both sides of  $a$ . When  $n$  increases, the density of the points on the line increases, as shown in Figure 6.2(b).

### 6.3.2 C-functions

Assuming that  $\mathcal{L}_{R^{(-1)}}$  is a quasi-definite linear functional, we denote its associated  $\mathcal{C}$ -function by  $F_{R^{(-1)}}$ . Multiplying (6.12) by  $z^k$ ,  $k \geq 1$ , and replacing in (2.39), we get

$$\sum_{k=1}^{\infty} c_{-k} z^k = \sum_{k=1}^{\infty} \tilde{c}_{-(k+1)} z^k + \sum_{k=1}^{\infty} \tilde{c}_{-(k-1)} z^k - (\alpha + \bar{\alpha}) \sum_{k=1}^{\infty} \tilde{c}_{-k} z^k,$$

$$F(z) - 1 = (z + z^{-1} - (\alpha + \bar{\alpha})) F_{R^{(-1)}}(z) + \tilde{c}_0 (z - z^{-1} + (\alpha + \bar{\alpha})) - 2\tilde{c}_{-1}.$$

Therefore

$$F_{R^{(-1)}}(z) = \frac{F(z) + (z^{-1} - z - (\alpha + \bar{\alpha})) \tilde{c}_0 + 2\tilde{c}_{-1} - 1}{z + z^{-1} - (\alpha + \bar{\alpha})}.$$



---

Notice that from (6.12),  $1 + (\alpha + \bar{\alpha})\tilde{c}_0 = \tilde{c}_1 + \tilde{c}_{-1}$ , and thus

$$F_{R^{(-1)}}(z) = \frac{zF(z) - \tilde{c}_0z^2 + (\tilde{c}_{-1} - \tilde{c}_1)z + \tilde{c}_0}{z^2 - (\alpha + \bar{\alpha})z + 1}, \quad (6.20)$$

which is equivalent to

$$F_{R^{(-1)}}(z) = \frac{zF(z)}{z^2 - (\alpha + \bar{\alpha})z + 1} + m_1 \frac{z+b}{z-b} + m_2 \frac{z+\bar{b}}{z-\bar{b}}, \quad (6.21)$$

where  $b, \bar{b}$  are the zeros of  $z^2 - (\alpha + \bar{\alpha})z + 1$ , with  $|b| = 1$ , and

$$m_1 = -\frac{1}{2} \left( \tilde{c}_0 + \frac{\Im(\tilde{c}_1)}{\Im(b)} \right), \quad m_2 = -\frac{1}{2} \left( \tilde{c}_0 - \frac{\Im(\tilde{c}_1)}{\Im(b)} \right).$$

On the other hand, from (6.21)

$$\begin{aligned} F_{R^{(-1)}}(z) &= \left( \frac{b}{(b-\bar{b})(z-b)} - \frac{\bar{b}}{(b-\bar{b})(z-\bar{b})} \right) F(z) \\ &\quad - m_1 (1 + \bar{b}z) \sum_{k=0}^{\infty} \frac{z^k}{b^k} - m_2 (1 + bz) \sum_{k=0}^{\infty} b^k z^k \\ &= \left( \sum_{k=1}^{\infty} \frac{b^k - \bar{b}^k}{b - \bar{b}} z^k \right) F(z) - m_1 \left( 1 + 2 \sum_{k=1}^{\infty} \bar{b}^k z^k \right) - m_2 \left( 1 + 2 \sum_{k=1}^{\infty} b^k z^k \right), \end{aligned}$$

and thus

$$\tilde{c}_0 + 2 \sum_{k=1}^{\infty} \tilde{c}_{-k} z^k = \left( \sum_{k=1}^{\infty} \frac{b^k - \bar{b}^k}{b - \bar{b}} z^k \right) \left( 1 + 2 \sum_{k=1}^{\infty} c_{-k} z^k \right) - m_1 \left( 1 + 2 \sum_{k=1}^{\infty} \bar{b}^k z^k \right) - m_2 \left( 1 + 2 \sum_{k=1}^{\infty} b^k z^k \right).$$

Therefore, comparing coefficients of  $z^n$  on both sides of the last expression, we have for  $n \geq 2$ ,

$$\tilde{c}_{-n} = \sum_{k=1}^{n-1} \frac{b^k - \bar{b}^k}{b - \bar{b}} c_{-(n-k)} + \frac{1}{2} \frac{b^n - \bar{b}^n}{b - \bar{b}} - m_1 \bar{b}^n - m_2 b^n.$$

Comparing the independent terms and the coefficients for  $z$  we can deduce (6.12) for

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

$n = 0$  and  $n = 1$ . Furthermore, denoting this transformation by  $\mathcal{F}_{R^{(-1)}}$ ,

$$\mathcal{F}_R(\alpha) \circ \mathcal{F}_{R^{(-1)}}(\alpha) = I, \quad \mathcal{F}_{R^{(-1)}}(\alpha) \circ \mathcal{F}_R(\alpha) = \mathcal{F}_U(b, \widehat{\mathbf{m}}_1) \circ \mathcal{F}_U(\bar{b}, \widehat{\mathbf{m}}_2),$$

that follows in a straightforward way from the definition of  $\overline{\mathcal{F}}_R$  and  $\mathcal{F}_{R^{(-1)}}$ .

Denoting  $H(z) = (\mathcal{F}_{R^{(-1)}}(\alpha) \circ \mathcal{F}_R(\alpha))[F(z)]$ ,

$$H(z) = \frac{zF_R(z) - \widetilde{c}_0 z^2 + (\widetilde{c}_{-1} - \widetilde{c}_1)z + \widetilde{c}_0}{z^2 - (\alpha + \bar{\alpha})z + 1} = F(z) + \widehat{\mathbf{m}}_1 \frac{z+b}{z-b} + \widehat{\mathbf{m}}_2 \frac{z+\bar{b}}{z-\bar{b}},$$

with  $\widehat{\mathbf{m}}_1 = \widetilde{\mathbf{m}}_1 + \mathbf{m}_1$ ,  $\widehat{\mathbf{m}}_2 = \widetilde{\mathbf{m}}_2 + \mathbf{m}_2$ , and

$$\widetilde{\mathbf{m}}_1 = \frac{1}{2} \left( 1 + \frac{\Im(c_1)}{\Im(b)} \right), \quad \widetilde{\mathbf{m}}_2 = \frac{1}{2} \left( 1 - \frac{\Im(c_1)}{\Im(b)} \right).$$

As in (6.8), given a finite composition of order  $k \in \mathbb{N}$  of  $\mathcal{L}_{R^{(-1)}}$  defined by

$$\mathcal{L} = \mathfrak{R} \left( \prod_{i=1}^k (z - \alpha_i) \right) \mathcal{L}_{R^{(-k)}}, \quad \alpha_i \in \mathbb{C}, \quad (6.22)$$

we can deduce

**Lemma 6.3.1.** *A spectral transformation given by  $\widetilde{F}(z) = \frac{F(z) + B(z)}{D(z)}$  such that  $B$  and  $D$  provide a true asymptotic behavior to (2.43) is equivalent to (6.22). Furthermore,  $B$  and  $D$  are given by*

$$B(z) = \widetilde{P}(z) - \widetilde{P}_*(z), \quad D(z) = \mathfrak{R}p(z), \quad (6.23)$$

where  $\widetilde{P}$  is the polynomial of second kind of

$$p(z) = \prod_{i=1}^k (z - \alpha_i)$$

with respect to the linear functional (6.22).

*Proof.* From Lemma 6.2.1 it is straightforward to show that the  $G$ -function  $F_{\mathcal{L}_{R^{(-1)}}}$  is the reciprocal of  $F_{\mathcal{L}_*}$ . However, in contrast to  $F_R$ ,  $F_{R^{(-1)}}$  contains three free parameters  $\alpha$ ,  $c_0$  and  $\Im c_1$ , while  $\Re c_1$  is determined by  $c_0$  and the choice of  $\alpha$ .

Conversely, if we start with a spectral transformation with  $A = 1$ ,  $B$  is a hermitian Laurent polynomial of degree one and  $C = 0$ , then  $D$  is a hermitian Laurent polynomial

---

of degree one with three restrictions for their coefficients. We hence get just (6.20). Using an analog of Lemma 6.2.1 we can complete the proof.  $\square$

Notice that for different values  $\alpha_1, \alpha_2, \dots, \alpha_r$  and  $\beta_1, \beta_2, \dots, \beta_r$ ,  $\mathcal{F}_{R^{(k)}}[F] = F_{R^{(k)}}$  and  $\mathcal{F}_{R^{(-k)}}[F] = F_{R^{(-k)}}$  we get

$$\mathcal{F}_{R^{(k)}}(\alpha_1, \alpha_2, \dots, \alpha_r) \circ \mathcal{F}_{R^{(-k)}}(\beta_1, \beta_2, \dots, \beta_r) = \mathcal{F}_{R^{(-k)}}(\beta_1, \beta_2, \dots, \beta_r) \circ \mathcal{F}_{R^{(k)}}(\alpha_1, \alpha_2, \dots, \alpha_r).$$

Moreover, for the same parameters, we have the following relations

$$\begin{aligned} (\mathcal{F}_{R^{(-k)}} \circ \mathcal{F}_{R^{(k)}})(\alpha_1, \alpha_2, \dots, \alpha_r) &= \mathcal{I} \\ (\mathcal{F}_{R^{(k)}} \circ \mathcal{F}_{R^{(-k)}})(\alpha_1, \alpha_2, \dots, \alpha_r) &= \mathcal{F}_U(\alpha_1, \dots, \alpha_r) \circ \mathcal{F}_U(\bar{\alpha}_1, \dots, \bar{\alpha}_r), \end{aligned}$$

where  $\mathcal{F}_U(\alpha_1, \dots, \alpha_k)$  is the so-called general Uvarov spectral transformation as the result of the addition of masses at the points  $z = \alpha_1, z = \alpha_2, \dots, z = \alpha_r$ ; see Appendix B.

## 6.4 Rational spectral transformations

Spectral transformations under the modification of a finite number of moments are given by

$$\widetilde{F}_j(z) = F(z) + E(z),$$

where  $E(z) = \sum_{j \in G} m_j z^j$  for  $m_j \in \mathbb{C}$  and  $G$  a finite subset of non-negative integer numbers.

Hence, a generator system of local spectral transformations follows immediately from Chapter 4.

**Theorem 6.4.1.** *A local spectral transformation can be obtained as a finite composition of spectral transformations associated with the linear functionals  $\mathcal{L}_j$  defined in (4.33).*

In order to prove our main results in the following lemma we characterize the polynomial coefficients of (2.43).

**Lemma 6.4.1.** *Only one of the following two statements holds:*

- i) *The polynomial coefficients in (2.43) are hermitian Laurent polynomials of the same degree such that*

$$A_*(z) = A(z), \quad B_*(z) = -B(z), \quad C_*(z) = -C(z), \quad D_*(z) = D(z).$$

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

ii) *The polynomial coefficients in (2.43) are self-reciprocal polynomials of the same degree such that*

$$A^*(z) = A, \quad B^*(z) = -B(z), \quad C^*(z) = -C(z), \quad D^*(z) = D(z).$$

*Proof.* Since  $F$  is the rational spectral transformation associated with (2.43) given by

$$F(z) = \frac{-D(z)\widetilde{F}(z) + B(z)}{C(z)\widetilde{F}(z) - A(z)}, \quad (6.24)$$

multiplying and dividing (6.24) for  $z^l$ , where  $l$  is the minimum non-negative integer number such that  $z^l A(z)$ ,  $z^l B(z)$ ,  $z^l C(z)$ , and  $z^l D(z)$  are polynomials, and using the characterization of orthogonal polynomials with respect to the functional  $\mathcal{L}$  (2.41)-(2.42), we immediately get

$$\begin{aligned} z^l(-\Phi_n(z)D(z) - \Omega_n(z)C(z))\widetilde{F}(z) - z^l(-\Phi_n(z)B(z) - \Omega_n(z)A(z)) &= O(z^{n+\nu}), \\ z^l(-\Phi_n^*(z)D(z) + \Omega_n^*(z)C(z))\widetilde{F}(z) + z^l(\Phi_n^*(z)B(z) - \Omega_n^*(z)A(z)) &= O(z^{n+1+\nu}), \end{aligned}$$

where  $\nu$  is a positive integer such that

$$z^l(A(z) - C(z)\widetilde{F}(z)) = O(z^\nu).$$

Therefore, we can deduce that the new polynomial coefficients  $z^l A(z)$ ,  $z^l B(z)$ ,  $z^l C(z)$ , and  $z^l D(z)$  satisfy the following relations

$$(z^l A(z))^* = z^l A(z), \quad (z^l B(z))^* = -z^l B(z), \quad (z^l C(z))^* = -z^l C(z), \quad \text{and} \quad (z^l D(z))^* = z^l D(z).$$

Suppose that  $l \neq 0$ . Then,  $A$ ,  $B$ ,  $C$ , and  $D$  are Laurent hermitian polynomials of the same degree  $l$ , which prove *i*). On the other hand, if we suppose that  $l = 0$ , then  $A$ ,  $B$ ,  $C$ , and  $D$  are self-reciprocal polynomials of the same degree. Thus, *ii*) follows.  $\square$

If we have a generic rational spectral transformation with self-reciprocal polynomial coefficients of odd degree, then it can not be transformed into a equivalent rational spectral transformation with hermitian Laurent polynomial coefficients. We use the following result concerning the symmetrization of sequence of orthogonal polynomials to study this problem.

**Theorem 6.4.2.** [133] *There exists one and only one sequence of monic polynomials*

---

$\{\Psi_n\}_{n \geq 0}$  orthogonal with respect to a hermitian linear functional such that

$$\Psi_{2n}(z) = \Phi_n(z^2), \quad n \geq 0.$$

Furthermore,

$$\Psi_{2n+1}(z) = z\Phi_n(z^2), \quad n \geq 0.$$

Notice that the odd Verblunsky coefficients for the new sequence,  $\{\Psi(0)_n\}_{n \geq 0}$ , are zero. A linear functional  $\mathcal{D}$  is said to be symmetric if all its moments of odd order are 0, i.e.,

$$\langle \mathcal{D}, z^{2n+1} \rangle = 0, \quad n \geq 0.$$

It is interesting to recall that in the real line case if we look for  $\{Q_n\}_{n \geq 0}$ , which is a sequence of monic orthogonal polynomials and such that  $Q_{2n+1}(x) = xP_n(x^2)$ , there is not a unique solution, i.e., we can find an infinity number of sequences of polynomials  $\{R_n\}_{n \geq 0}$  such that  $Q_{2n}(x) = R_n(x^2)$ .

**Theorem 6.4.3.** *A generic rational spectral transformation with self-reciprocal polynomial coefficients of odd degree has symmetric generators to the corresponding symmetric rational spectral transformation which has hermitian Laurent polynomial coefficients.*

*Proof.* Let  $\{\tilde{\Phi}_n\}_{n \geq 0}$  be the sequence of monic orthogonal polynomials associated with the  $C$ -functions,  $F$ , a rational spectral transformation of  $F$  given by (2.43). We assume that  $A, B, C$ , and  $D$  are self-reciprocal polynomials of odd degree. From Theorem 6.4.2 there is an unique sequence of monic orthogonal polynomials  $\{\psi_n\}_{n \geq 0}$  (respectively  $\{\tilde{\psi}_n\}_{n \geq 0}$ ) such that

$$\tilde{\Psi}_{2n}(z) = \tilde{\Phi}_n(z^2), \quad \Psi_{2n}(z) = \Phi_n(z^2), \quad n \geq 0,$$

with respect to the symmetric linear functional  $\mathcal{D}$  (respectively  $\tilde{\mathcal{D}}$ ).

On the other hand, the corresponding symmetric  $C$ -functions associated with the symmetric functionals  $\mathcal{D}$  and  $\tilde{\mathcal{D}}$  are  $F_{\mathcal{D}}(z) = F(z^2)$  and  $F_{\tilde{\mathcal{D}}}(z) = \tilde{F}(z^2)$ , respectively. Therefore,

$$F_{\tilde{\mathcal{D}}}(z) = \frac{A(z^2)F_{\mathcal{D}}(z) + B(z^2)}{C(z^2)F_{\mathcal{D}}(z) + D(z^2)}$$

is a rational spectral transformation associated with  $F_{\mathcal{D}}$  with self-reciprocal polynomial coefficients of even degree. Thus,  $F_{\tilde{\mathcal{D}}}$  is equivalent to a rational spectral transformation

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

with hermitian Laurent polynomial coefficients.  $\square$

According to the previous theorem, in the next section we consider only hermitian Laurent polynomial coefficients.

### Example: Aleksandrov transformation

Notice that if the coefficients  $A$ ,  $B$ ,  $C$ , and  $D$  of (2.43) are constant, then  $A$  and  $D$  are real numbers and  $B$  and  $C$  are pure imaginary numbers. An example of this is the Aleksandrov transformation [178].

Let  $\{\Phi_n(0)\}_{n \geq 1}$  be the family of Verblunsky coefficients, and let  $\lambda$  be a complex number with  $|\lambda| = 1$ . When we consider a new family of Verblunsky coefficients defined by  $\{\Phi_n^\lambda(0)\}_{n \geq 1}$ , with  $\Phi_n^\lambda(0) = \lambda \Phi_n(0)$ ,  $n \geq 0$ , the resulting transformation is called Aleksandrov transformation. The  $C$ -functions are related by

$$F^\lambda(z) = \frac{(\lambda + 1)F(z) + \lambda - 1}{(\lambda - 1)F(z) + \lambda + 1}.$$

### 6.4.1 Generator system for rational spectral transformations

**Theorem 6.4.4.** *A generic linear spectral transformation can be obtained as a finite composition of spectral transformations associated with a modification of the functional by the real part of a polynomial and its inverse.*

*Proof.* Let  $H = \mathcal{F}[F]$  be the  $C$ -function obtained from (2.39) after a generic linear spectral transformation with hermitian Laurent polynomial coefficients. Hence, we can apply to  $F$  the finite composition of  $\mathcal{L}_R$  given in Lemma 6.2.1. Thus, we get a new  $C$ -function as a result of the composition

$$\mathcal{F}_{R^{(k)}}[H(z)] = \frac{1/2(\Re p(z))A(z)F(z) + 1/2(\Re p(z))B(z) + (P(z) - P_*(z))D(z)}{D(z)}$$

with some polynomials such that the zeros of the polynomial  $p$  can be chosen as the parameters of the spectral transformation. Hence, we can always choose  $\Re p = \tau D$ , where  $\tau$  is a constant. Thus, we get

$$\mathcal{F}_{R^{(k)}}[H(z)] = \tau A(z)F(z) + \tau B(z) + P(z) - P_*(z). \quad (6.25)$$

This transformation is reduced to the case considered in (6.10). Therefore, from Lemma 6.2.1 the  $C$ -function  $\mathcal{F}_{R^{(k)}}[H]$  is obtained from (2.39) by means of a finite composition

of (2.44).

On the other hand, from (6.25) and using a finite composition of (2.45), we get the  $C$ -function

$$H(z) = \mathcal{F}_{R^{(-k)}} [\mathcal{F}_{R^{(k)}} [\mathcal{F}[F(z)]]],$$

and so our statement holds.  $\square$

**Theorem 6.4.5.** *A generic rational spectral transformation can be obtained as a finite composition of linear and associated canonical spectral transformations.*

*Proof.* Assuming hermitian Laurent polynomial as coefficients of the perturbed linear functional, the application of backward transformations with even degree  $2k$  to the real spectral transformation (2.43) of degree  $k$  yields a new rational spectral transformation where the transformed Laurent polynomial  $\widetilde{C}$  is hermitian and is given by

$$\widetilde{C}(z) = z^k (A(z) + C(z)) (\widetilde{\Phi}_{2k})_*(z) + z^{-k} (C(z) - A(z)) \widetilde{\Phi}_{2k}(z).$$

Notice that the polynomial  $\widetilde{\Phi}_{2k}$  can be chosen in an arbitrary way. Indeed, instead of choosing arbitrary  $2k$  Verblunsky parameters we can choose the polynomial  $\widetilde{\Phi}_{2k}$  satisfying  $|\widetilde{\Phi}_{2k}(0)| \neq 1$ , and from the Schur-Cohn-Jury criterion we obtain a sequence of complex numbers  $\widetilde{\Phi}_{2k}(0), \dots, \widetilde{\Phi}_1(0)$  with modulus different of 1. Let  $(\Phi_n(0))_{n \geq 1}$  be the Verblunsky coefficients of the hermitian linear functional associated with (2.43). Then, by Favard's theorem  $(\widetilde{\Phi}_i(0))_{i=1}^{2k} \cup (\Phi_n(0))_{n \geq 1}$  arises as a new sequence of Verblunsky coefficients of a hermitian linear functional. Notice that it is unique.

On the other hand, in order to preserve the hermitian character of  $\widetilde{C}$ , the principal leading coefficients of  $A$  and  $C$  are different and not symmetric with respect to the origin. Thus,

$$\deg(A(z) + C(z)) = \deg(C(z) - A(z)) = k.$$

Moreover, without loss of generality, the polynomial  $z^k(A(z) + C(z))$  evaluated at  $z = 0$  can be chosen in such a way that its modulus is different from one. Therefore, we can choose the polynomial  $\widetilde{\Phi}_{2k}$  such that

$$\widetilde{\Phi}_{2k}(z) = -z^k(C(z) + A(z)).$$

Using Lemma 6.4.1 we obtain the reciprocal polynomial

$$(\widetilde{\Phi}_{2k})_*(z) = z^{-k}(C(z) - A(z)),$$

## 6. GENERATORS OF RATIONAL SPECTRAL TRANSFORMATIONS

---

leading to  $\tilde{C} = 0$ . But this means that we can reduce our rational spectral transformation to a linear spectral transformation and the result follows.  $\square$



# Chapter 7

## Conclusions and open problems

'Would you tell me, please, which way I ought to go from here?'

'That depends a good deal on where you want to get to,' said the Cat.

'I don't much care where —' said Alice.

'Then it doesn't matter which way you go,' said the Cat.

'— so long as I get SOMEWHERE,' Alice added as an explanation.

'Oh, you're sure to do that,' said the Cat, 'if you only walk long enough.'

— L. Carroll. *Alice's Adventures in the Wonderland*. Random House, New York, 1865

### 7.1 Conclusions

The results showed in this work have been focussed on the study of spectral transformations of linear functionals.

We have presented an alternative approach to the theory of orthogonal polynomials, giving a central role to continued fractions. The results have been proved under the assumption that the Verblunsky coefficients  $\{\Phi_n(0)\}_{n \geq 0}$  satisfy  $|\Phi_n(0)| > 1$ . As a consequence, we have obtained specific properties of the corresponding Hankel and Toeplitz determinants which we have used to deduce necessary and sufficient conditions for the existence of the moment problem associated with Chebyshev polynomials of the first kind.

We also proved that the zeros of orthogonal polynomial  $\Phi_n(\omega; \cdot)$  satisfy, for  $\Phi_n(0) > 1$ ,  $n \geq 1$ ,

$$0 < z_{n,1}(\omega) < z_{n,2}(\omega) < \dots < z_{n,n-1}(\omega) < z_{n,n}(\omega), \quad \omega < 1,$$

## 7. CONCLUSIONS AND OPEN PROBLEMS

---

$$z_{n,1}(\omega) < 0 < z_{n,2}(\omega) < \dots < z_{n,n-1}(\omega) < z_{n,n}(\omega), \quad \omega > 1.$$

and, for  $\Phi_n(0) < 1$ ,  $n \geq 1$ ,

$$z_{n,1}(\omega) < z_{n,2}(\omega) < \dots < z_{n,n-1}(\omega) < z_{n,n}(\omega) < 0, \quad \omega < 1,$$

$$z_{n,1}(\omega) < z_{n,2}(\omega) < \dots < z_{n,n-1}(\omega) < 0 < z_{n,n}(\omega), \quad \omega > 1.$$

The properties of the zeros allowed us to define a step function, and by using Helly's selection principle we have showed that an rsq-definite linear functional has an integral representation in terms of a measure that belongs to the class  $\mathcal{S}^3(0, 1, b)$ . This result played a key role to find conditions for the determinacy of our moment problem.

Motivated by applications in integrable systems, we have obtained explicit expressions of the orthogonal polynomials associated with the perturbations of a hermitian linear functional in terms of the first one. In the same direction, we defined a perturbation of a quasi-definite linear functional by the addition of the first derivative of the Dirac linear functional when its support is a point on the unit circle or two points symmetric with respect to the unit circle. In both cases we get

$$\lim_{n \rightarrow \infty} \frac{\Psi_n(z)}{\Phi_n(z)} = 1, \quad z \in \mathbb{C} \setminus \overline{\mathbb{D}},$$

where  $\{\Psi_n\}_{n \geq 0}$  is the sequence of orthogonal polynomials associated with the corresponding perturbed functional. Nevertheless, we are far from achieving our initial goals.

We have analyzed the previous results for asymptotics of the discrete Sobolev polynomials and their zeros when the mass tends to infinity. The study of these systems was motivated by the search of efficient algorithms for computing Fourier expansions of a function in terms of discrete Sobolev orthogonal polynomials. In this direction very few results are known in the literature. We have stated that the resulting polynomials tend to a linear combination of polynomial perturbations of several orders, i.e.,

$$\phi_n(z), (z - \alpha)\phi_{n-1}(z, d\sigma_1), \dots, (z - \alpha)^{j+1}\phi_{n-j-1}(z, d\sigma_{j+1}).$$

We have characterized the eigenvalues of the Hessenberg matrix  $(\mathbf{H}_\Psi)_n$  associated with the discrete Sobolev polynomials, i.e., the zeros of the discrete Sobolev polynomials, as the eigenvalues of a rank one perturbation of the Hessenberg matrix associated with the measure  $\sigma$ , i.e.,

$$(\mathbf{H}_\Psi)_n = \mathbf{L}_n(\mathbf{H}_n - \mathbf{A}_n)\mathbf{L}_n^{-1},$$

---

where  $\mathbf{A}_n$  and  $\mathbf{L}_n$  are known. As a consequence, the zeros of  $\Psi_n$  are the eigenvalues of the matrix  $\mathbf{H}_n - \mathbf{A}_n$ , a rank one perturbation of the matrix  $\mathbf{H}_n$ . We also proved a characterization for the limit of these discrete Sobolev polynomials, when the mass tends to infinity as extremal polynomials. The results have been deduced for measures on the Nevai class, so in this sense they are sharp. We also showed that, as an analog of the real line case, a generic rational spectral transformation of the  $C$ -function is a finite composition of four canonical spectral transformations which we have studied in detail. To obtain this result it was essential our classification and characterization of the polynomial coefficients of rational spectral transformations obtained from the characterization of orthogonal polynomials with respect to a functional given by Peherstorfer and Steinbauer. All our results are based on the strong relationship between the theory of orthogonal polynomials on the real line and on the unit circle.

## 7.2 Open problems

In this section we formulate some open questions which have arisen during our research.

### Zeros of Szegő-type polynomials

The Szegő-type polynomials [112; 113] are orthogonal with respect to a moment linear functional  $\mathcal{T}$ , such that their moments

$$c_n = \langle \mathcal{T}, z^n \rangle = c_{-n}, \quad n \geq 0,$$

are all complex. If the linear functional  $\mathcal{T}$  is such that  $\{c_n\}_{n \in \mathbb{Z}}$  is real and  $(-1)^{n(n+1)/2} \det \mathbf{T}_n > 0$ ,  $n \geq 0$ , then the zeros of associated polynomials have been studied in Chapter 3. We were unable to obtain the localization and asymptotic behavior of the zeros of Szegő-type orthogonal polynomials. In [112; 113] the authors establish that the hypergeometric functions

$$\Phi_n(b; z) = \frac{(2b+1)_n^+}{(b+1)_n^+} {}_2F_1(-n, b+1; 2b+2; 1-z), \quad 2b \neq \mathbb{Z}_-, \quad n \geq 0,$$

are the Szegő-type polynomials with respect to a moment functional with moments  $\{c_n(b)\}_{n \geq 0}$ ,

$$c_0(b) = 2, \quad c_n(b) = 2 \frac{(-b)_n^+}{(b+1)_n^+} = c_{-n}(b), \quad n \geq 1.$$

## 7. CONCLUSIONS AND OPEN PROBLEMS

Figure 7.1 shows the behavior of the zeros of  $\Phi_n(b; \cdot)$  for fixed  $n = 100$  and several values of the parameter  $b$ , namely  $b = 1000$  (green triangles),  $b = 100i$  (blue discs),  $b = 5 + 6i$  (purple square), and  $b = 11$  (yellow diamonds).




Figure 7.1: Zeros of the Szegő-type polynomial  ${}_2F_1(-100, b+1; 2b+2; 1-z)$

### Perturbations on anti-diagonals of Hankel matrices

Given  $j \geq 0$  it is natural to ask if the corresponding perturbation  $\mathcal{M}_j$ , studied in Chapter 4, preserves the positive definiteness of  $\mathcal{M}$ . Of course, the necessary and sufficient conditions are given in Theorem 4.1.1. However, if one is interested in the existence of a neighborhood  $(\tau_1, \tau_2)$  such that the functional  $\mathcal{M}_j$  is positive definite for every  $m_j \in (\tau_1, \tau_2)$ , then to determine such an interval can be very complicated. An open problem is to analyze if there exists a different approach that allows one to determine the values of  $m_j$  such that  $\mathcal{M}_j$  is positive definite. Certainly, the interval  $(\tau_1, \tau_2)$  should depend essentially on the initial functional  $\mathcal{M}$  and the point  $a$ .

Another question that might be of interest is if given two positive definite moment functionals  $\mathcal{M}$  and  $\widetilde{\mathcal{M}}$ , then there exists a sequence of perturbations  $\mathcal{M}_{j_k}$  such that we can obtain  $\widetilde{\mathcal{M}}$  from the consecutive applications of those perturbations to  $\mathcal{M}$ , i.e.,

$$\mathcal{M} \xrightarrow{\mathcal{M}_{j_1}} \mathcal{M}^{(1)} \xrightarrow{\mathcal{M}_{j_2}} \mathcal{M}^{(2)} \xrightarrow{\mathcal{M}_{j_3}} \dots \xrightarrow{\mathcal{M}_{j_k}} \mathcal{M}^{(k)} \rightarrow \widetilde{\mathcal{M}},$$

---

with the condition that the positiveness must be preserved in each step. As an example, consider the linear functionals  $\mathcal{M}^1$  and  $\mathcal{M}^2$ , associated with the Chebyshev polynomials of first and second kind, respectively. When using the basis  $\{1, (x-1), (x-1)^2, \dots\}$ , it is easy to see that one of the sequences of moments can be obtained from the other one by means of a shift. Thus, as explained in Chapter 4, one can go from  $\mathcal{M}^1$  to  $\mathcal{M}^2$  applying a sequence of perturbations  $\mathcal{M}_{j_k} = \mathcal{M}_k$ ,  $k = 0, 1, \dots$ , with  $a = 1$ , and  $m_{j_k}$ . However, proceeding in such a way, the positive definiteness would be lost after second step. Nevertheless, another sequence  $\mathcal{M}_{j_k}$  which preserves the positive definiteness may still exist.

### Zeros of Sobolev orthogonal polynomials on the real line

Several people have studied the zeros of the polynomials  $S_n(\lambda, \mathbf{c}, r; \cdot)$ , with  $\lambda = (\lambda_0, \lambda_1, \dots, \lambda_r)$  and  $\mathbf{c} = (c_0, c_1, \dots, c_r)$ , orthogonal with respect to the discrete Sobolev inner product of the form

$$\langle p, q \rangle_{\mathcal{D}_r} = \int_I p(x)q(x)d\mu(x) + \sum_{i=0}^r \lambda_i p^{(i)}(c_i)q^{(i)}(c_i), \quad c_i \in \mathbb{R}, \quad \lambda_i \geq 0, \quad r \geq 0, \quad (7.1)$$

where  $\mu$  is a positive Borel measure supported on  $I \subseteq \mathbb{R}$ . In the Appendix A we consider a particular case of (7.1).

We now formulate some questions: fix  $\lambda_i$ ,  $i = 0, 1, \dots, k-1, k+1, \dots, r$ , in (7.1), and consider the zeros of  $S_n$  as functions of  $\lambda_k$ . Then, how many zeros of  $S_n$  lie in the interior of the convex hull of  $I$ ? Moreover, are these zeros monotonic functions with respect to the parameter  $\lambda_k$ ? Do they converge when  $\lambda_k$  goes to infinity? If so, what is the speed of convergence? For some partial results towards these problems, for specific measures and vectors  $\lambda$ , we refer to [5; 6; 140; 157], [159] and the references therein. The most general result in this direction was obtained in [4]. There it is stated that every  $S_n(\lambda, \mathbf{c}, r; \cdot)$  possesses at least  $n - r$  zeros in the interior of the convex hull of  $I$ , when  $c_0 = c_1 = \dots = c_r$ .

### Infinity discrete Sobolev inner products on a rectifiable Jordan curve or arc

The sequences of orthogonal polynomials associated with an orthogonality measure with finitely many point masses outside a curve or arc have also been studied by Kaliaguine and Kononova [103; 104; 105]. The case of an infinity number of mass point for measures of the Szegő supported on the real line is considered by Peherstorfer

## 7. CONCLUSIONS AND OPEN PROBLEMS

---

and Yuditskii in [156]. Let  $f(z)$  be a function of one variable and let

$$Z = (\underbrace{z_1, \dots, z_1}_{l_1}, \underbrace{z_2, \dots, z_2}_{l_2}, \dots, \underbrace{z_m, \dots, z_m}_{l_m}, \dots)$$

be an infinite vector. Denote

$$\mathbf{f}(Z) = (f(z_1), \dots, f^{(l_1)}(z_1), \dots, f(z_m), \dots, f^{(l_m)}(z_m), \dots).$$

The infinite discrete Sobolev inner product in the complex plane is defined by

$$\langle f, g \rangle_{S_I} = \int_C f(z) \overline{g(z)} \omega(z) |dz| + \mathbf{f}(Z) \mathbf{A} \mathbf{g}(Z)^H, \quad (7.2)$$

where  $C$  is a rectifiable Jordan curve or arc in the complex plane;  $\mathbf{A}_M$ , the principal  $M \times M$  hermitian submatrix of the infinity matrix  $\mathbf{A}$ , is quasi-definite;  $z_i \in \Omega$ ,  $i = 1, 2, \dots$ , and  $\Omega$  denotes the connect component of  $\mathbb{C} \setminus C$  such that  $\infty \in \Omega$ . It is thus fundamental to ask: What can be said about the asymptotic behavior of discrete Sobolev orthogonal polynomial with respect to the inner product (7.2)?

### Toda lattices

The study of integrable systems on the unit circle is not so performant as in the real line case, and its development started at the end of the previous century [9], [65], when the system of non-linear differential-difference equations (1.7) was studied. The main focus was based on the spectral theory of the GGT matrix  $\mathbf{H}$ . It was found that the resulting GGT matrices satisfy a Lax equation

$$\mathbf{H}' = [(\mathbf{H} + \mathbf{H}^{-1})_+, \mathbf{H}],$$

More recently, some developments in this direction have been done; see Chapter 1. In [84], the author considers the equations (1.7) with a similar approach, but using an alternative matrix representation of the multiplication operator for Laurent orthogonal polynomials, the CMV matrix. The corresponding CMV matrices satisfy a Lax equation

$$\mathbf{C}' = [\mathbf{B}, \mathbf{C}],$$

where  $\mathbf{B}$  is an upper triangular matrix with two non-zero diagonals above the main diagonal, whose entries are also expressed in terms of the Verblunsky coefficients. In this situation, the non-linear differential-difference equation of their associated sequence of

---

monic orthogonal polynomials satisfy

$$\Phi'_n(z, t) = \Phi_{n+1}(z, t) - (z + \Phi_{n+1}(0, t)\overline{\Phi_n(0, t)})\Phi_n(z, t) - (1 - |\Phi_n(0, t)|^2)\Phi_{n-1}(z, t),$$

and the orthogonality measure associated with the Verblunsky coefficients is given by (1.8).

An interesting open problem is the study of Schur flows associated with the canonical linear spectral transformation of  $C$ -functions obtained in Chapter 6. Another interesting question is to analyze the analog of the Lax equations for the corresponding GGT and CMV matrices associated with such perturbations.

## **7. CONCLUSIONS AND OPEN PROBLEMS**

---



## Appendix A

# Discrete Sobolev orthogonal polynomials on the real line

Let  $\{Q_n(\lambda, c, j; \cdot)\}_{n \geq 0}$  be the sequence of orthogonal polynomials with respect to the discrete Sobolev inner product

$$\langle p, q \rangle_{\mathcal{D}_1} = \int_a^b p(x)q(x)d\mu(x) + \lambda p^{(j)}(c)q^{(j)}(c), \quad c \notin (a, b), \quad \lambda \in \mathbb{R}_+, \quad j \geq 0, \quad (\text{A.1})$$

where  $\mu$  is a positive Borel measure supported in the interval  $(a, b)$  (either  $a$  or  $b$  can be infinity) and  $p, q$  are polynomials with real coefficients. In what follows we assume all orthogonal polynomials, both those with respect to  $\mu$  and the Sobolev ones, to be monic.

In this appendix we prove that the zeros of discrete Sobolev orthogonal polynomials are monotonic functions of the parameter  $\lambda$  and establish their asymptotics when either  $\lambda$  converges to zero or to infinity. The precise location of the extreme zeros is also analyzed.

### A.1 Monotonicity and asymptotics of zeros

Let  $x_{n,k}(\lambda, c, j)$ ,  $k = 1, \dots, n$ , be the zeros of  $Q_n(\lambda, c, j; \cdot)$ . For  $\lambda = 0$  or  $n \leq j$ , the polynomials  $P_n = Q_n(\lambda, c, j; \cdot)$  are orthogonal with respect to the inner product

$$\langle p, q \rangle_{\mu} = \int_a^b p(x)q(x)d\mu(x).$$

When  $\lambda > 0$  and  $n > j$  some natural questions arise. Are the zeros of  $Q_n(\lambda, c, j; \cdot)$

## A. DISCRETE SOBOLEV ORTHOGONAL POLYNOMIALS

---

all real and do they belong to  $(a, b)$ ? If so, do the zeros of  $Q_n(c, \lambda, j; \cdot)$  interlace with the zeros of  $P_n$ ? Moreover, are the zeros  $x_{n,k}(\lambda, c, j)$  monotonic functions with respect to the parameter  $\lambda$ ? Do the zeros  $x_{n,k}(\lambda, c, j)$  converge when  $\lambda$  goes to infinity? If so, what is the speed of convergence?

The answer of the first two questions was given in [140]. He proved that the polynomial  $Q_n(\lambda, c, j; \cdot)$  possesses  $n$  real simple zeros and at most one of them is located outside the interval  $(a, b)$ . In addition, he showed the following interlacing property. If  $c \leq a$ , then

$$x_{n,1}(\lambda, c, j) < x_{n,1} < x_{n,2}(\lambda, c, j) < x_{n,2} < \cdots < x_{n,n}(\lambda, c, j) < x_{n,n}. \quad (\text{A.2})$$

If  $c \geq b$ , then

$$x_{n,1} < x_{n,1}(\lambda, c, j) < x_{n,2} < x_{n,2}(\lambda, c, j) < \cdots < x_{n,n} < x_{n,n}(\lambda, c, j). \quad (\text{A.3})$$

In this appendix our main contribution deals with the remaining questions posed above. Our result is general in three aspects: the measure involved is any positive Borel measure, the point  $c$  is any value outside  $(a, b)$ , and  $j$  is any positive integer. We obtain a new interlacing property, the monotonicity of  $x_{n,k}(\lambda, c, j)$  with respect to  $\lambda$ , as well as their convergence – when  $\lambda$  tends to infinity – to the zeros of some polynomial with a speed of convergence of order  $1/\lambda$ . For this propose we define the polynomial  $G_n(c, j; \cdot)$  by

$$G_n(c, j; x) = P_n(x) - \frac{P_n^{(j)}(c)}{K_{n-1}^{(j,j)}(c, c)} K_{n-1}^{(j,0)}(c, x), \quad (\text{A.4})$$

where  $K_{n-1}^{(r,s)}$  denotes the generalized Kernel polynomial of degree  $n-1$ ,

$$K_{n-1}^{(r,s)}(x, y) = \sum_{k=0}^{n-1} \frac{P_k^{(r)}(x) P_k^{(s)}(y)}{\langle P_k, P_k \rangle_\mu}.$$

Notice that when  $r = s = 0$ ,  $K_{n-1} = K_{n-1}^{(0,0)}$  is the usual Kernel polynomial.

**Theorem A.1.1.** *Let  $\lambda > 0$  and  $y_{n,1}(c, j), \dots, y_{n,n}(c, j)$  be the zeros of the polynomial  $G_n(c, j; \cdot)$  defined by (A.4). For every  $n > j$  and  $c \leq a$ , then*

$$y_{n,1}(c, j) < x_{n,1}(\lambda, c, j) < x_{n,1} < \cdots < y_{n,n}(c, j) < x_{n,n}(\lambda, c, j) < x_{n,n}.$$

Moreover, each  $x_{n,k}(\lambda, c, j)$  is a decreasing function of  $\lambda$ , for each  $k = 1, \dots, n$ .

---

On the other hand, if  $c \geq b$ , then

$$x_{n,1} < x_{n,1}(\lambda, c, j) < y_{n,1}(c, j) < \cdots < x_{n,n} < x_{n,n}(\lambda, c, j) < y_{n,n}(c, j). \quad (\text{A.5})$$

In addition, each  $x_{n,k}(\lambda, c, j)$  is an increasing function of  $\lambda$ , for  $1 \leq k \leq n$ .

In both cases, when  $\lambda$  goes to infinity,

$$\lim_{\lambda \rightarrow \infty} x_{n,k}(\lambda, c, j) = y_{n,k}(c, j)$$

and

$$\lim_{\lambda \rightarrow \infty} \lambda(y_{n,k}(c, j) - x_{n,k}(\lambda, c, j)) = \frac{P_n(y_{n,k}(c, j))}{K_{n-1}^{(j,j)}(c, c)G_n'(c, j; y_{n,k}(c, j))}. \quad (\text{A.6})$$

*Proof.* Let us consider the following expression for the discrete Sobolev orthogonal polynomial [131],

$$Q_n(\lambda, c, j; x) = P_n(x) - \frac{\lambda P_n^{(j)}(c)}{1 + \lambda K_{n-1}^{(j,j)}(c, c)} K_{n-1}^{(j,0)}(c, x). \quad (\text{A.7})$$

Considering the normalization

$$\tilde{Q}_n(\lambda, c, j; x) = (1 + \lambda K_{n-1}^{(j,j)}(c, c)) Q_n(\lambda, c, j; x),$$

we derive a simple representation for the  $n$ -th discrete Sobolev orthogonal polynomial

$$\tilde{Q}_n(\lambda, c, j; x) = P_n(x) + \lambda K_{n-1}^{(j,j)}(c, c) G_n(c, j; x).$$

Observe that  $G_n(c, j; x) = \lim_{\lambda \rightarrow \infty} Q_n(\lambda, c, j; x)$  is independent of  $\lambda$ . Then, evaluating the above expression in the zeros of  $P_n$  and  $Q_n(\lambda, c, j; \cdot)$ , we conclude that

$$\begin{aligned} \operatorname{sgn}(\tilde{Q}_n(\lambda, c, j; x_{n,k})) &= \operatorname{sgn}(G_n(c, j; x_{n,k})), \\ \operatorname{sgn}(P_n(x_{n,k}(\lambda, c, j))) &= -\operatorname{sgn}(G_n(c, j; x_{n,k}(\lambda, c, j))), \end{aligned}$$

for every  $k = 1, \dots, n-1$ . Therefore, using (A.2) and (A.3),  $G_n(c, j; \cdot)$  changes sign  $n-1$  times at the zeros of  $Q_n(\lambda, c, j; \cdot)$  and  $P_n$ . In other words, each interval  $(x_{n,k}, x_{n,k+1})$  and  $(x_{n,k}(\lambda, c, j), x_{n,k+1}(\lambda, c, j))$ ,  $1 \leq k \leq n-1$ , contains one zero of  $G_n(c, j; \cdot)$ . It remains to find the location of one zero of  $G_n(c, j; \cdot)$ . Taking into account that the polynomial  $G_n(c, j; \cdot)$  has a positive leader coefficient and the location of the point  $c$  with respect to

## A. DISCRETE SOBOLEV ORTHOGONAL POLYNOMIALS

---

the interval  $(a, b)$ , we obtain the inequalities stated.

To prove the monotonicity of the zeros  $x_{n,k}(\lambda, c, j)$  with respect to  $\lambda$ , for every non-negative  $\varepsilon$ , we consider the polynomial

$$\widetilde{Q}_n(\lambda + \varepsilon, c, j; x) = P_n(x) + (\lambda + \varepsilon)K_{n-1}^{(j,j)}(c, c)G_n(c, j; x) = \widetilde{Q}_n(\lambda, c, j; x) + \varepsilon K_{n-1}^{(j,j)}(c, c)G_n(c, j; x).$$

Thus, for  $\varepsilon > 0$ ,

$$\widetilde{Q}_n(\lambda + \varepsilon, c, j; x_{n,k}(\lambda, c, j)) = \varepsilon K_{n-1}^{(j,j)}(c, c)G_n(c, j; x_{n,k}(\lambda, c, j)),$$

and

$$\widetilde{Q}_n(\lambda + \varepsilon, c, j; x_{n,k+1}(\lambda, c, j)) = \varepsilon K_{n-1}^{(j,j)}(c, c)G_n(c, j; x_{n,k+1}(\lambda, c, j))$$

have opposite sign. Therefore, each interval  $(x_{n,k}(\lambda, c, j), x_{n,k+1}(\lambda, c, j))$  contains at least one zero of  $\widetilde{Q}_n(\lambda + \varepsilon, c, j; x)$ . In addition, if  $c \leq a$ , then

$$\operatorname{sgn}\left(\widetilde{Q}_n(\lambda + \varepsilon, c, j; x_{n,1}(\lambda, c, j))\right) = (-1)^{n+1}, \quad \lim_{x \rightarrow -\infty} \widetilde{Q}_n(\lambda + \varepsilon, c, j; x) = \begin{cases} +\infty, & n \text{ even;} \\ -\infty, & n \text{ odd.} \end{cases}$$

Hence,

$$x_{n,1}(\lambda + \varepsilon, c, j) < x_{n,1}(\lambda, c, j) < \cdots < x_{n,n}(\lambda + \varepsilon, c, j) < x_{n,n}(\lambda, c, j).$$

In the other situation, if  $c \geq b$ , then

$$\operatorname{sgn}\left(\widetilde{Q}_n(\lambda + \varepsilon, c, j; x_{n,n}(\lambda, c, j))\right) = -1, \quad \lim_{x \rightarrow +\infty} \widetilde{Q}_n(\lambda + \varepsilon, c, j; x) = +\infty.$$

Hence,

$$x_{n,1}(\lambda, c, j) < x_{n,1}(\lambda + \varepsilon, c, j) < \cdots < x_{n,n}(\lambda, c, j) < x_{n,n}(\lambda + \varepsilon, c, j).$$

It remains to prove the limit relations stated in the theorem. For this purpose, we define the polynomial  $\widehat{Q}_n(\lambda, c, j; \cdot)$  by

$$\widehat{Q}_n(\lambda, c, j; x) = \frac{1}{\lambda} P_n(x) + K_{n-1}^{(j,j)}(c, c)G_n(c, j; x).$$

---

Notice that the zeros of  $Q_n(\lambda, c, j; \cdot)$  and  $\widehat{Q}_n(\lambda, c, j; \cdot)$  are the same for each  $\lambda > 0$ . Since

$$\lim_{\lambda \rightarrow \infty} \widehat{Q}_n(\lambda, c, j; x) = K_{n-1}^{(j,j)}(c, c)G_n(c, j; x),$$

by Hurwitz's Theorem [188], the zeros  $x_{n,k}(\lambda, c, j)$  of  $Q_n(\lambda, c, j; \cdot)$  converge to the zeros  $y_{n,k}(c, j)$  of  $G_n(c, j; \cdot)$  when  $\lambda$  tends to infinity, that is,

$$\lim_{\lambda \rightarrow \infty} x_{n,k}(\lambda, c, j) = y_{n,k}(c, j), \quad k = 1, \dots, n.$$

On the other hand, by the Mean Value Theorem, there exist real numbers  $\theta_{n,k}$  between  $y_{n,k}(c, j)$  and  $x_{n,k}(\lambda, c, j)$ ,  $k = 1, \dots, n$ , such that

$$\frac{G_n(c, j; y_{n,k}(c, j)) - G_n(c, j; x_{n,k}(\lambda, c, j))}{y_{n,k}(c, j) - x_{n,k}(\lambda, c, j)} = G'_n(c, j; \theta_{n,k}),$$

or, equivalently,

$$\lambda(y_{n,k}(c, j) - x_{n,k}(\lambda, c, j)) = \frac{P_n(x_{n,k}(\lambda, c, j))}{K_{n-1}^{(j,j)}(c, c)G'_n(c, j; \theta_{n,k})}.$$

Since  $\lim_{\lambda \rightarrow \infty} x_{n,k}(\lambda, c, j) = y_{n,k}(c, j)$  and  $\theta_{n,k}$  is located between  $y_{n,k}(c, j)$  and  $x_{n,k}(\lambda, c, j)$  we also have  $\lim_{\lambda \rightarrow \infty} \theta_{n,k} = y_{n,k}(c, j)$ . Thus

$$\lim_{\lambda \rightarrow \infty} \lambda(y_{n,k}(c, j) - x_{n,k}(\lambda, c, j)) = \lim_{\lambda \rightarrow \infty} \frac{P_n(x_{n,k}(\lambda, c, j))}{K_{n-1}^{(j,j)}(c, c)G'_n(c, j; \theta_{n,k})} = \frac{P_n(y_{n,k}(c, j))}{K_{n-1}^{(j,j)}(c, c)G'_n(c, j; y_{n,k}(c, j))},$$

and our statements hold. □

We used for the proof of Theorem A.1.1 the technique developed in [159] and the references therein concerning the zeros of a linear combination of two polynomials of the same degree with interlacing zeros. We emphasize an interesting consequence of this theorem. It says that, when  $\lambda$  goes from zero to infinity, each zero  $x_{n,k}(\lambda, c, j)$  runs monotonically over the entire interval  $(y_{n,k}(c, j), x_{n,k})$  or  $(x_{n,k}, y_{n,k}(c, j))$ . In other words, if  $c \leq a$ , then

$$x_{n,k}(\lambda, c, j) \in (y_{n,k}(c, j) = \sup_{\lambda} \{x_{n,k}(\lambda, c, j)\}, x_{n,k} = \inf_{\lambda} \{x_{n,k}(\lambda, c, j)\})$$

## A. DISCRETE SOBOLEV ORTHOGONAL POLYNOMIALS

---

and, if  $c \geq a$ , then

$$x_{n,k}(\lambda, c, j) \in (x_{n,k} = \inf_{\lambda} \{x_{n,k}(\lambda, c, j)\}, y_{n,k}(c, j) = \sup_{\lambda} \{x_{n,k}(\lambda, c, j)\}).$$

Meijer [140] obtained the inequalities (A.2) and (A.3) for  $j = 1$  using another technique. Furthermore, for  $n \geq 3$  he showed that for some choice of  $c$  in  $(a, b)$ ,  $Q_n(\lambda, c, 1; \cdot)$  has two complex zeros, if  $\lambda$  is sufficiently large. Earlier Marcellán, Pérez, and Piñar [129] had shown the interlacing properties (A.2) and (A.3) when  $j = 1$ . Moreover, the monotonicity of  $x_{n,k}(\lambda, c, 1)$  with respect to  $\lambda$ , and the convergence of one of the extreme zeros to  $c$  when  $n$  goes to  $+\infty$  were established in [129].

In the sequel we analyze the location of the smallest (resp. greatest) zero of  $Q_n(\lambda, c, j; \cdot)$  with respect to the point  $a$  (resp.  $b$ ).

**Corollary A.1.1.** *Let  $n > j$  and  $\lambda > 0$ .*

*i) If  $c \leq a$  and  $y_{n,1}(c, j) < a$ , then the smallest zero  $x_{n,1}(\lambda, c, j)$  satisfies*

$$\begin{aligned} x_{n,1}(\lambda, c, j) &> a, & \lambda < \lambda_0, \\ x_{n,1}(\lambda, c, j) &= a, & \lambda = \lambda_0, \\ x_{n,1}(\lambda, c, j) &< a, & \lambda > \lambda_0, \end{aligned}$$

where

$$\lambda_0 = \lambda_0(n, a, c, j) = \frac{P_n(a)}{K_{n-1}^{(j,0)}(c, a)P_n^{(j)}(c) - K_{n-1}^{(j,j)}(c, c)P_n(a)}.$$

*ii) If  $c \geq b$  and  $y_{n,n}(c, j) > b$ , then the largest zero  $x_{n,n}(\lambda, c, j)$  satisfies*

$$\begin{aligned} x_{n,n}(\lambda, c, j) &< b, & \lambda < \lambda_0, \\ x_{n,n}(\lambda, c, j) &= b, & \lambda = \lambda_0, \\ x_{n,n}(\lambda, c, j) &> b, & \lambda > \lambda_0, \end{aligned}$$

where  $\lambda_0 = \lambda_0(n, b, c, j)$ .

The proofs are an immediate consequence of determining the value of the polynomial  $Q_n(\lambda, c, j; \cdot)$  via (A.7), together with the fact that  $G_n(c, j; a)P_n(a) < 0$  if  $y_{n,1}(c, j) < a$  or  $G_n(c, j; b)P_n(b) < 0$  if  $y_{n,n}(c, j) > b$ . Observe that for  $c \notin [a, b]$  we derive explicitly the value  $\lambda_0$  of the mass, such that for  $\lambda > \lambda_0$  one of the zeros is located outside  $(a, b)$ .

A similar result about the mutual location of  $c$  and the extreme zeros of  $Q_n(\lambda, c, j; \cdot)$  is the following corollary.

**Corollary A.1.2.** *Let  $n > j$  and  $\lambda > 0$ .*

---

i) If  $c \leq a$  and  $y_{n,1}(c, j) < c$ , then the smallest zero  $x_{n,1}(\lambda, c, j)$  satisfies

$$\begin{aligned} x_{n,1}(\lambda, c, j) &> c, & \lambda < \lambda_1, \\ x_{n,1}(\lambda, c, j) &= c, & \lambda = \lambda_1, \\ x_{n,1}(\lambda, c, j) &< c, & \lambda > \lambda_1, \end{aligned}$$

$$\text{where } \lambda_1 = \lambda_1(n, c, j) = \frac{P_n(c)}{K_{n-1}^{(j,0)}(c, c)P_n^{(j)}(c) - K_{n-1}^{(j,j)}(c, c)P_n(c)}.$$

ii) If  $c \geq b$  and  $y_{n,n}(c, j) > c$ , then the largest zero  $x_{n,n}(\lambda, c, j)$  satisfies

$$\begin{aligned} x_{n,n}(\lambda, c, j) &< c, & \lambda < \lambda_1, \\ x_{n,n}(\lambda, c, j) &= c, & \lambda = \lambda_1, \\ x_{n,n}(\lambda, c, j) &> c, & \lambda > \lambda_1, \end{aligned}$$

$$\text{where } \lambda_1 = \lambda_1(n, c, j).$$

It means that, depending on the value of the parameter  $\lambda$ , one zero can be located outside the interval  $(\min\{a, c\}, \max\{b, c\})$ .

### A.1.1 Jacobi polynomials

Let  $\{Q_n^{(\alpha, \beta)}(\lambda, c, j; \cdot)\}_{n \geq 0}$  be the polynomials which are orthogonal with respect to the discrete Sobolev inner product (A.1) with  $d\mu = d\mu(\alpha, \beta; \cdot)$ ; see Chapter 1. Let us denote by  $x_{n,k}(\alpha, \beta)$  and  $x_{n,k}(\lambda, c, j; \alpha, \beta)$  the zeros of  $P_n^{(\alpha, \beta)}$  and  $Q_n^{(\alpha, \beta)}(\lambda, c, j; \cdot)$ , respectively.

To illustrate the results obtained in Corollary A.1.1 and A.1.2 we consider two figures. In the Figure A.1(a) we consider  $n = 3$ ,  $\alpha = -1/2$ ,  $\beta = 1/2$ ,  $j = 2$ , and  $c = 1$  for some values of  $\lambda$ . In the Figure A.1(b) we take the same values of  $n$ ,  $\alpha$ ,  $\beta$ , and  $j$  but we choose  $c = 2$ , that is, now  $c$  is not an endpoint of  $(-1, 1)$  and we vary the parameter  $\lambda$ . We see that in both figures at least one zero of

$$Q_3^{(-1/2, 1/2)}(\lambda, c, 2; x) = x^3 - \frac{384c\lambda + \pi}{2(64\lambda + \pi)}x^2 + \frac{192c\lambda - 96\lambda - \pi}{2(64\lambda + \pi)}x + \frac{384c\lambda + \pi}{8(64\lambda + \pi)}$$

is outside of the support  $(-1, 1)$ . Moreover, one zero coincides with  $b = 1$  when  $\lambda = \lambda_0 = \pi/(128(3c - 1))$  and with  $c$  when  $\lambda = \lambda_1 = \pi(1 - 4c - 4c^2 + 8c^3)/(256c^2(4c - 3))$ . In Figure A.1(a) observe that since  $b = c = 1$  then  $\lambda_0 = \lambda_1$ .

We also provide two tables illustrating the monotonicity of the zeros of  $Q_n^{(\alpha, \beta)}(\lambda, c, j; \cdot)$  as functions of  $\lambda$ , and the convergence of these zeros to the zeros of  $G_n^{(\alpha, \beta)}(c, j; \cdot)$ , which are the polynomials defined in (A.4). In Table A.1 we can observe the behavior of the zeros of  $Q_4^{(\alpha, \beta)}(\lambda, c, j; \cdot)$  when  $\alpha = -1/2$ ,  $\beta = 1/2$ ,  $j = 2$  and  $c = 1$ , for some values of

## A. DISCRETE SOBOLEV ORTHOGONAL POLYNOMIALS




Figure A.1: Location of the smallest and greatest zero of  $Q_n(\lambda, c, j; x)$

$\lambda$ . It is quite clear that the zeros are increasing functions of  $\lambda$ , and they converge to the zeros of  $G_4^{(-1/2, 1/2)}(1, 2; \cdot)$ , which are  $-0.696751$ ,  $0.0803371$ ,  $0.859082$ , and  $1.69964$ .

Table A.1: Zeros of  $Q_4^{(-1/2, 1/2)}(\lambda, 1, 2; x) = x^4 - (1/2 + 2400\lambda/(\pi + 1664\lambda))x^3 + [-3/4 + 960\lambda/(\pi + 1664\lambda)]x^2 + [1/4 + 1320\lambda/(\pi + 1664\lambda)]x + 1/16 - 240\lambda/(\pi + 1664\lambda)$  for some values of  $\lambda$

$\lambda$	$x_{4,1}(\lambda, 1, 2; -1/2, 1/2)$	$x_{4,2}(\lambda, 1, 2; -1/2, 1/2)$
1/3000	-0.74365	-0.09607
1/2000	-0.73703	-0.07079
$\lambda_0 = \lambda_1 = \pi/4096$	-0.72955	-0.04152
1	-0.69680	0.08019
10	-0.69676	0.08032
100	-0.69675	0.08034
$\lambda$	$x_{4,3}(\lambda, 1, 2; -1/2, 1/2)$	$x_{4,4}(\lambda, 1, 2; -1/2, 1/2)$
1/3000	0.59496	0.96120
1/2000	0.63516	0.97465
$\lambda_0 = \lambda_1 = \pi/4096$	0.68774	1
1	0.85899	1.69721
10	0.85907	1.69940
100	0.85908	1.69962

In Table A.2 we present the zeros of  $Q_4^{(\alpha, \beta)}(\lambda, c, j; \cdot)$  when  $\alpha = -1/2$ ,  $\beta = 1/2$ ,  $j = 2$ ,  $c = 2$  for several choices of  $\lambda$ . In this table, since the zeros of  $G_4^{(-1/2, 1/2)}(2, 2; \cdot)$  are  $-0.659576$ ,  $0.160050$ ,  $0.886989$ , and  $3.76418$ , we observe that the zeros have the same monotonic and asymptotic behavior when the point  $c$  does not coincide with an endpoint of the interval  $(-1, 1)$ .



Table A.2: Zeros of  $Q_4^{(-1/2,1/2)}(\lambda, 2, 2; x) = x^4 - (1/2 + 28512\lambda/(\pi + 7808\lambda))x^3 + (-3/4 + 12960\lambda/(\pi + 7808\lambda))x^2 + (1/4 + 14904\lambda/(\pi + 7808\lambda))x + 1/16 - 3240\lambda/(\pi + 7808\lambda)$  for some values of  $\lambda$

$\lambda$	$x_{4,1}(\lambda, 2, 2; -1/2, 1/2)$	$x_{4,2}(\lambda, 2, 2; -1/2, 1/2)$
$\lambda_0 = \pi/54400$	-0.71792	-0.02081
1/3000	-0.67724	0.11537
$\lambda_1 = 17\pi/133376$	-0.67465	0.12258
1	-0.65958	0.16003
10	-0.65958	0.16005
100	-0.65958	0.16005
$\lambda$	$x_{4,3}(\lambda, 2, 2; -1/2, 1/2)$	$x_{4,4}(\lambda, 2, 2; -1/2, 1/2)$
$\lambda_0 = \pi/54400$	0.69706	1
1/3000	0.87010	1.84630
$\lambda_1 = 17\pi/133376$	0.87350	2
1	0.88699	3.76274
10	0.88699	3.76403
100	0.88699	3.76416

## A.1.2 Laguerre polynomials

Let  $x_{n,k}(\lambda, c, j; \alpha)$  be the zeros of the polynomials  $\{Q_n^{(\alpha)}(\lambda, c, j; \cdot)\}_{n \geq 0}$  which are orthogonal with respect to the discrete Sobolev inner product (A.1) with  $d\mu = d\mu(\alpha; \cdot)$ ; see Chapter 1.

To illustrate the behavior of  $x_{n,k}(\lambda, c, j; \alpha)$  we show two figures and two tables. We plot  $Q_3^{(\alpha)}(\lambda, c, j; x)$  for  $\alpha, c$ , and  $j$  fixed and vary  $\lambda$ , to show that, depending on the value of  $\lambda$ , the smallest zero can be less than or equal to  $c$ . In Figure A.2(a) we choose  $\alpha = 1$ ,  $j = 2$ , and  $c = 0$ , that is,  $c$  is an endpoint of the orthogonality interval. In Figure A.2(b) we choose  $\alpha = 1$ ,  $j = 2$ , and  $c = -2$ .

In order to illustrate the behavior of the zeros of the polynomials  $Q_4^{(\alpha)}(\lambda, c, j; \cdot)$  presented in Theorem A.1.1, we present some numerical computations in Tables A.3 and A.4. In the first one we show the zeros  $x_{4,k}(\lambda, c, j; \alpha)$ , with  $\alpha = 1$ ,  $c = 0$  and  $j = 2$  for several choices of  $\lambda$ . Observe that, since the point  $c$  is on the left side of the interval of orthogonality, all the zeros of these polynomials are decreasing functions of  $\lambda$ , and they converge to the zeros of  $G_4^{(1)}(0, 2; \cdot)$ , defined in (A.4), which are  $-3.63913$ ,  $1.16543$ ,  $4.00543$ , and  $9.23750$ .

In Table A.4 we choose the same values of  $\alpha$  and  $j$ , but in this case  $c = -2$ . The zeros  $x_{4,k}(\lambda, c, j; \alpha)$  are also monotonic functions and converge to the zeros of  $G_4^{(1)}(-2, 2; \cdot)$ , which are  $-7.76110$ ,  $1.07579$ ,  $3.76689$ , and  $8.77556$ .

## A. DISCRETE SOBOLEV ORTHOGONAL POLYNOMIALS




Figure A.2: Location of the smallest and greatest zero of  $Q_n(\lambda, c, j; x)$

Table A.3: Zeros of  $Q_4^{(1)}(\lambda, 0, 2; x) = x^4 - (20 - 120\lambda/(3 + 13\lambda))x^3 + (360/(3 + 13\lambda))x^2 - (240 - 5040\lambda/(3 + 13\lambda))x + 120 - 3600\lambda/(3 + 13\lambda)$  for some values of  $\lambda$  ( $\lambda_0 = \lambda_1 = 3/17$ )

$\lambda$	$x_{4,1}(\lambda, 0, 2; 1)$	$x_{4,2}(\lambda, 0, 2; 1)$	$x_{4,3}(\lambda, 0, 2; 1)$	$x_{4,4}(\lambda, 0, 2; 1)$
1/100	0.71843	2.49658	5.60038	10.8012
1/10	0.39195	1.89442	4.85101	10.0719
$\lambda_0 = \lambda_1$	0	1.60262	4.57292	9.82446
1	-2.21826	1.22608	4.11868	9.37350
10	-3.46229	1.17104	4.01682	9.25187
100	-3.62101	1.16599	4.00657	9.23895
1000	-3.63731	1.16549	4.00554	9.23765

Table A.4: Zeros of  $Q_4^{(1)}(\lambda, -2, 2; x) = x^4 - (20 - 396\lambda/(3 + 28\lambda))x^3 + (120 - 5016\lambda/(3 + 28\lambda))x^2 - (240 - 15840\lambda/(3 + 28\lambda))x + 120 - 11088\lambda/(3 + 28\lambda)$  for some values of  $\lambda$  ( $\lambda_0 = 15/322$  and  $\lambda_1 = 471/3854$ )

$\lambda$	$x_{4,1}(\lambda, 0, 2; 1)$	$x_{4,2}(\lambda, 0, 2; 1)$	$x_{4,3}(\lambda, 0, 2; 1)$	$x_{4,4}(\lambda, 0, 2; 1)$
1/100	0.660855	2.33685	5.33702	10.4580
$\lambda_0$	0	1.58118	4.50358	9.62953
1/10	-1.47545	1.26126	4.13306	9.25354
$\lambda_1$	-2	1.219141	4.06667	9.17819
10	-7.62158	1.07713	3.77041	8.78110
100	-7.74700	1.07593	3.76724	8.77611
1000	-7.75969	1.07580	3.76693	8.77561

## Appendix B

# Uvarov spectral transformation

In this appendix we deal with a transformation of a quasi-definite functional by the addition of Dirac's linear functionals supported on  $r$  different points located either on the unit circle or on its complement. Consider the quasi-definite linear functional  $\mathcal{L}$  introduced in (2.23), and let  $\mathcal{L}_U$  be the linear functional such that its associated bilinear functional satisfies

$$\langle f, g \rangle_{\mathcal{L}_U} = \langle f, g \rangle_{\mathcal{L}} + \sum_{i=1}^r m_i f(\alpha_i) \overline{g(\alpha_i)}, \quad m_i \in \mathbb{R} \setminus \{0\}, \quad |\alpha_i| = 1.$$

Necessary and sufficient conditions for the regularity of the perturbed linear functional  $\mathcal{L}_U$  are deduced. We also obtain the corresponding linear  $C$ -functions.

### B.1 Mass points on the unit circle

Using an analog method to the one used in [54], we can show

**Theorem B.1.1.** *The following statements are equivalent:*

- i)  $\mathcal{L}_U$  is a quasi-definite linear functional.
- ii) The matrix  $\mathbf{D}_r^{-1} + \mathbb{K}_{n-1}$  is non-singular, and

$$\mathbf{k}_n + \Phi_n^H (\mathbf{D}_r^{-1} + \mathbb{K}_{n-1})^{-1} \Phi_n \neq 0, \quad n \geq 1.$$

Moreover, the sequence of monic polynomials orthogonal with respect to  $\mathcal{L}_U$  is

## B. UVAROV SPECTRAL TRANSFORMATION

---

given by

$$\Psi_n(z) = \Phi_n(z) - \mathbf{K}_{n-1}(z) \left( \mathbf{D}_r^{-1} + \mathbb{K}_{n-1} \right)^{-1} \Phi_n, \quad n \geq 1, \quad (\text{B.1})$$

with

$$\mathbf{K}_{n-1}(z) = [K_{n-1}(z, \alpha_1), K_{n-1}(z, \alpha_2), \dots, K_{n-1}(z, \alpha_r)], \quad \mathbf{D}_r = \text{diag}[\mathbf{m}_1, \mathbf{m}_2, \dots, \mathbf{m}_r],$$

$$\Phi_n = [\Phi_n(\alpha_1), \Phi_n(\alpha_2), \dots, \Phi_n(\alpha_r)]^T,$$

and

$$\mathbb{K}_{n-1} = \begin{bmatrix} K_{n-1}(\alpha_1, \alpha_1) & K_{n-1}(\alpha_1, \alpha_2) & \cdots & K_{n-1}(\alpha_1, \alpha_r) \\ K_{n-1}(\alpha_2, \alpha_1) & K_{n-1}(\alpha_2, \alpha_2) & \cdots & K_{n-1}(\alpha_2, \alpha_r) \\ \vdots & \vdots & \ddots & \vdots \\ K_{n-1}(\alpha_r, \alpha_1) & K_{n-1}(\alpha_r, \alpha_2) & \cdots & K_{n-1}(\alpha_r, \alpha_r) \end{bmatrix}.$$

*Proof.* First, let us assume that  $\mathcal{L}_U$  is a quasi-definite linear functional and denote by  $\{\Psi_n\}_{n \geq 0}$  its corresponding sequence of monic orthogonal polynomials. Thus,

$$\Psi_n(z) = \Phi_n(z) + \sum_{k=0}^{n-1} \lambda_{n,k} \Phi_k(z), \quad \lambda_{n,k} = -\frac{1}{\mathbf{K}_k} \sum_{i=1}^r m_i \Psi_n(\alpha_i) \overline{\Phi_k(\alpha_i)}, \quad n \geq 1.$$

Then, we have

$$\Psi_n(z) = \Phi_n(z) - \sum_{i=1}^r m_i \Psi_n(\alpha_i) K_{n-1}(z, \alpha_i). \quad (\text{B.2})$$

In particular, for  $j = 1, \dots, r$ , we have the following system of  $r$  linear equations and  $r$  unknowns  $\Psi_n(\alpha_j)$ ,  $j = 1, 2, \dots, r$ ,

$$\Psi_n(\alpha_j) = \Phi_n(\alpha_j) - \sum_{i=1}^r m_i \Psi_n(\alpha_i) K_{n-1}(\alpha_j, \alpha_i).$$

Therefore,

$$\begin{bmatrix} 1 + m_1 K_{n-1}(\alpha_1, \alpha_1) & m_2 K_{n-1}(\alpha_1, \alpha_2) & \cdots & m_r K_{n-1}(\alpha_1, \alpha_r) \\ m_1 K_{n-1}(\alpha_2, \alpha_1) & 1 + m_2 K_{n-1}(\alpha_2, \alpha_2) & \cdots & m_r K_{n-1}(\alpha_2, \alpha_r) \\ \vdots & \vdots & \ddots & \vdots \\ m_1 K_{n-1}(\alpha_r, \alpha_1) & m_2 K_{n-1}(\alpha_r, \alpha_2) & \cdots & 1 + m_r K_{n-1}(\alpha_r, \alpha_r) \end{bmatrix} \Psi_n = \Phi_n,$$

where  $\Psi_n = [\Psi_n(\alpha_1), \Psi_n(\alpha_2), \dots, \Psi_n(\alpha_r)]$ . In other words,

$$(\mathbb{K}_{n-1} \mathbf{D}_r + \mathbf{I}_r) \Psi_n = \Phi_n.$$

Since  $\mathcal{L}_U$  is assumed to be quasi-definite, the matrix  $\mathbb{K}_{n-1} \mathbf{D}_r + \mathbf{I}_r$  is non-singular and, therefore, (B.1) follows from (B.2).

On the other hand, assume *ii*) holds. For  $0 \leq k \leq n-1$ , we have

$$\begin{aligned} \langle \Psi_n, \Phi_k \rangle_{\mathcal{L}_U} &= \left\langle \Phi_n(z) - \sum_{i=1}^r m_i \Psi_n(\alpha_i) K_{n-1}(z, \alpha_i), \Phi_k(z) \right\rangle + \sum_{i=1}^r m_i \Psi_n(\alpha_i) \overline{\Phi_k(\alpha_i)} \\ &= - \sum_{i=1}^r m_i \Psi_n(\alpha_i) \langle K_{n-1}(z, \alpha_i), \Phi_k(z) \rangle + \sum_{i=1}^r m_i \Psi_n(\alpha_i) \overline{\Phi_k(\alpha_i)} = 0, \end{aligned}$$

using the reproducing kernel property in the last expression. Furthermore,

$$\begin{aligned} \langle \Psi_n, \Phi_n \rangle_{\mathcal{L}_U} &= \mathbf{k}_n + \sum_{i=1}^r m_i \Psi_n(\alpha_i) \overline{\Phi_n(\alpha_i)} = \mathbf{k}_n + \Phi_n^H \mathbf{D}_r \Psi_n \\ &= \mathbf{k}_n + \Phi_n^H (\mathbb{K}_{n-1} + \mathbf{D}_r^{-1})^{-1} \Phi_n \neq 0, \end{aligned}$$

which proves that  $\{\Psi_n\}_{n \geq 0}$ , defined by (B.1), is the sequence of monic polynomials orthogonal with respect to  $\mathcal{L}_U$ .  $\square$

Notice that for  $r = 1$ , the regularity condition for  $\mathcal{L}_U$  becomes  $1 + m_1 K_{n-1}(\alpha_1, \alpha_1) \neq 0$ ,  $n \geq 0$ , as shown in [54].

**Theorem B.1.2.** *For  $z \in \mathbb{D}$ , the C-function associated with  $\mathcal{L}_U$  is*

$$F_U(z) = F(z) + \sum_{i=1}^N m_i \left( \frac{\alpha_i + z}{\alpha_i - z} \right).$$

*Proof.* Denoting  $\tilde{c}_{-k} = \langle \mathcal{L}_U, z^{-k} \rangle$ , we have

$$F_U(z) = \tilde{c}_0 + 2 \sum_{k=1}^{\infty} \tilde{c}_{-k} z^k = F(z) + \sum_{i=1}^r m_i \left( \frac{\alpha_i + z}{\alpha_i - z} \right),$$

i.e.,  $F_U(z)$  has simple poles at  $z = \alpha_i$ .  $\square$

## B.2 Mass points outside the unit circle

The next step is to consider a perturbation  $\mathcal{L}_D$  such that its corresponding bilinear functional satisfies

$$\langle f, g \rangle_{\mathcal{L}_D} = \langle f, g \rangle_{\mathcal{L}} + \sum_{i=1}^r \left( m_i f(\alpha_i) \bar{g}(\alpha_i^{-1}) + \bar{m}_i f(\bar{\alpha}_i^{-1}) \overline{g(\alpha_i)} \right), \quad m_i \in \mathbf{C} \setminus \{0\}, \quad |\alpha_i| \neq 0, 1. \quad (\text{B.3})$$

By analogy with the previous case, we can obtain the following result.

**Theorem B.2.1.** *The following statements are equivalent:*

i)  $\mathcal{L}_D$  is a quasi-definite linear functional.

ii) The matrix  $\widetilde{\mathbf{D}}_{2r}^{-1} + \widetilde{\mathbf{K}}_{n-1}$  is non-singular, and

$$\mathbf{k}_n + \widetilde{\Phi}_n^H \left( \widetilde{\mathbf{D}}_{2r}^{-1} + \widetilde{\mathbf{K}}_{n-1} \right)^{-1} \widetilde{\Psi}_n \neq 0, \quad n \geq 1. \quad (\text{B.4})$$

Moreover, the corresponding sequence of monic polynomials orthogonal with respect to  $\mathcal{L}_\Omega$  is given by

$$\widetilde{\Psi}_n(z) = \Phi_n(z) - \widetilde{\mathbf{K}}_{n-1}(z) \left( \widetilde{\mathbf{D}}_{2r}^{-1} + \widetilde{\mathbf{K}}_{n-1} \right)^{-1} \widetilde{\Phi}_n, \quad n \geq 1, \quad (\text{B.5})$$

with

$$\begin{aligned} \widetilde{\Phi}_n(\alpha) &= [\Phi_n(\alpha_1), \dots, \Phi_n(\alpha_r), \Phi_n(\bar{\alpha}_1^{-1}), \dots, \Phi_n(\bar{\alpha}_r^{-1})]^T, \quad \widetilde{\mathbf{D}}_{2r} = \text{diag}\{m_1, \dots, m_r, \bar{m}_1, \dots, \bar{m}_r\}, \\ \widetilde{\mathbf{K}}_{n-1} &= [K_{n-1}(z, \alpha_1), \dots, K_{n-1}(z, \alpha_r), K_{n-1}(z, \bar{\alpha}_1^{-1}), \dots, K_{n-1}(z, \bar{\alpha}_r^{-1})], \end{aligned}$$

$$\begin{aligned} \widetilde{\mathbf{K}}_{n-1} &= \left[ \begin{array}{c|c} \mathbb{R}_{n-1}(\alpha, \alpha) & \mathbb{R}_{n-1}(\alpha, \bar{\alpha}^{-1}) \\ \hline \mathbb{R}_{n-1}(\bar{\alpha}^{-1}, \alpha) & \mathbb{R}_{n-1}(\bar{\alpha}^{-1}, \bar{\alpha}^{-1}) \end{array} \right], \\ \mathbb{R}_{n-1}(\alpha, \alpha) &= \begin{bmatrix} K_{n-1}(\alpha_1, \alpha_1) & \cdots & K_{n-1}(\alpha_1, \alpha_r) \\ \vdots & \ddots & \vdots \\ K_{n-1}(\alpha_r, \alpha_1) & \cdots & K_{n-1}(\alpha_r, \alpha_r) \end{bmatrix}. \end{aligned}$$

Proceeding as in the proof of Theorem B.1.2, we obtain

---

**Theorem B.2.2.** For  $z \in \mathbb{D}$ , the  $C$ -function associated with  $\mathcal{L}_D$  is

$$F_D(z) = F(z) + \sum_{i=1}^r \left( m_i \frac{\alpha_i + z}{\alpha_i - z} + \bar{m}_i \frac{\bar{\alpha}_i^{-1} + z}{\bar{\alpha}_i^{-1} - z} \right),$$

i.e.,  $F_D(z)$  has simple poles at  $z = \alpha_i$  and  $z = \bar{\alpha}_i^{-1}$ .

## **B. UVAROV SPECTRAL TRANSFORMATION**

---



# Bibliography

- [1] M. J. ABLOWITZ AND J. F. LADIK. Nonlinear differential-difference equations. *J. Math. Phys.*, **16**:598–603, 1975.
- [2] M. J. ABLOWITZ AND J. F. LADIK. Nonlinear differential-difference equations and Fourier analysis. *J. Math. Phys.*, **17**:1011–1018, 1976.
- [3] N. I. AKHIEZER. *The Classical Moment Problem and Some Related Questions in Analysis*. Hafner, New York, 1965. Russian original, 1961.
- [4] M. ALFARO, G. LÓPEZ, AND M. L. REZOLA. Some properties of zeros of Sobolev-type orthogonal polynomials. *J. Comput. Appl. Math.*, **69**:171–179, 1996.
- [5] M. ALFARO, F. MARCELLÁN, M. L. REZOLA, AND A. RONVEAUX. On orthogonal polynomials of Sobolev type: Algebraic properties and zeros. *SIAM J. Math. Anal.*, **23**:737–757, 1992.
- [6] M. ALFARO, J. J. MORENO-BALCÁZAR, A. PEÑA, AND M. L. REZOLA. A new approach to the asymptotics for Sobolev orthogonal polynomials. *J. Approx. Theory*, **163**:460–480, 2010.
- [7] R. ÁLVAREZ-NODARSE, F. MARCELLÁN, AND J. PETRONILHO. WKB approximation and Krall-type orthogonal polynomials. *Acta Appl. Math.*, **54**:27–58, 1998.
- [8] R. ÁLVAREZ-NODARSE AND J. PETRONILHO. On Krall-type discrete polynomials. *J. Math. Anal. Appl.*, **10**:335–347, 2004.
- [9] G. S. AMMAR AND W. B. GRAGG. Schur flows for orthogonal Hessenberg matrices. Hamiltonian and gradient flows, algorithms and control. *Algorithms and Control, Fields Inst. Commun.*, **3**:27–34, 1994.

## BIBLIOGRAPHY

---

- [10] J. ARVESÚ, R. ÁLVAREZ-NODARSE, AND F. MARCELLÁN. Modifications of quasi-definite linear functionals via addition of delta and derivatives of delta Dirac functions. *Indag. Math. (N.S.)*, **15**[1]:1–20, 2004.
- [11] R. ASKEY. *Orthogonal Polynomials and Special Functions*. SIAM, Philadelphia, PA, 1975. Regional Conference Series in Applied Mathematics, Vol. 21.
- [12] S. BARNETT. *Polynomials and Linear Control Systems*. Marcel Dekker, New York, 1983.
- [13] G. BAXTER. Polynomials defined by difference system. *J. Math. Anal. Appl.*, **2**:223–263, 1961.
- [14] S. BELMEHDI AND F. MARCELLÁN. Orthogonal polynomials associated to some modifications of a linear functional. *Appl. Anal.*, **46**:1–24, 1992.
- [15] C. BERG. Fibonacci numbers and orthogonal polynomials. *Arab. J. Math. Sci.*, **17**:75–88, 2011.
- [16] F. BEUKERS. Legendre polynomials in irrationality proofs. *Bull. Austral. Math. Soc.*, **22**:431–438, 1980.
- [17] S. BOCHNER. Über Sturm-Liouvillesche polynomsysteme. *Math. Zeit.*, **29**:730–736, 1929.
- [18] A. BÖTTCHER AND S. M. GRUDSKY. *Matrix Spectral Properties of Banded Toeplitz Matrices*. SIAM, Philadelphia, PA, 2005.
- [19] C. F. BRACCIALI, J. H. McCABE, AND A. SRI RANGA. On a symmetry in strong distributions. *J. Comput. Appl. Math.*, **105**:187–198, 1999.
- [20] A. BRANQUINHO, A. FOULQUIÉ, AND F. MARCELLÁN. Asymptotic behavior of Sobolev-type orthogonal polynomials on a rectifiable Jordan curve or arc. *Constr. Approx.*, **18**:161–182, 2002.
- [21] A. BRANQUINHO, L. GOLINSKII, AND F. MARCELLÁN. Orthogonal polynomials and rational modifications of the Lebesgue measure on the unit circle. *Complex Var. Theory Appl.*, **38**:137–154, 1999.
- [22] C. BREZINSKI. *History of Continued Fractions and Padé Approximants*. Springer-Verlag, New York, 1991. Springer Series in Computational Mathematics, Vol. 12.

- 
- [23] M. G. DE BRUIN. A tool for locating zeros of orthogonal polynomials in Sobolev inner product spaces. *J. Comput. Appl. Math.*, **49**:27–35, 1993.
- [24] A. BULTHEEL AND M. VAN BAREL. *Linear Algebra, Rational Approximation and Orthogonal Polynomials*. Elsevier Science Publ., Amsterdam, 1997. Studies in Computational Mathematics, Vol. 6.
- [25] A. CACHAFEIRO, F. MARCELLÁN, AND C. PÉREZ. Lebesgue perturbation of a quasi-definite Hermitian functional. The positive definite case. *Linear Algebra Appl.*, **369**:235–250, 2003.
- [26] A. CACHAFEIRO, F. MARCELLÁN, AND C. PÉREZ. Orthogonal polynomials with respect to a sum of a arbitrary measure and a Bernstein-Szegő measure. *Adv. Comput. Math.*, **26**:81–104, 2007.
- [27] M. CÁMARA, J. FÁBREGA, M. A. FIOL, AND E. GARRIGA. Some families of orthogonal polynomials of a discrete variable and their applications to graphs and codes. *Electron. J. Combin.*, **16**[1]:R83, 2009.
- [28] M. J. CANTERO. *Polinomios ortogonales sobre la circunferencia unidad. Modificaciones de los parámetros de Schur*. PhD thesis, Departamento de Matemáticas, Universidad de Zaragoza, 1997. In Spanish.
- [29] M. J. CANTERO, L. MORAL, AND L. VELÁZQUEZ. Five-diagonal matrices and zeros of orthogonal polynomials on the unit circle. *Linear Algebra Appl.*, **362**:29–56, 2003.
- [30] M. J. CANTERO, L. MORAL, AND L. VELÁZQUEZ. Direct and inverse polynomial perturbations of hermitian linear functionals. *J. Approx. Theory*, **163**:988–1028, 2011.
- [31] C. CARATHÉODORY. Über den variabilitätsbereich der Koeffizienten von Potenzreihen die gegebene Werter nicht annehmen. *Math. Ann.*, **64**:95–115, 1907.
- [32] K. CASTILLO. A new approach to relative asymptotic behavior for discrete Sobolev-type orthogonal polynomials on the unit circle. *Appl. Math. Lett.*, **25**:1000–1004, 2012f.
- [33] K. CASTILLO, D. K. DIMITROV, L. GARZA, AND F. R. RAFAELI. Perturbations on the antidiagonals of Hankel matrices. 2012h. Submitted.

## BIBLIOGRAPHY

---

- [34] K. CASTILLO, L. GARZA, AND F. MARCELLÁN. On a direct Uvarov-Chihara problem and some extensions. Submitted.
- [35] K. CASTILLO, L. GARZA, AND F. MARCELLÁN. Laurent polynomials perturbations of linear functional. An inverse problem. *Electron. Trans. Numer. Anal.*, **36**:83–98, 2010.
- [36] K. CASTILLO, L. GARZA, AND F. MARCELLÁN. Linear spectral transformations, Hessenberg matrices, and orthogonal polynomials. *Rend. Circ. Mat. Palermo*, **82**[2]:3–26, 2010.
- [37] K. CASTILLO, L. GARZA, AND F. MARCELLÁN. A new linear spectral transformation associated with derivatives of Dirac linear functionals. *J. Approx. Theory*, **163**:1834–1853, 2011.
- [38] K. CASTILLO, L. GARZA, AND F. MARCELLÁN. Perturbations on the subdiagonals of Toeplitz matrices. *Linear Algebra Appl.*, **434**:1563–1579, 2011.
- [39] K. CASTILLO, L. GARZA, AND F. MARCELLÁN. Asymptotic behavior of Sobolev orthogonal polynomials on the unit circle. *Integral Transforms Spec. Funct.*, 2012a. DOI: 10.1080/10652469.2011.649751.
- [40] K. CASTILLO, L. GARZA, AND F. MARCELLÁN. Zeros of discrete Sobolev orthogonal polynomials on the unit circle. *Numer. Algorithms*, **60**:669–681, 2012b.
- [41] K. CASTILLO, R. L. LAMBLÉM, F. R. RAFAELI, AND A. SRI RANGA. Szegő polynomials and para-orthogonal polynomials on the real line: Zeros and canonical spectral transformations. *Math. Comp.*, **81**:2229–2249, 2012d.
- [42] K. CASTILLO, R. L. LAMBLÉM, AND A. SRI RANGA. On a moment problem associated with Chebyshev polynomials. *Appl. Math. Comput.*, **218**:9571–9574, 2012e.
- [43] K. CASTILLO AND F. MARCELLÁN. Generators of rational spectral transformations for nontrivial  $\mathcal{C}$ -functions. *Math. Comp.*, 2012c. In Press.
- [44] K. CASTILLO, M. V. MELLO, AND F. R. RAFAELI. Monotonicity and asymptotics of zeros of Sobolev type orthogonal polynomials: A general case. *Appl. Numer. Math.*, **62**:1663–1671, 2012g.

- [45] P. L. CHEBYSHEV. Sur les fractions continues. *J. Math. Pures Appl.*, **3**[2]:289–323, 1885. Translation of a Russian paper published in 1885; reprinted in *Oeuvres de P. L. Tchebychef*, tome 1, A. Markoff and N. Sonin (eds.), St Petersburg 203–230, 1899; reprinted Chelsea, New York, 1961.
- [46] T. S. CHIHARA. *An Introduction to Orthogonal Polynomials*. Gordon and Breach, New York-London-Paris, 1978. Mathematics and Its Applications, Vol. 13.
- [47] E. B. CHRISTOFFEL. Über die Gaussische Quadratur und eine Verallgemeinerung derselben. *J. Reine Angew. Math.*, **55**:61–82, 1858.
- [48] A. K. COMMON AND J. H. McCABE. The symmetric strong moment problem. *J. Comput. Appl. Math.*, **67**:327–341, 1996.
- [49] J. B. CONWAY. *Functions of One Complex Variable I*. Springer-Verlag, New York, 1978.
- [50] C. W. CRYER. Über die Jacobischen Polynome und zwei verwandte Polynomklassen. *Math. Zeit.*, **39**:634–638, 1935.
- [51] C. W. CRYER. Rodrigues formula and the classical orthogonal polynomials. *Bull. Un. Mat. Ital.*, **25**:1–11, 1970.
- [52] A. P. DA SILVA AND A. SRI RANGA. Polynomials generated by three term recurrence relation. Bounds for complex zeros. *Linear Algebra Appl.*, **397**:299–324, 2005.
- [53] G. DARBOUX. Mémoire sur l’approximation des fonctions de très-grands nombres, et sur une classe étendue de développements en série. *Liouville J.*, **4**[3]:5–56, 377–416, 1878.
- [54] L. DARUIS, J. HERNÁNDEZ, AND F. MARCELLÁN. Spectral transformations for Hermitian Toeplitz matrices. *J. Comput. Appl. Math.*, **202**:155–176, 2007.
- [55] D. DAY AND L. ROMERO. Roots of polynomials expressed in terms of orthogonal polynomials. *SIAM J. Numer. Anal.*, **43**[5]:1969–1987, 2005.
- [56] L. DE BRANGES. A proof of the Bieberbach conjecture. *Acta Math.*, **154**[1]:137–152, 1985.
- [57] P. A. DEIFT. *Orthogonal Polynomials and Random Matrices: A Riemann-Hilbert Approach*. New York University, Courant Institute of Mathematical Sciences,

## BIBLIOGRAPHY

---

- New York; Amer. Math. Soc., Providence, Rhode Island, 1999. Courant Lecture Notes, Vol. 3.
- [58] P. DELSARTE AND Y. GENIN. The split Levinson algorithm. *IEEE Trans. Acoust. Speech Signal Process.*, **34**:470–478, 1986.
- [59] P. DELSARTE AND Y. GENIN. On the role of orthogonal polynomials on the unit circle in digital signal processing applications. *Orthogonal polynomials: Theory and Practice*, **294**:115–133, 1990. P. Nevai (ed.), Kluwer Academic Publishers, Dordrecht-Boston-London, Serie C: Mathematical and Physical Sciences.
- [60] P. DELSARTE AND Y. GENIN. Tridiagonal approach to the algebraic environment of Toeplitz matrices, Part I: Basic results. *SIAM J. Matrix Anal.*, **12**:220–238, 1991.
- [61] D. K. DIMITROV, F. MARCELLÁN, AND F. R. RAFAELI. Monotonicity of zeros of Laguerre-Sobolev-type orthogonal polynomials. *J. Math. Anal. Appl.*, **368**:80–89, 2010.
- [62] D. K. DIMITROV, M. V. MELLO, AND F. R. RAFAELI. Monotonicity of zeros of Jacobi-Sobolev type orthogonal polynomials. *Appl. Numer. Math.*, **60**:263–276, 2010b.
- [63] S. ELHAY, G. H. GOLUB, AND J. KAUTSKY. Updating and downdating of orthogonal polynomials with data fitting applications. *SIAM J. Matrix Anal. Appl.*, **12**:327–353, 1991.
- [64] J. FAVARD. Sur les polynômes de Tchebycheff. *C. R. Acad. Sci. Paris*, **200**:2052–2055, 1935.
- [65] L. FAYBUSOVICH AND M. GEKHTMAN. On Schur flows. *J. Phys. A: Math. Gen.*, **32**:4671–4680, 1999.
- [66] L. FÉJER. Über trigonometrische Polynome. *J. Reine Angew. Math.*, **146**:53–82, 1915.
- [67] L. FÉJER. Über die Lage der Nullstellen von Polynomen, die aus Minimumforderungen gewisser Art entspringen. *Math. Ann.*, **85**:41–48, 1922.
- [68] E. FERMI, J. PASTA, AND S. ULAM. *Studies of Nonlinear Problems*. University of Chicago Press, 1965. Collected Works of Enrico Fermi.

- [69] H. FLASCHKA. Discrete and periodic illustrations of some aspects of the inverse method. *Dynamical Systems, Theory and Applications, Lecture Notes in Phys.*, **38**:441–466, 1975.
- [70] A. FOULQUIÉ, F. MARCELLÁN, AND K. PAN. Asymptotic behavior of Sobolev-type orthogonal polynomials on the unit circle. *J. Approx. Theory*, **100**:345–363, 1999.
- [71] G. FREUD. *Orthogonal Polynomials*. Akadémiai Kiadó, Pergamon Press, Budapest, 1971.
- [72] L. GARZA. *Transformaciones espectrales, funciones de Carathéodory y polinomios ortogonales en la circunferencia unidad*. PhD thesis, Departamento de Matemáticas, Universidad de Carlos III de Madrid, 2008. In Spanish.
- [73] W. GAUTSCHI. *Orthogonal Polynomials: Computation and Approximation*. Oxford University Press, 2004.
- [74] YA. L. GERONIMUS. On the polynomials orthogonal with respect to a given number sequence. *Zap. Mat. Otdel. Khar'kov. Univers. i NII Mat. i Mehan.*, **17**:3–18, 1940.
- [75] YA. L. GERONIMUS. On the polynomials orthogonal with respect to a given number sequence and a theorem by W. Hahn. *Izv. Akad. Nauk. SSSR*, **4**:215–228, 1940.
- [76] YA. L. GERONIMUS. On some difference equations and corresponding systems of orthogonal polynomials. *Izv. Akad. Nauk SSSR, Ser. Mat.*, **5**:203–210, 1943. In Russian.
- [77] YA. L. GERONIMUS. On polynomials orthogonal on the unit circle, on trigonometric moment problem, and on allied Carathéodory and Schur functions. *Sb. Math.*, **15**:99–130, 1944. In Russian.
- [78] YA. L. GERONIMUS. *Orthogonal Polynomials on the Unit Circle and Their Applications*. Amer. Math. Soc. Translation, 1954. no. 104.
- [79] YA. L. GERONIMUS. *Orthogonal Polynomials: Estimates, Asymptotic Formulas, and Series of Polynomials Orthogonal on the Unit Circle and on An Interval*. Consultants Bureau, New York, 1961.

## BIBLIOGRAPHY

---

- [80] F. GESZTESY AND B. SIMON.  $m$ -functions and inverse spectral analysis for finite and semi-infinite Jacobi matrices. *J. Anal. Math.*, **73**:267–297, 1997.
- [81] E. GODOY AND F. MARCELLÁN. An analogue of the Christoffel formula for polynomial modification of a measure on the unit circle. *Boll. Un. Mat. Ital.*, **5**[A]:1–12, 1991.
- [82] E. GODOY AND F. MARCELLÁN. Orthogonal polynomials and rational modifications of measures. *Canad. J. Math.*, **45**:930–943, 1993.
- [83] E. GODOY, F. MARCELLÁN, L. SALTO, AND A. ZARZO. Perturbations of discrete semiclassical functionals by Dirac masses. *Integral Transforms Spec. Funct.*, **5**[1-2]:19–46, 1997.
- [84] L. B. GOLINSKII. Schur flows and orthogonal polynomials on the unit circle. *Sb. Math.*, **197**[8]:1145–1165, 2007.
- [85] A. A. GONCHAR. On the convergence of Padé approximants for some classes of meromorphic functions. *Math. USSR Sbornik*, **26**:555–575, 1975.
- [86] W. B. GRAGG. Positive definite Toeplitz matrices, the Arnoldi process for isometric operators, and Gaussian quadrature on the unit circle. *J. Comput. Appl. Math.*, **46**:183–198, 1993. Russian original in Numerical Methods of Linear Algebra, pp. 16–32, Moskov. Gos. Uni., Moscow, 1982.
- [87] U. GRENANDER AND G. SZEGÖ. *Toeplitz Forms and Their Applications*. 2nd edition, Chelsea, New York, 1984. 1st edition, Univ. California Press, Berkeley-Los Angeles, 1958.
- [88] H. HAMBURGER. Über eine Erweiterung des Stieltjesschen Momentproblems I–III. *Math. Ann.*, **81**, **82**, **82**:235–319, 120–164, 168–187, 1920, 1921, 1921.
- [89] F. HAUSDORFF. Momentenprobleme für ein endliches Intervall. *Math. Z.*, **16**:220–248, 1923.
- [90] E. HEINE. *Handbuch der Kugelfunctionen, Theorie und Anwendungen*. G. Reimer, Berlin, 1878, 1881. vols 1,2.
- [91] G. HERGLOTZ. Über potenzreihen mit positive, reellem teil im Einheitspreis. *Ber. Ver. Ges. Wiss. Leipzig*, **63**:501–511, 1911.



- [92] R. HORN AND C. JOHNSON. *Matrix Analysis*. Cambridge University Press, Cambridge, 1990.
- [93] M. E. H. ISMAIL. *Classical and Quantum Orthogonal Polynomials in One Variable*. Cambridge University Press, Cambridge, 2009. Encyclopedia in Mathematics and its Applications, Vol. 98.
- [94] C. G. J. JACOBI. Über die Reduction der quadratischen Formen auf die kleinste Anzahl Glieder. *J. Reine Angew. Math.*, **39**:290–292, 1848.
- [95] C. G. J. JACOBI. Untersuchungen über die Differentialgleichung der Hypergeometrische Reihe. *J. Reine Angew. Math.*, **56**:149–165, 1859.
- [96] W. B. JONES, O. NJÅSTAD, AND W. J. THRON. Orthogonal Laurent polynomials and the strong moment problem. *J. Math. Anal. Appl.*, **98**:528–554, 1984.
- [97] W. B. JONES, O. NJÅSTAD, AND W. J. THRON. Continued fractions associated with trigonometric and other strong moment problems. *Constr. Approx.*, **2**:197–211, 1986.
- [98] W. B. JONES, O. NJÅSTAD, AND W. J. THRON. Perron–Carathéodory continued fractions. *Lecture Notes in Math.*, **1237**:188–206, 1987. J. Gilewicz, M. Pindor, and W. Siemaszko (eds.), *Rational Approximation and Its Applications to Mathematics and Physics*, Springer-Verlag, New York.
- [99] W. B. JONES, O. NJÅSTAD, AND W. J. THRON. Moment theory, orthogonal polynomials, quadrature, and continued fractions associated with the unit circle. *Bull. London Math. Soc.*, **21**:113–152, 1989.
- [100] W. B. JONES AND W. J. THRON. *Continued Fractions. Analytic Theory and Applications*. Addison-Wesley, Reading, Massachusetts, 1980. Encyclopedia of Mathematics and Its Applications, Vol. 11.
- [101] W. B. JONES, W. J. THRON, AND H. WAADELAND. A strong Stieltjes moment problem. *Trans. Amer. Math. Soc.*, **261**:503–528, 1980.
- [102] T. KAILATH, A. VIEIRA, AND M. MORF. Inverse Toeplitz operators, innovations, and orthogonal polynomials. *SIAM Rev.*, **20**:106–119, 1978.
- [103] V. A. KALIAGUINE. On asymptotics of  $L^p$  extremal polynomials on a complex curve ( $0 < p < \infty$ ). *J. Approx. Theory*, **74**:226–236, 1993.

## BIBLIOGRAPHY

---

- [104] V. A. KALIAGUINE. A note on the asymptotics of orthogonal polynomials on a complex arc: the case of a measure with a discrete part. *J. Approx. Theory*, **80**:138–145, 1995.
- [105] V. A. KALIAGUINE AND A. A. KONONOVA. On the asymptotic of polynomials orthogonal on a system of curves with respect to a measure with discrete part. *St. Petersburg Math. J.*, **21**:217–230, 2010.
- [106] A. YA. KHINTCHINE. *Continued Fractions*. Noordhoff, Groningen, 1963.
- [107] T. H. KJELDSSEN. The early history of the moment problem. *Historia Math.*, **20**:19–44, 1993.
- [108] D. J. KORTEWEG AND F. DE VRIES. On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. *Philos. Mag.*, **39**:422–443, 1895.
- [109] A. M. KRALL. *Hilbert Space, Boundary Value Problems and Orthogonal Polynomials*. Birkhäuser Verlag, Basel-Boston-Berlin, 2002. Operator Theory: Advances and Applications, Vol. 133.
- [110] H. L. KRALL. On derivatives of orthogonal polynomials. *Bull. Amer. Math. Soc.*, **42**:423–428, 1936.
- [111] H. L. KRALL. On orthogonal polynomials satisfying a certain fourth order differential equation. *Pennsylvania State College Stud.*, **6**:1–24, 1940.
- [112] R. L. LAMBLÉM. *Polinômios tipo Szegő*. PhD thesis, Department of Computer Science and Statistics, State University of São Paulo, 2011. In Portuguese.
- [113] R. L. LAMBLÉM, J. H. McCABE, M. A. PIÑAR, AND A. SRI RANGA. Szegő type polynomials and para-orthogonal polynomials. *J. Math. Anal. Appl.*, **370**:30–41, 2010.
- [114] A. M. LEGENDRE. Recherches sur l’attraction des sphéroïdes homogènes. *Mém. Math. Phys. Prés. à l’Acad. Roy. Sci. (Paris) par divers savants*, **10**:411–434, 1785.
- [115] E. LEVIN AND D. S. LUBINSKY. Universality limits involving orthogonal polynomials on the unit circle. *Comput. Methods Funct. Theory*, **7**:543–561, 2007.

- [116] X. LI AND F. MARCELLÁN. On polynomials orthogonal with respect to Sobolev inner products. *Pacific J. Math.*, **175**:127–146, 1996.
- [117] X. LI AND F. MARCELLÁN. Representation of orthogonal polynomials for modified measures. *Commun. Anal. Theory Contin. Fract.*, **7**:9–22, 1999.
- [118] G. LÓPEZ, F. MARCELLÁN, AND W. VAN ASSCHE. Relative asymptotics for orthogonal polynomials with respect to a discrete Sobolev inner product. *Constr. Approx.*, **11**:107–137, 1995.
- [119] L. LORENTZEN AND H. WAADELAND. *Continued Fractions with Applications*. Elsevier Science Publ., Amsterdam, 1992. Studies in Computational Mathematics, Vol. 3.
- [120] L. LORENTZEN AND H. WAADELAND. *Continued Fractions: Convergence Theory*. Atlantis Studies in Math., Atlantis Press, 2008. Vol. 1.
- [121] A. K. LOUIS AND F. NATTERER. Mathematical problems of computerized tomography. *Proc. IEEE*, **71**:379–390, 1983.
- [122] W. MAGNUS AND S. WINKLER. *Hill's Equation*. Interscience Publishers John Wiley and Sons, New York-London-Sydney, 1966.
- [123] F. MARCELLÁN, J. S. DEHESA, AND A. RONVEAUX. On orthogonal polynomials with perturbed recurrence relations. *J. Comput. Appl. Math.*, **30**:203–212, 1990.
- [124] F. MARCELLÁN AND J. HERNÁNDEZ. Geronimus spectral transforms and measures on the complex plane. *J. Comput. Appl. Math.*, **217**:441–456, 2008.
- [125] F. MARCELLÁN AND P. MARONI. Sur l'adjonction d'une masse de Dirac à une forme régulière et semiclassique. *Ann. Mat. Pura ed Appl.*, **162**[IV]:1–22, 1992.
- [126] F. MARCELLÁN AND L. MORAL. Sobolev-type orthogonal polynomials on the unit circle. *Appl. Math. Comput.*, **128**:107–137, 2002.
- [127] F. MARCELLÁN, F. PEHERSTORFER, AND R. STEINBAUER. Orthogonality properties of linear combinations of orthogonal polynomials, I. *Adv. Comput. Math.*, **5**:281–295, 1996.
- [128] F. MARCELLÁN, F. PEHERSTORFER, AND R. STEINBAUER. Orthogonality properties of linear combinations of orthogonal polynomials, II. *Adv. Comput. Math.*, **7**:401–428, 1997.

## BIBLIOGRAPHY

---

- [129] F. MARCELLÁN, T. E. PÉREZ, AND M. A. PIÑAR. On zeros of Sobolev-type polynomials. *Rend. Mat. Appl.*, **7**:455–473, 1992.
- [130] F. MARCELLÁN AND F. R. RAFAELI. Monotonicity and asymptotic of zeros of Laguerre-Sobolev-type orthogonal polynomials of higher derivatives. *Proc. Amer. Math. Soc.*, **139**:3929–3936, 2011.
- [131] F. MARCELLÁN AND A. RONVEAUX. On a class of polynomials orthogonal with respect to a discrete Sobolev inner product. *Indag. Math. (N. S.)*, **1**:451–464, 1990.
- [132] F. MARCELLÁN AND A. RONVEAUX. Orthogonal polynomials and Sobolev inner products. A bibliography. *Universidad Carlos III de Madrid*, 2012.
- [133] F. MARCELLÁN AND G. SANSIGRE. Orthogonal polynomials on the unit circle: Symmetrization and quadratic decomposition. *J. Approx. Theory*, **65**:109–119, 1991.
- [134] F. MARCELLÁN AND W. VAN ASSCHE. Relative asymptotics for orthogonal polynomials with a Sobolev inner product. *J. Approx. Theory*, **72**:193–209, 1993.
- [135] P. MARONI. Une théorie algébrique des polynômes orthogonaux: Applications aux polynômes orthogonaux semiclassiques. *MACS Ann. Comput. Appl. Math.*, **9**:95–130, 1991.
- [136] A. MARTÍNEZ-FINKELSHTEIN AND B. SIMON. Asymptotics of the  $L^2$  norm of derivatives of OPUC. *J. Approx. Theory*, **163**:747–773, 2011.
- [137] A. MATÉ, P. NEVAI, AND V. TOTIK. Extensions of Szegő’s theory of orthogonal polynomials II. *Constr. Approx.*, **3**:51–72, 1987.
- [138] H. MCKEAN AND P. VAN MOERBEKE. The spectrum of Hill’s equation. *Invent. Math.*, **30**[3]:217–274, 1975.
- [139] R. MCWEENY. *Symmetry: An Introduction to Group Theory and Its Applications*. Dover, New York, 2002.
- [140] H. G. MEIJER. Zero distribution of orthogonal polynomials in a certain discrete sobolev space. *J. Math. Anal. Appl.*, **172**:520–532, 1993.
- [141] G. MEURANT AND G. H. GOLUB. *Matrices, Moments and Quadrature with Applications*. Princeton University Press, Princeton, 2010.

- [142] P. VAN MOERBEKE. The spectrum of Jacobi matrices. *Invent. Math.*, **37**:45–81, 1976.
- [143] J. MOSER. Three integrable hamiltonian systems connected with isospectral deformations. *Adv. Math.*, **16**:197–220, 1975a.
- [144] J. MOSER. Finitely many mass points on the line under the influence of an exponential potential - an integrable system. *Dynamical systems. Theory and Applications, Lecture Note Phys*, **38**:467–497, 1975b.
- [145] A. MUKAIHIRA AND Y. NAKAMURA. Schur flow for orthogonal polynomials on the unit circle and its integrable discretization. *J. Comput. Appl. Math.*, **139**:75–94, 2002.
- [146] I. NENCIU. Lax pairs for Ablowitz-Ladik system via orthogonal polynomials on the unit circle. *Int. Math. Res. Not.*, **11**:647–686, 2005.
- [147] P. NEVAI. *Orthogonal Polynomials*. Amer. Math. Soc. Providence, Rhode Island, 1979. Memoirs of the Amer. Math. Soc., Vol. 213.
- [148] A. F. NIKIFOROV AND V. B. UVAROV. *Special Functions of Mathematical Physics*. Birkhäuser, Boston, 1988.
- [149] E. M. NIKISHIN AND V. N. SOROKIN. *Rational Approximations and Orthogonality*. Transl. Math. Monogr., 1991. Russian original, 1988.
- [150] S. P. NOVIKOV, S. V. MANAKOV, L. P. PITAEVSKII, AND V. E. ZAKHAROV. *Theory of Solitons: The Inverse Scattering Method*. Plenum, New York, 1984. Russian original, 1980.
- [151] F. PEHERSTORFER. Finite perturbations of orthogonal polynomials. *J. Comput. Appl. Math.*, **44**:275–302, 1992.
- [152] F. PEHERSTORFER. A special class of polynomials orthogonal on the unit circle including the associated polynomials. *Constr. Approx.*, **12**:161–185, 1996.
- [153] F. PEHERSTORFER. On Toda lattices and orthogonal polynomials. *J. Comput. Appl. Math.*, **133**:519–534, 2001.
- [154] F. PEHERSTORFER, V. P. SPIRIDONOV, AND A. S. ZHEDANOV. Toda chain, Stieltjes function, and orthogonal polynomials. *Theoret. Math. Phys.*, **151**[1]:505–528, 2007.

## BIBLIOGRAPHY

---

- [155] F. PEHERSTORFER AND R. STEINBAUER. Characterization of orthogonal polynomials with respect to a functional. *J. Comput. Appl. Math.*, **65**:339–355, 1995.
- [156] F. PEHERSTORFER AND P. YUDITSKII. Asymptotic of orthogonal polynomials in the presence of a denumerable set of mass points. *Proc. Amer. Math. Soc.*, **129**:3213–3220, 2001.
- [157] T. E. PÉREZ AND M. PIÑAR. Global properties of zeros for Sobolev-type orthogonal polynomials. *J. Comput. Appl. Math.*, **49**:225–232, 1993.
- [158] V. F. PISARENKO. The retrieval of harmonics from a covariance function. *Geophys. J. Royal Austral. Soc.*, **33**:347–366, 1973.
- [159] F. R. RAFAELI. *Zeros de polinômios ortogonais na recta real*. PhD thesis, Department of Computer Science and Statistics, State University of São Paulo, 2010. In Portuguese.
- [160] E. A. RAKHMANOV. On the asymptotics of the ratio of orthogonal polynomials I, II. *Sb. Math.*, **32**, **46**:199–214, 105–118, 1977, 1983. Russian original in *Mat. Sb.*, 103(145):237–252, 1977; *Mat. Sb.*, 118(160):104–117, 1982.
- [161] A. SRI RANGA, E. X. L. ANDRADE, AND J. McCABE. Some consequence of symmetry in strong distributions. *J. Math. Anal. Appl.*, **193**:158–168, 1995.
- [162] R. RASALA. The Rodrigues formula and polynomial differential operators. *J. Math. Anal. Appl.*, **84**:443–482, 1981.
- [163] M. RAUSSEN AND C. SKAU. Interview with John Milnor. *Notices Amer. Math. Soc.*, **59**[3]:400–408, 2012.
- [164] F. RIESZ. Sur les opérations fonctionnelles linéaires. *C. R. Acad. Sci. Paris*, **149**:974–997, 1909.
- [165] F. RIESZ. Sur certains systèmes singuliers d'équations intégrales. *Ann. École. Norm. Sup.*, **28**[3]:32–62, 1911.
- [166] L. RIESZ. Über ein Problem des Herrn Carathéodory. *J. Reine Angew. Math.*, **146**:83–87, 1915.
- [167] A. M. ROCKET AND P. SZÜSZ. *Continued Fractions*. World Scientific, Singapore, 1992.

## BIBLIOGRAPHY

---

- [168] A. RONVEAUX AND W. VAN ASSCHE. Upward extension of the Jacobi matrix for orthogonal polynomials. *J. Approx. Theory*, **86**:335–357, 1996.
- [169] E. J. ROUTH. On some properties of certain solutions of a differential equation of the second order. *Proc. London Math. Soc.*, **16**:245–262, 1884.
- [170] W. RUDIN. *Real and Complex Analysis*. McGraw-Hill, New York, 1987. 3rd edition.
- [171] E. B. SAFF AND V. TOTIK. What parts of a measure's support attract zeros of the corresponding orthogonal polynomials? *Proc. Amer. Math. Soc.*, **114**:185–190, 1995.
- [172] W. SCHOUTENS. *Stochastic Processes and Orthogonal Polynomials*. Springer-Verlag, New York, 2000.
- [173] I. SCHUR. Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind, I. *J. Reine Angew. Math.*, **147**:205–232, 1917. I. Gohberg (ed.), English translation in I. Schur Methods in Operator Theory and Signal Processing, pp. 31-59, Operator Theory: Advances and Applications, 18, Birkhäuser, Basel, 1986.
- [174] I. SCHUR. Über Potenzreihen, die im Innern des Einheitskreises beschränkt sind, II. *J. Reine Angew. Math.*, **148**:122–145, 1918. I. Gohberg (ed.), English translation in I. Schur Methods in Operator Theory and Signal Processing, pp. 66-88, Operator Theory: Advances and Applications, 18, Birkhäuser, Basel, 1986.
- [175] J. A. SHOHAT AND J. D. TAMARKIN. *The Problem of Moments*. Amer. Math. Soc., New York, 1943.
- [176] B. SIMON. The classical moment problem as a self-adjoint finite difference operator. *Adv. Math.*, **137**:82–203, 1998.
- [177] B. SIMON. Analogs of the  $m$ -function in the theory of orthogonal polynomials on the unit circle. *J. Comput. Appl. Math.*, **171**:411–424, 2004.
- [178] B. SIMON. *Orthogonal Polynomials on the Unit Circle*. Amer. Math. Soc. Providence, Rhode Island, 2005. Amer. Math. Soc. Coll. Publ. Series, Vol. 54, 2 vols.
- [179] B. SIMON. CMV matrices: Five years after. *J. Comput. Appl. Math.*, **208**:120–154, 2007.

## BIBLIOGRAPHY

---

- [180] B. SIMON. *Szegő's Theorem and Its Descendants: Spectral Theory for  $L^2$  Perturbations of Orthogonal Polynomials*. Princeton University Press, Princeton, 2011.
- [181] H. STAHL AND V. TOTIK. *General Orthogonal Polynomials*. Cambridge University Press, Cambridge, 1992.
- [182] T. J. STIELTJES. Recherches sur les fractions continues. *Ann. Frac. Sci. Toulouse*, **8**, 9:J1–J122, A1–A47, 1894, 1895.
- [183] C. SUÁREZ. *Polinomios ortogonales relativos a modificaciones polinomiales de funcionales regulares hermitianos*. PhD thesis, Departamento de Matemática Aplicada, Universidad de Vigo, 1993. In Spanish.
- [184] P. K. SUETIN. Fundamental properties of polynomials orthogonal on a contour. *Russian Math. Surveys*, **21**:35–83, 1966.
- [185] G. SZEGŐ. Beiträge zur Theorie der Toeplitzschen formen. *Math. Z.*, **6**:167–202, 1920.
- [186] G. SZEGŐ. Beiträge zur Theorie der Toeplitzschen Formen, II. *Math. Z.*, **9**:167–190, 1921.
- [187] G. SZEGŐ. Über orthogonale Polynome, die zu einer gegebenen Kurve der komplexen Ebene gehören. *Math. Z.*, **9**:218–270, 1921.
- [188] G. SZEGŐ. *Orthogonal Polynomials*. Amer. Math. Soc. Providence, Rhode Island, 4th Edition, 1975. Amer. Math. Soc. Coll. Publ. Series, Vol. 24.
- [189] C. TÁSIS. *Propiedades de polinomios ortogonales relativos a la circunferencia unidad*. PhD thesis, Departamento de Matemáticas, Universidad de Cantabria, 1989. In Spanish.
- [190] A. V. TEPLYAEV. The pure point spectrum of random orthogonal polynomials on the unit circle. *Soviet Math. Dokl.*, **44**:407–411, 1992. Russian original in Dokl. Akad. Nauk SSSR 320:49-53, 1991.
- [191] M. TODA. *Theory of Nonlinear Lattices*. Springer-Verlag, Berlin, 1989. Springer Series in Solid-State Sciences Vol. 20.



## BIBLIOGRAPHY

---

- [192] R. USMANI. Inversion of a triangular Jacobi matrix. *Linear Algebra Appl.*, **212/213**:413–414, 1994.
- [193] S. VERBLUNSKY. On positive harmonic functions: A contribution to the algebra of Fourier series. *Proc. London Math. Soc.*, **38**[2]:125–157, 1935.
- [194] L. VINET AND A. ZHEDANOV. Szegő polynomials on the real axis. *Integral Transforms Spec. Funct.*, **8**:149–164, 1999.
- [195] L. VOLKOVYSKI, G. LUNTS, AND I. ARAMANOVICH. *Problems in the Theory of Functions of a Complex Variable*. Mir, Moscow, 1972.
- [196] H. S. WALL. *Analytic Theory of Continued Fractions*. Van Nostrand, New York, 1948. The University Series in Higher Mathematics, Vol. 1.
- [197] J. WALLIS. *The Arithmetic of Infinitesimals*. Springer-Verlag, New York-Oxford, 1656. Translation and introduction by J. A. Stedall of *Arithmetica Infinitorum*, 2004.
- [198] D.S. WATKINS. Some perspectives on the eigenvalue problem. *SIAM Rev.*, **35**:430–471, 1993.
- [199] M. S. WEBSTER. Orthogonal polynomials with orthogonal derivatives. *Bull. Amer. Math. Soc.*, **44**:880–888, 1938.
- [200] B. WENDROFF. On orthogonal polynomials. *Proc. Amer. Math. Soc.*, **12**:554–555, 1961.
- [201] M. L. WONG. Generalized bounded variation and inserting point masses. *Constr. Approx.*, **30**:1–15, 2009.
- [202] A. ZHEDANOV. Rational spectral transformations and orthogonal polynomials. *J. Comput. Appl. Math.*, **85**:67–86, 1997.



